

## Use of Raman spectroscopy to characterize and distinguish minerals of the alunite supergroup

Nicolas Maubec, Abdeltif Lahfid, Catherine Lerouge, Guillaume Wille, Karine Michel

► **To cite this version:**

Nicolas Maubec, Abdeltif Lahfid, Catherine Lerouge, Guillaume Wille, Karine Michel. Use of Raman spectroscopy to characterize and distinguish minerals of the alunite supergroup. Geo-Raman Xth Meeting, Jun 2012, Nancy, France. pp.Mineralogy and Gemmology. hal-00694193

**HAL Id: hal-00694193**

**<https://hal-brgm.archives-ouvertes.fr/hal-00694193>**

Submitted on 3 May 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Use of Raman spectroscopy to characterize and distinguish minerals of the alunite supergroup

Maubec, Nicolas\*, Lahfid, Abdeltif\*, Lerouge, Catherine\*, Wille, Guillaume\*, Michel, Karine\*

\* BRGM, 3 Avenue Claude Guillemin, BP 36009, 45060 Orléans cédex 2, France

The alunite supergroup consists of more than 50 representative minerals with the general formula  $A B_3 (XO_4)_2 (OH)_6$ , where A may be occupied by Na, K, Ca, Pb, Ba, Sr, Ce, La, LREE, U ..., B is the trivalent cation,  $Al^{3+}$  or  $Fe^{3+}$ , but also  $Cu^{2+}$  or  $Zn^{2+}$  and X is the tetrahedral anion, S, P or As [1, 2]. The minerals of this supergroup form numerous complex solid solutions which crystallize in a rhombohedral system, space group  $R\bar{3}m$ . The general structure can be described as a combination of  $BO_2(OH)_4$  – octahedra and  $XO_4$  – tetrahedra. The ions in A-site are located between the  $BO_2(OH)_4$  – octahedral sheets, in an icosahedral site.

The minerals of the alunite supergroup occur in a wide range of geological environments, including metamorphic, sedimentary, magmatic systems and soils. Their large domain of stability, the complex solid solutions between the numerous end-members and their capacity to incorporate heavy metals such as As, Pb, Ba and radionuclides such as Sr, Th, U in their lattice, makes them minerals of interest and potential tools in mineral exploration and resources, but also as traps for pollutants.

Although Raman spectra of synthetic and natural minerals of the alunite supergroup have already been published (e.g.[3, 4]), the attributions of some bands are still debated. It is the same for the interpretations of the wavenumber variations of some vibrational bands, with the elemental composition, which are sometimes superficial.

In this work, acquisition of new Raman spectra was performed on natural minerals differing from one to another by their composition in A-, B- and X-sites. The minerals used are from advanced argillic zones hosted by volcanic rocks in Bulgaria [5] and are defined as K-alunite ( $KAl_3(SO_4)_2(OH)_6$ ), Na-alunite ( $NaAl_3(SO_4)_2(OH)_6$ ), K-jarosite ( $KFe_3(SO_4)_2(OH)_6$ ) and woodhouseite ( $CaAl_3(SO_4)(PO_4)(OH)_6$ ). Their Raman spectra are presented in figure 1. Comparison of these spectra one to one (K-alunite versus Na-rich alunite for Na-K substitution, K-alunite versus K-jarosite for Al-Fe substitution, and K-Na alunite versus woodhouseite for  $SO_4$ - $PO_4$  substitution) combined with crystalline parameters determined by Sato et al. [6] has brought new insight to the interpretation of the vibrational spectra and band assignments.

Thus, the region below  $650\text{ cm}^{-1}$  is complex because of bands due to various deformation modes of  $XO_4$ , as well as vibrations of B-O and B-OH bonds and framework deformations. The region around  $1000 - 1200\text{ cm}^{-1}$  is mainly due to  $XO_4$  stretching vibrations and above  $3000\text{ cm}^{-1}$ , the bands present are assigned to OH stretching vibrations [4, 7].

The influence of the ions, in A-, B- and X-sites, is highlighted by variations of the position of certain vibrations. The substitutions of  $K^+$  by  $Na^+$ , in A-site, are characterized by wavenumber shifts of the vibrations involving the hydroxyl groups, the B-OH and O-H bonds. The B-OH stretching vibrations occur at  $381$  and  $395\text{ cm}^{-1}$  in the K- and Na-alunite spectra, respectively, whereas two O-H stretching bonds appear at  $3480$  and  $3508\text{ cm}^{-1}$  in K-alunite and at  $3459$  and  $3487\text{ cm}^{-1}$  in Na-alunite. These differences are due to both a modification around the hydroxyl groups, such as the interatomic distances A-OH which are shorter in presence of Na, and a difference in the length of the O-H bonds which is longer in the Na-alunite.

The substitutions of  $Al^{3+}$  by  $Fe^{3+}$ , in B-site, cause changes in the position of almost all bands toward lower wavenumbers. These differences are due to the size of  $Fe^{3+}$  ( $0.65\text{ \AA}$  [8]) relative to that of  $Al^{3+}$

(0.54 Å [8]) which induces significant structural changes including the B-O, B-OH and X-O bond lengths and the interaction strengths around the hydroxyl groups.

The wavenumbers of the vibrations assigned to  $XO_4$  groups decrease significantly with increasing the size of the ion in X-site. Thus, the  $PO_4^{3-}$  vibrations, such as the stretching modes ( $990\text{ cm}^{-1}$ ), occur at lower wavenumbers than  $SO_4^{2-}$  ( $1024\text{ cm}^{-1}$ ), as the result of a decrease of the X-O bond force due to a longer P-O bond lengths than S-O bonds. In addition, the present study allows observing that bands are broader in the woodhouseite mineral due to both a disorder in this minerals induced by the presence of different  $XO_4$  groups and a reduction of symmetry of this group which gives a greater number of bands with adjacent positions.

With literature about the Raman spectroscopic characterization of other minerals from the alunite supergroup, it was also possible to generalize the substitution effects on the wavenumbers of several vibrations with other cations in the different sites and to show that this technique could be a powerful tool to identify and distinguish minerals of this supergroup.




Figure 1 : Raman spectra of natural minerals of the alunite supergroup.

#### References:

- [1] Botinelly T. Review of minerals of alunite-jarosite, beudantite, and plumbogummite groups. *Journal of Research of the Us Geological Survey* 1976;4:213.
- [2] Scott KM. Solid-solution in, and classification of, gossan-derived members of the alunite-jarosite family, northwest Queensland, Australia. *American Mineralogist* 1987;72:178.
- [3] Serna CJ, Cortina CP, Ramos JVG. Infrared and Raman-study of alunite jarosite compounds. *Spectrochimica Acta Part a-Molecular and Biomolecular Spectroscopy* 1986;42:729.
- [4] Frost RL, Wills RA, Weier ML, Martens W, Kloprogge JT. A Raman spectroscopic study of alunites. *Journal of Molecular Structure* 2006;785:123.
- [5] Lerouge C, Kunov A, Flehoc C, Georgieva S, Hikov A, Lescuyer JL, Petrunov R, Velinova N. Constraints of stable isotopes on the origin of alunite from advanced argillic alteration systems in Bulgaria. *Journal of Geochemical Exploration* 2006;90:166.
- [6] Sato E, Nakai I, Miyawaki R, Matsubara S. Crystal structures of alunite family minerals: beaverite, corkite, alunite, natroalunite, jarosite, svanbergite, and woodhouseite. *Neues Jahrbuch Fur Mineralogie-Abhandlungen* 2009;185:313.
- [7] Breiting DK, Kriegelstein R, Bogner A, Schwab RG, Pimpl TH, Mohr J, Schukow H. Vibrational spectra of synthetic minerals of the alunite and crandallite type. *Journal of Molecular Structure* 1997;408:287.
- [8] Shannon RD. Revised effective ionic-radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallographica Section A* 1976;32:751.