

The 2004 Parkfield earthquake (Mw 6) post-seismic surface displacement observed by coupling ERS and ENVISAT InSAR between 2005 and 2010.

Guillaume Bacques, Daniel Raucoules, Marcello de Michele, Hideo Aochi, Frédérique Rolandone

► **To cite this version:**

Guillaume Bacques, Daniel Raucoules, Marcello de Michele, Hideo Aochi, Frédérique Rolandone. The 2004 Parkfield earthquake (Mw 6) post-seismic surface displacement observed by coupling ERS and ENVISAT InSAR between 2005 and 2010.. Wegener 2012, Sep 2012, Strasbourg, France. hal-00729072

HAL Id: hal-00729072

<https://hal-brgm.archives-ouvertes.fr/hal-00729072>

Submitted on 7 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The 2004 Parkfield earthquake (Mw 6) post-seismic surface displacement observed by coupling ERS and ENVISAT InSAR between 2005 and 2010.

Session : Seismotectonics and the Earthquake Cycle

Guillaume Bacques, Daniel Raucoules*, Marcello De Michele*, Hideo Aochi*, Frédérique Rolandone***

**BRGM, Bureau de Recherches Géologiques et Minières, Orléans, France*

***ISTEP, CNRS UMR 7193, Université Pierre et Marie Curie-Paris VI, Paris, France*

We used the ability of InSAR technique to provide precise displacement map in line of sight to characterise the post-seismic deformation following the 28 September 2004 (Mw 6) Parkfield earthquake between 2005 and 2010 (San Andreas Fault, California). Measuring the post-seismic displacement distribution and its spatio-temporal evolution is one of the most important steps to constrain the slip budget upon the fault plain and consequently to improve our knowledge concerning the seismic cycle. The expected 2004 Parkfield event, by its time delay and its magnitude, has been classified as an outsider compare to the previous events and led researchers to reconsider the validity of the seismic cycle at Parkfield (e.g. W. H. Bakun et al, Nature, 2005). Up to now, the analysis of seismic and space-geodetic data covering all the seismic cycle around the 2004 event seconded with numerical models provided new elements such as the co-seismic and post-seismic slip distribution and its evolution upon the fault plain (e.g. S. Barbot et al, Science, 2012). Thanks to its high spatial resolution, the use of InSAR could provide complementary new insight on the shallower part of the fault slip behavior. In this study, we propose to use both ERS and ENVISAT interferograms to increase the temporal data sampling on the Parkfield section of the San Andreas Fault. Firstly, we combined 5 years of available SAR acquisitions including both ERS-2 and Envisat covering the post-seismic period from 2005 to 2010 to provide stacks of unwrapped interferograms that represent the estimation of the mean velocity per year. Secondly, we stacked selected interferogram (both from ERS2 and Envisat) in the attempt to measure the temporal evolution of the surface velocities associated with the post seismic surface displacement. Here we present the first result from our study. The measured detailed space distribution and the decay of the displacement velocity rate during the post-seismic period is used as input for slip budget assessment.