



HAL
open science

Pb-Zn-Cd-Hg multi isotopic characterization of the Loire River Basin, France

Anne-Marie Desaulty, Romain Millot, David Widory, Catherine Guerrot,
Christophe Innocent, Xavier Bourrain, Gideon Bartov, T.M. Johnson

► **To cite this version:**

Anne-Marie Desaulty, Romain Millot, David Widory, Catherine Guerrot, Christophe Innocent, et al..
Pb-Zn-Cd-Hg multi isotopic characterization of the Loire River Basin, France. Goldschmidt 2013,
Aug 2013, Florence, Italy. hal-00812010

HAL Id: hal-00812010

<https://hal-brgm.archives-ouvertes.fr/hal-00812010>

Submitted on 11 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

**Pb-Zn-Cd-Hg multi isotopic
characterization of the Loire River Basin,
France**

A.M. DESAULTY¹, R. MILLOT¹, D. WIDORY^{1,2},
C. GUERROT¹, C. INNOCENT¹, X. BOURRAIN³,
G. BARTOV⁴, T.M. JOHNSON⁴

¹BRGM, Laboratory Division, Orléans, France

(*correspondance : am.desaulty@brgm.fr)

²UQAM/GEOTOP, Earth Atm. Sci. Dept., Montréal, Canada

³Agence de l'Eau Loire Bretagne, Orléans, France

⁴University of Illinois, Department of Geology, Urbana, USA

The Loire River in Central France is approximately 1010 km long and drains an area of 117 800 km². Upstream, the Loire river flows following a south to north direction from the Massif Central down to the city of Orléans, 650 km from its source. The Loire River is one of the main European riverine inputs to the Atlantic Ocean. Over time, its basin has been exposed to numerous sources of anthropogenic metal pollutions, such as metal mining, food industry, agriculture and domestic inputs. The Loire River basin is thus an excellent study site to develop new isotope systematics for tracking anthropogenic sources of metal pollutions.

We have chosen to analyze the isotope composition of cadmium (Cd), zinc (Zn), mercury (Hg) and lead (Pb). These heavy metals have been chosen for their toxicity, even at low levels, to humans. The main goal of this study is to characterize 1) the sources and behavior of these heavy metals in the aquatic environment, and 2) their spatial distribution using a multi-isotope approach. Sources responsible for the release of Pb-Zn-Cd-Hg in the Loire basin were sampled and analysed for their concentrations and corresponding isotope compositions. We also analyzed river waters and suspended solids samples, known to play an important role in the transport of heavy metals through river systems. Biota samples (mussels, oysters), as natural accumulators of metal pollutants, were also analyzed.

All trace elements were analysed in the BRGM laboratories using a Q-ICPMS. Pb-Zn-Cd isotope compositions were measured using a Neptune MC-ICPMS at the BRGM. Hg isotope compositions were measured at the U. of Illinois using a Nu Plasma MC-ICPMS. To analyse Zn and Cd, we carried out a two-steps analytical development: 1) a chromatographic separation, followed by 2) analysis on the MC-ICPMS.

The results showed that, on their own, each of these isotope systematics reveals important information about the geogenic or anthropogenic origin of these metals. Considered together, they are however providing a more integrated understanding of the overall budgets of these pollutants at the scale of the Loire River Basin.