

Delta97/95Mo variation in molybdenites: a link with mineralizing processes ?

Noémie Breillat, Catherine Guerrot, Philippe Négrel, Eric Marcoux

► **To cite this version:**

Noémie Breillat, Catherine Guerrot, Philippe Négrel, Eric Marcoux. Delta97/95Mo variation in molybdenites: a link with mineralizing processes ?. 10th Applied Isotope Geochemistry Conference, Sep 2013, Budapest, Hungary. hal-00835823

HAL Id: hal-00835823

<https://hal-brgm.archives-ouvertes.fr/hal-00835823>

Submitted on 19 Jun 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

$\delta^{97/95}\text{Mo}$ VARIATION IN MOLYBDENITES: A LINK WITH MINERALIZING PROCESSES?

Noemie Breillat^{1,2} – Catherine Guerrot¹ – Philippe Negrel¹ – Eric Marcoux²

¹ BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France, e-mail: n.breillat@brgm.fr

² Université d'Orléans, ISTO, UMR 7327, 45071 Orléans, France,

Molybdenum (Mo) isotopic compositions are usually studied to reconstruct (paleo-) redox conditions in ocean (e.g., Siebert *et al.*, 2003). Some studies also focus on isotopic compositions of Mo in molybdenite in order to decipher a possible genetic link between isotopic variations and mineralizing processes. This is also the aim of our work based on a wide molybdenite databank. We propose here a comparison of $\delta^{97/95}\text{Mo}$ of 101 molybdenite samples from different localities, different types of occurrences and different ages. This databank of 101 molybdenites, presently analyzed, represents various types of mineralization as samples from 5 skarns, 7 porphyry deposits, 9 pegmatites, 5 granites, 2 greisens, 7 perigranitic veins and 9 alpine-type fissure veins have been studied. Molybdenite sample ages vary from 5Ma to 2.7Ga.

The Mo isotopic composition has been determined on molybdenites using a MC-ICP-MS Neptune after *aqua regia* dissolution and adjustment to $[\text{Mo}] = 1\mu\text{g.g}^{-1}$. Mass bias was corrected by using Zr as dopant and standard-sample-standard bracketing. The $\delta^{97/95}\text{Mo}$ ratios have been normalized to NIST 3134. The external reproducibility is 0.07‰ (2 σ).

The overall range of the $\delta^{97/95}\text{Mo}$ ratio in the 101 molybdenite samples varies from -0.58 to 0.89‰. The $\delta^{97/95}\text{Mo}$ ratio is higher for the molybdenites formed in alpine-type fissure veins, greisens and perigranitic veins than for the molybdenites issued from granites, pegmatites, porphyry deposits and skarns.

The crystallization temperature can explain this difference as alpine-type fissure veins, greisens and perigranitic veins crystallize at lower temperatures than the other types of deposits. For some occurrences, the $\delta^{97/95}\text{Mo}$ has been performed on several molybdenite samples and variability at occurrence scale has been observed. For example, in Azegour skarn (Morocco), the $\delta^{97/95}\text{Mo}$ varies from -0.40 to 0.32‰ (n= 29) and in “Ravin de la Ruine” alpine-type fissure veins (France) from -0.08 to 0.77‰ (n=3). As noted by Hannah *et al.* (2007), no correlations appear between $\delta^{97/95}\text{Mo}$ and age of the deposits. Further Re-Os dating, Pb and S isotopic compositions in particular deposits (e.g. Azegour skarn) will help to decipher the origins of these variations.

References

1. C. Siebert, T.F. Nägler, F. von Blanckenburg and J.D. Kramers, “Molybdenum isotope records as a potential new proxy for paleoceanography”, *Earth and Planetary Science Letters*, 211, pp.159-171, 2003.
2. J.L. Hannah, H.J. Stein, M.E. Wieser, J.R. de Laeter and M.D. Varner, “Molybdenum isotope variations in molybdenite: Vapor transport and Rayleigh fractionation of Mo”, *Geology*, 35-8, pp.703-706, 2007.