

**Fluid geochemistry monitoring and fractured zones
characterization in the GRT1 borehole (ECOGEI project,
Rittershoffen, Alsace, France)**

Chrystel Dezayes, Bernard Sanjuan, Frédérick Gal, Catherine Lerouge

► **To cite this version:**

Chrystel Dezayes, Bernard Sanjuan, Frédérick Gal, Catherine Lerouge. Fluid geochemistry monitoring and fractured zones characterization in the GRT1 borehole (ECOGEI project, Rittershoffen, Alsace, France). Deep Geothermal Days, Apr 2014, Paris, France. hal-00934757

HAL Id: hal-00934757

<https://hal-brgm.archives-ouvertes.fr/hal-00934757>

Submitted on 22 Jan 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fluid geochemistry monitoring and fractured zones characterization in the GRT1 borehole (ECOGI project, Rittershoffen, Alsace, France)

C. Dezayes, B. Sanjuan, F. Gal, C. Lerouge

BRGM – 3, avenue Claude Guillemin – BP 36009 – 45060 Orléans Cedex 2.

The GRT1 borehole has been drilled in 2012, in the framework of the ECOGI project. This project, associating Roquette frères, Electricité de Strasbourg and Caisse des Dépôts, aims at producing geothermal steam to dry starch in the Roquette factory located close to the Rhine river. This borehole is located within the Rhine Graben, 6 km south-east to the Soultz site, the first experimental EGS power plant in the world.

According to its Soultz experience and its knowledge of the deep fluid circulation, BRGM has led a research project to acquire new knowledge about the fracture network and the nature of fluid in the deeper part of the Rhine Graben. In this framework, during the drilling, Helium and Radon gas contents were monitored; fluids were sampled and analyzed, also including during the hydraulic tests. The fracture network was characterized based on the borehole image UBI and fracture filling minerals were determined.

The Radon concentration remained at low levels and was correlated with the basement geology rather than the deep circulations. On the contrary, the Helium concentration showed high enrichment in relation with fractures observed on the borehole images, as this was observed during the drilling of the Soultz wells (Sanjuan *et al.*, 2006).

The fluids discharged from the borehole and collected during the drilling and hydraulic tests are probably still perturbed by the drilling mud. However, the chemical analyses show a high salinity and characteristics similar to those of the Soultz deep native geothermal brine. Like in Soultz, these results suggest that the fluid is in equilibrium within deep sedimentary formations at the temperature of about $220 \pm 20^\circ\text{C}$ (Sanjuan *et al.*, 2010).

A structural analysis has been performed based on the observation of borehole images UBI recorded in the deeper part of the basin, from Keuper to Permian and the top of the granitic basement. In the sedimentary part, the orientation of fractures is rather homogeneous, mainly in the rhenan directions. We note that in the Muschelkalk, the density of fractures is high and organized in cluster. Two large fractures have been identified in this Muschelkalk part which have shown fluid lost during the drilling. In the contrary, in the Buntsandstein, the density of fractures is less important and very regularly, with the major part of fractures deepening to the west. It is the same in the upper part of the Permian, which corresponds to the Annweiler sandstones. The lower Permian and the top of granitic basement are intensely fractured. The orientation of fractures is very scattered, comparatively to the sedimentary part, including hercynian directions, with high dip to the west. Several large fractures showing fluid circulation during the drilling have been observed with a sub-meridian direction and a dip weaker than for the meso-fractures. As at Soultz, the fracture network is constituted by meso-fractures organized in cluster and crossed cut by large fractures (Dezayes *et al.*, 2010).

A mineralogical determination of cutting sampling within the high fractured zones and the large fractures shows two stages of micro-fracturing at the grain scale: a cataclase stage observed in all samples with microcrystalline quartz and illite, and a vein filling stage with quartz, carbonate and anhydrite observed only at the top of the granite. These two stages have been observed at Soultz but the barite constitutes the mineralogical of vein filling (Genter & Traineau, 1996) instead of the anhydrite. This difference can be due to a variation of the nature of the alteration fluid or a variation of the physic-chemical conditions of precipitation.

The correlation of the different geochemical, mineralogical and structural analyses permits us to highlight fluid circulations in the borehole. These fluid circulations are controlled by two or three fracture zones showing also paleo-circulation indices.

References :

Dezayes C., Genter A., Valley B. (2010a) - Structure of the low naturally fractured geothermal reservoir at Soultz. *CR Geosciences*, Vol.342, n°7-8, p.517-530.

Genter A. & Traineau H. (1996) – Analysis of macroscopic fractures in the HDR geothermal well EPS-1, Soultz-sous-Forêts, France. *J. Volc. Geoth. Res.*, 72, 121-141.

Sanjuan B., Millot R., Dezayes Ch. and Brach M. (2010) - Main characteristics of the deep geothermal brine (5 km) at Soultz-sous-Forêts (France) determined using geochemical and tracer test data. *CR. Geoscience*, 342, 546-559.

Sanjuan B., Pinault J.L., Rose P., Gérard A., Brach M., Braibant G., Cruzet C., Foucher J.C., Gautier A., Touzelet S. (2006) - Geochemical fluid characteristics and main achievements about tracer tests at Soultz-sous-Forêts (France). BRGM/RP-54776-FR, 64 p., 22 fig., 5 tables, 3 ann.