



HAL
open science

Lithium isotopes in the Loire River Basin, France

Romain Millot, Anne-Marie Desauty

► **To cite this version:**

Romain Millot, Anne-Marie Desauty. Lithium isotopes in the Loire River Basin, France. Goldschmidt Conference 2014, Jun 2014, Sacramento, United States. pp.1. hal-00940153

HAL Id: hal-00940153

<https://brgm.hal.science/hal-00940153>

Submitted on 31 Jan 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

Lithium isotopes in the Loire River Basin, France

ROMAIN MILLOT¹, ANNE-MARIE DESAULTY²

¹ BRGM, Laboratory Division, Orléans, France
(r.millot@brgm.fr)

² BRGM, Laboratory Division, Orléans, France
(am.desaulty@brgm.fr)

Assessing the behaviour of lithium and the distribution of Li isotopes during river weathering is of major importance for studying water/rock interactions at the surface of the Earth. Lithium (⁶Li ~ 7.5% and ⁷Li ~ 92.5%) is a fluid-mobile element and, due to the large relative mass difference between its two stable isotopes, it is subject to significant low temperature mass fractionation which provides key information on the nature of erosion processes.

The Loire River in central France is approximately 1010 km long and drains a surface area of 117 800 km². In the upper basin, the bedrock is old plutonic rock overlain by much younger volcanic rocks. In the intermediate basin, the Loire River drains the sedimentary series of the Paris Basin, mainly carbonate deposits. Then, the lower Loire basin drains pre-Mesozoic basement of the Armorican Massif and its overlying Mesozoic to Cenozoic sedimentary deposits.

In the rivers of the Loire Basin, Li isotopes were measured using the Neptune MC-ICP-MS, ⁷Li/⁶Li ratios were normalized to the L-SVEC standard solution (NIST SRM 8545) following the standard-sample bracketing method.

Lithium concentrations in river waters of the Loire River main stream and the main tributaries span a wide range from 2.0 to 46.5 µg/L, whereas δ⁷Li are comprised between +5.0 and +13.3‰.

There is a clear contrast between the headwaters upstream and rivers located downstream in the lowlands, with a significant decrease of the δ⁷Li with the distance from the source. This feature is observed for both low (summer time) and high water stages (spring time). In addition, one of the major tributaries in the Massif Central (the Allier River) is clearly influenced by inputs from mineralized waters resulting of hydrothermal activities having lower δ⁷Li values. More generally, the Li isotopic composition measured in the rivers of the Loire Basin shows that ⁷Li is enriched in the dissolved load. On going analysis will focus on suspended sediments and the distribution of lithium and its isotopes between dissolved and solid products of erosion at the scale of the whole Loire River Basin.