



**HAL**  
open science

## Assessment of a sequential extraction protocol by examining solution chemistry and mineralogical evolution

Nicolas Maubec, H el ene Pauwels, Herv e No el, Xavier Bourrat

► **To cite this version:**

Nicolas Maubec, H el ene Pauwels, Herv e No el, Xavier Bourrat. Assessment of a sequential extraction protocol by examining solution chemistry and mineralogical evolution. European Geosciences Union General Assembly 2015, Apr 2015, Vienne, Austria. pp. EGU2015 - 5860. hal-01111349

**HAL Id: hal-01111349**

**<https://hal-brgm.archives-ouvertes.fr/hal-01111349>**

Submitted on 30 Jan 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



## Assessment of a sequential extraction protocol by examining solution chemistry and mineralogical evolution

Nicolas Maubec (1), H el ene Pauwels (1), Herv e No el (2), and Xavier Bourrat (1)

(1) BRGM, 3 Avenue Claude Guillemin, 45060 ORLEANS C edex 2, FRANCE, (2) G eo-Hyd, 101 rue Jacques Charles, 45160 OLIVET, FRANCE

Knowledge of the behavior of heavy metals, such as copper and zinc in sediments, is a key factor to improve the management of rivers. The mobility of these metals, which may be harmful to the environment, depends directly on their concentration and speciation, which in turn depend on physico-chemical parameters such as mineralogy of the sediment fraction, pH, redox potential, salinity etc ... (Anderson et al., 2000; Sterckeman et al., 2004; Van Oort et al., 2008).

Several methods based on chemical extractions are currently applied to assess the behavior of heavy metals in soils and sediments. Among them, sequential extraction procedure is widely used in soil and sediment science and provides details about the origin, biological and physicochemical availability, mobilization and transports of trace metals elements. It is based on the use of a series of extracting reagents to extract selectively heavy metals according to their association within the solid phase (Cornu and Clozel, 2000) including the following different fraction: exchangeable, bound to carbonates, associated to oxides (reducible fraction), linked to organic matter and sulfides (oxidizable fraction) as well as silicate minerals so called residual fraction (Hickey and Kittrick, 1984; Tessier et al., 1979). Consequently sequential extraction method is expected to simulate a lot of potential natural and anthropogenic modifications of environmental conditions (Arey et al., 1999; Brannon and Patrick, 1987; Hickey and Kittrick, 1984; La Force et al., 1999; Tessier et al., 1979).

For three decades, a large number of protocols has been proposed, characterized by specific reagents and experimental conditions (concentrations, number of steps, extraction orders and solid/solution ratio) (Das et al., 1995; Gomez Ariza et al., 2000; Quevauviller et al., 1994; Rauret, 1998; Tack and Verloo, 1995), but it appeared that several of them suffer from a lack of selectivity of applied reagents: besides target ones, some of them are able to leach several solid phases.

In this context, the aim of the present study is to investigate the effectiveness and the selectivity of different reagents for metal extraction from target geochemical fraction. It is based on solid analyses with the use of X-ray diffraction and a scanning electron microscopy (SEM) coupled to a microRaman spectrometer in conjunction with chemical analyses of extracting solutions at each step. This methodology provides the opportunity to assess more accurately the effect of each reagent. The study focuses on extraction of Cu and Zn from sediment samples collected at two sites from river banks and characterized by presence of Quartz, Feldspar K, Micas, Kaolinite but with differences regarding accessory phases (pyrite, organic matter, iron oxyhydroxide, calcite). The interaction of the samples with eight different reagents was assessed and compared (Ca(NO<sub>3</sub>)<sub>2</sub> and CaCl<sub>2</sub> for the exchangeable fraction; buffered solutions of sodium acetate/acetic acid at pH = 5.5 and pH = 5 for the acido-soluble fraction; hydroxylamine hydrochloride and a solution of ammonium oxalate/oxalic acid for reducible fraction; hydrogen peroxide and sodium hypochlorite for the oxidizable fraction. In-depth characterization of solid residue at each step allowed proposing the best protocol for both metals.

Anderson, P., Davidson, C. M., Duncan, A. L., Littlejohn, D., Ure, A. M., and Garden, L. M. (2000). Column leaching and sorption experiments to assess the mobility of potentially toxic elements in industrially contaminated land. *Journal of Environmental Monitoring*, 2.

Arey, J. S., Seaman, J. C., and Bertsch, P. M. (1999). Immobilization of uranium in contaminated sediments by hydroxyapatite addition. *Environmental Science & Technology*, 33, 337-342.

Brannon, J. M., and Patrick, W. H. (1987). Fixation, transformation, and mobilization of arsenic in sediments. *Environmental Science & Technology*, 21, 450-459.

Cornu, S., and Clozel, B. (2000). Extractions s equentielles et sp eci ation des  el ements trace m etalliques dans les sols naturels. *Analyse critique*, 7, 179-189.

Das, A. K., Chakraborty, R., Cervera, M. L., and Delaguardia, M. (1995). Metal speciation in solid matrices.

Talanta, 42.

Gomez Ariza, J. L., Giraldez, I., Sanchez-Rodas, D., and Morales, E. (2000). Selectivity assessment of a sequential extraction procedure for metal mobility characterization using model phases. *Talanta*, 52, 545-554.

Hickey, M. G., and Kittrick, J. A. (1984). Chemical partitioning of cadmium, copper, nickel and zinc in soils and sediments containing high-levels of heavy metals. *Journal of Environmental Quality*, 13, 372-376.

La Force, M. J., Fendorf, S., Li, G. C., and Rosenzweig, R. F. (1999). Redistribution of trace elements from contaminated sediments of Lake Coeur d'Alene during oxygenation. *Journal of Environmental Quality*, 28, 1195-1200.

Quevauviller, P., Rauret, G., Muntau, H., Ure, A. M., Rubio, R., Lopez-sanchez, J. F., Fiedler, H. D., and Griepink, B. (1994). Evaluation of a sequential extraction procedure for the determination of extractable trace-metal contents in sediments. *Fresenius Journal of Analytical Chemistry*, 349.

Rauret, G. (1998). Extraction procedures for the determination of heavy metals in contaminated soil and sediment. *Talanta*, 46(3), 449-455.

Sterckeman, T., Douay, F., Baize, D., Fourier, H., Proix, N., and Schwartz, C. (2004). Factors affecting trace element concentrations in soils developed on recent marine deposits from northern France. *Applied Geochemistry*, 19.

Tack, F. M. G., and Verloo, M. G. (1995). Chemical speciation and fractionation in soil and sediment heavy-metal analysis - a review. *International Journal of Environmental Analytical Chemistry*, 59, 225-238.

Tessier, A., Campbell, P. G. C., and Bisson, M. (1979). Sequential extraction procedure for the speciation of particulate trace-metals. *Analytical Chemistry*, 51.

Van Oort, F., Jongmans, A. G., Lamy, I., Baize, D., and Chevallier, P. (2008). Impacts of long-term waste-water irrigation on the development of sandy Luvisols: consequences for metal pollutant distributions. *European Journal of Soil Science*, 59.