



# Spectral induced polarization of Na-montmorillonite dispersions

Philippe Leroy, Guillaume Mériguet, S Huisman, E Zimmermann, M Weigand, C Tournassat, Mohamed Bizi, A. Kemna

## ► To cite this version:

Philippe Leroy, Guillaume Mériguet, S Huisman, E Zimmermann, M Weigand, et al.. Spectral induced polarization of Na-montmorillonite dispersions. Goldschmidt 2016, Jun 2016, Yokohama, Japan. 2016. <hal-01280362>

**HAL Id: hal-01280362**

**<https://hal-brgm.archives-ouvertes.fr/hal-01280362>**

Submitted on 29 Feb 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Spectral induced polarization of Na-montmorillonite dispersions

P. LEROY<sup>1\*</sup>, G. MERIGUET<sup>2</sup>, S. HUISMAN<sup>3</sup>,  
E. ZIMMERMANN<sup>3</sup>, M. WEIGAND<sup>4</sup>, C. TOURNASSAT<sup>1</sup>,  
M. BIZI<sup>1</sup> AND A. KEMNA<sup>4</sup>

<sup>1</sup> BRGM, 45060 Orléans, France

(\*correspondance: p.leroy@brgm.fr)

<sup>2</sup> PHENIX, UPMC, UMR 8234, 75252 Paris, France

(guillaume.meriguet@upmc.fr)

<sup>3</sup> Forschungszentrum, 52425 Jülich, Germany

<sup>4</sup> University of Bonn, 53115 Bonn, Germany

## Introduction

Montmorillonite has remarkable adsorption properties because of its high specific surface area and surface charge density. Its surface charge distribution is also anisotropic. The basal faces have a negative and permanent surface charge due to isomorphic substitutions in the crystal and the edge faces have a pH-dependent surface charge due to amphoteric surface sites. We propose low-frequency complex impedance measurements of Na-Mt dispersions in the unexplored frequency range [Hz kHz] to get information on surface charge distribution, size and shape of Na-Mt dispersions at various particles concentrations (weight fractions 0.5 to 5%), salinities (from  $10^{-4}$  to  $10^{-2}$  M NaCl), and pH (5 to 10).

## Experiment and results

Complex impedance measurements were performed using a cylindrical sample holder of length 30 cm, diameter 1 cm, two brass electrodes at the top and bottom of the sample holder for current injection, two brass electrodes for electric potential measurement, and the spectral induced polarization (SIP) apparatus developed by Zimmermann *et al.* [1]. The maximum instrument accuracy is a phase shift between injected current and measured electrical potential difference of 0.1 mrad. Electrochemical polarization of the electrical double layer around Mt aggregates was observed [2]. Dielectric permittivity spectra were inverted to obtain relaxation time and aggregate size distributions using the inversion code developed by Weigand and Kemna [3]. Depending on particles weight fractions and water chemical composition, changes of the surface electrical properties, size and shape of Mt aggregates were observed.

[1] Zimmermann *et al.* (2008) *Meas. Sci. Technology* **19**, 1-9.

[2] Arroyo *et al.* (2000) *J. Colloid Interface Sci.* **229**, 118-

122. [3] Weigand and Kemna (2016) *Comp. & Geo.* **86**, 34-45.