

Natural gas seeps in the French Alps: Sources and pathways

Wolfram Kloppmann, Michaela Blessing, Eric Proust, Frederick Gal, Gaëtan Bentivegna, Henry Benoît, Pierrick Défossez, Catherine Lerouge, Pauline Humez, Bernhard Mayer, et al.

► **To cite this version:**

Wolfram Kloppmann, Michaela Blessing, Eric Proust, Frederick Gal, Gaëtan Bentivegna, et al.. Natural gas seeps in the French Alps: Sources and pathways. EGU General Assembly 2016, Apr 2016, Vienne, Austria. pp. EGU2016-14484, 2016. <hal-01296648>

HAL Id: hal-01296648

<https://hal-brgm.archives-ouvertes.fr/hal-01296648>

Submitted on 1 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Natural gas seeps in the French Alps: Sources and pathways

Wolfram Kloppmann (1), Michaela Blessing (1), Eric Proust (1), Frédéric Gal (1), Gaetan Bentivegna (1), Benoit Henry (1), Pierrick Defossez (1), Lerouge Catherine (1), Pauline Humez (2), Bernhard Mayer (2), Romain Millot (1), and Eric Gaucher (3)

(1) BRGM, Orléans, France (w.kloppmann@brgm.fr), (2) Department of Geoscience, University of Calgary, Calgary, Alberta, Canada, (3) Total, Pau, France

Natural gas emanations are part of the geochemical baseline to take into account when assessing global greenhouse gas emissions and potential impacts of conventional and unconventional gas exploration and exploitation on groundwater. Examples of such natural gas macro-seeps are known in several parts of the world (Etiope *et al.*, 2009). Only a limited number of them have been characterized for their gas and isotopic compositions. Such analyses can provide essential information for baseline studies, providing insight in the sources (biogenic vs. thermogenic or modified thermogenic) and pathways of such seeps and may allow for distinction of natural seeps from stray gas leakage associated with human activities.

Here, we report gas concentrations and multi-isotope data ($\delta^{13}\text{C}$ and $\delta^2\text{H}$ of methane and ethane, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ of CO_2 , $^3\text{He}/^4\text{He}$ ratio) of two gas seeps in the French subalpine chains, both in a similar geological and structural position within Middle Jurassic claystones along the eastern border of the large synclinal structures of the Vercors and the Chartreuse massifs (Moss, 1992). The “ardent fountain” (*fontaine ardente*) of Le Gua, 30 km south of Grenoble has most likely the longest continuous written record of existence of any individual natural gas seep, mentioned explicitly as early as the first quarter of the 5th century (Augustin of Hippo (St. Augustin), approx. 426) This natural seep was described in the past as a “wet seep” associated with a spring, whereas the second investigated seep, Rochasson near Meylan north of Grenoble, is a dry seep.

Both seeps contain methane and ethane with thermogenic C and H isotope signatures, comparable with a seep in the Northern Swiss Alps at Giswil (Etiope *et al.*, 2010) but with a higher dryness ($\text{C}_1/(\text{C}_2+\text{C}_3) > 1000$) for the Le Gua seep, possibly due to molecular fractionation upon advective fluid+gas migration (Etiope *et al.*, 2009). Maturity (R_0) of the reservoir rocks deduced from $\delta^{13}\text{C}_{(\text{CH}_4)}$, $\delta^{13}\text{C}_{(\text{C}_2\text{H}_6)}$ is similar to values found by Moss (1992) for West-alpine Oxfordian shales. CO_2 contents of $< 9\%$ could stem from methane oxidation. He isotope ratios are radiogenic with only minor mantle contributions, as it is typical for the Western Alps (Marty *et al.*, 1992). This preliminary investigation reveals that thermogenic natural gas can migrate naturally to the surface in significant flux rates over millenaries without anthropogenic causes, in cases that appropriate leakage pathways exist. This study was co-funded by French ANR and Canadian NSERC.

ND_ABSTRACT@

Etiope G., Feyzullayev A., Baciuc C. L. (2009) *Marine and Petroleum Geology*, **26**, 333-344.

Etiope G., Zwahlen C., Anselmetti F. S., Kipfer R., Schubert C. J. (2010) *Geofluids*, **10**, 476-485.

Marty B., Onions R. K., Oxburgh E. R., Martel D., Lombardi S. (1992) *Tectonophysics*, **206**, 71-78.

Moss S. (1992) *Journal of the Geological Society*, **149**, 503-515.

Augustin of Hippo (St. Augustin) (approx. 426) *De Civitate Dei contra paganos*, Vol. Liber XXI, chap. VII