

**Natural CH<sub>4</sub> gas seeps in the French Alps:  
characteristics, typology and contribution to CH<sub>4</sub>  
natural emissions to the atmosphere**

Frédéric Gal, Wolfram Kloppmann, Eric Proust, Gaëtan Bentivegna, Pierrick Défossez, Bernard Mayer, Eric C. Gaucher

► **To cite this version:**

Frédéric Gal, Wolfram Kloppmann, Eric Proust, Gaëtan Bentivegna, Pierrick Défossez, et al.. Natural CH<sub>4</sub> gas seeps in the French Alps: characteristics, typology and contribution to CH<sub>4</sub> natural emissions to the atmosphere. 13th International Conference on Greenhouse Gas Control Technologies, GHGT-13, Nov 2016, Lausanne, Switzerland. hal-01325608

**HAL Id: hal-01325608**

**<https://hal-brgm.archives-ouvertes.fr/hal-01325608>**

Submitted on 2 Jun 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Natural CH<sub>4</sub> gas seeps in the French Alps: characteristics, typology and contribution to CH<sub>4</sub> natural emissions to the atmosphere.

GAL Frédéric<sup>1</sup>, KLOPPMANN Wolfram<sup>1</sup>, PROUST Eric<sup>1</sup>, BENTIVEGNA Gaëtan<sup>1</sup>, DEFOSSEZ Pierrick<sup>1</sup>, MAYER Bernard<sup>2</sup>, GAUCHER Eric C.<sup>3</sup>

<sup>1</sup>BRGM, 3 Avenue Claude Guillemin, 45060 Orléans, France

<sup>2</sup>University of Calgary, Department of Geoscience, 2500 University Dr. NW, Calgary, Alberta, Canada T2N 1N4

<sup>3</sup>TOTAL, CSTJF, Avenue Larribau, 64018 Pau, France

### Abstract

Quantification of greenhouse gases that naturally leak into the atmosphere has to be considered in lien with mitigation of CO<sub>2</sub> emissions, underground storage of CO<sub>2</sub> and conventional and unconventional gas exploitation. Among the long lived greenhouse gases, methane is one of the most critical specie regarding its global warming potential, far greater than the one of CO<sub>2</sub>. Methane can be released in the atmosphere through human activities (agricultural practices, landfills) but it can also be released from natural processes occurring *e.g.* in wetlands. Another source is constituted by natural gas seeps existing all over the world. Their contribution has been quantified in Europe to account for about a third of the yearly methane emissions (Etiope, 2009, Atmospheric Environment 43, 1430-1443).

The global methane budget still needs completion and refining, taking into account undocumented macro-seeps and, in particular, micro-seeps, difficult to measure and to account for. Such unreported seeps are located in the French Alps, near Grenoble. Although some of these emanations are known since Roman times, they have not been investigated since, except in the early XX<sup>th</sup> century for potential industrial exploitation of CH<sub>4</sub>. Here we report first investigations at the *Fontaine Ardente* (Le Gua) and the *Rochasson* (Meylan) natural gas seeps. Investigations were designed to understand the patterns of gas leak and obtain samples for isotope characterisations. The present study is focused on site investigations.

These two gas seeps are set within Middle Jurassic clays formations along the eastern border of the large synclinal structures of the Vercors and the Chartreuse massifs. The *Fontaine Ardente* seep occurs in a small thalweg and has been intensively modified in the past in the aim to protect the main vent from the small stream flowing nearby. The *Rochasson* gas seep has been discovered only recently (40 years ago) and is probably the consequence of a small landslide that occurred in the clayey formations.

Investigations on site include soil gas quantification in the field using portable infrared gas analysers for CO<sub>2</sub>, CH<sub>4</sub> and O<sub>2</sub> concentrations. Subsequent characterisation of the gas phase was done once a day using gas chromatography (N<sub>2</sub>, Ar, helium and alkanes up to C<sub>3</sub>H<sub>8</sub>). Some tentative

evaluations of the CH<sub>4</sub> and CO<sub>2</sub> fluxes were done using a plastic funnel and the record of gas accumulation into this pseudo-chamber.

The gas of *Fontaine Ardente* seep is CH<sub>4</sub>-rich (84%) with significant contribution of CO<sub>2</sub> (11.5%) and minor contribution of air or deep-seated nitrogen (4% N<sub>2</sub> and <1% O<sub>2</sub>). Traces of C<sub>2</sub>H<sub>6</sub> are present (max. 0.06%) together with helium (max. 0.025%). The gas of *Rochasson* seep has nearly the same composition in CH<sub>4</sub>, O<sub>2</sub> and N<sub>2</sub>, but is depleted in CO<sub>2</sub> (<0.5 %), enriched in C<sub>2</sub>H<sub>6</sub> (2%) and in C<sub>3</sub>H<sub>8</sub> (0.33%). Rough evaluation of the gas flux at the mouth of the seeps suggests that fluxes may reach 130 kg/day/m<sup>2</sup> of CH<sub>4</sub> and 38 kg/day/m<sup>2</sup> of CO<sub>2</sub> at *Fontaine Ardente* and 4.3 kg/day/m<sup>2</sup> of CH<sub>4</sub> at *Rochasson*. The value for *Fontaine Ardente* seems to be in reasonable agreement with previous evaluation of the leakage rate established around 130 l/min of CH<sub>4</sub> during borehole drilling in the 90's. As such, this gas seep can be considered as an important one regarding the database established by Etiope (2009).

Even if high fluxes are measured at the mouth of the gas seeps, leakage of CH<sub>4</sub> – and CO<sub>2</sub> – is not spatially restricted to the vent itself. Soil gas investigations allowed defining at very detailed scale (1 m grid) the geometry of gas emanation.

For *Fontaine Ardente*, it is possible to define an area of several square meters (6x4 m) where CH<sub>4</sub> concentrations at 1 m depth exceed 70%. In the centre, CH<sub>4</sub> concentrations are greater than 80%, with high CO<sub>2</sub> (> 10%), the presence of C<sub>2</sub>H<sub>6</sub> and a strong depletion in O<sub>2</sub> and N<sub>2</sub>. The replacement of one gas phase by another is not homogeneous even at such small scale. Samples taken near the gas seep show CH<sub>4</sub>/O<sub>2</sub>, CH<sub>4</sub>/N<sub>2</sub> and CO<sub>2</sub>/O<sub>2</sub> relationships suggesting that O<sub>2</sub> and N<sub>2</sub> replace CH<sub>4</sub> and CO<sub>2</sub>. For each mole of CH<sub>4</sub> introduced in the system, 1/5 mole of O<sub>2</sub> and 4/5 mole of N<sub>2</sub> disappear, balanced by CO<sub>2</sub> (1 mole of CO<sub>2</sub> introduced and 2 moles of O<sub>2</sub> disappeared). With some distance (a few meters) from the seep, the relation is different as CH<sub>4</sub> concentrations remain under 8% while O<sub>2</sub> concentrations decrease from 21% to 8%. Once this pivot point achieved, then the CH<sub>4</sub> concentration rises exponentially. This second mechanism is not yet fully understood – nonetheless, points affected by this process are distributed in a halo shape suggesting that the seep, with high CH<sub>4</sub> flux, sweeps away atmospheric gases, whereas with some distance, there may be some mixing with soil gas. Halo distributions are known around some CO<sub>2</sub> gas vents (Gal *et al.*, 2011, IJGGC 5, 1099-1118).

Situation is different for the *Rochasson* seep. The CH<sub>4</sub> + C<sub>2</sub>H<sub>6</sub> vent is narrower (2x1 m) and associated with strong O<sub>2</sub> and N<sub>2</sub> depletion. High CO<sub>2</sub> concentrations are not measured above the vent but few meters away. This may be the consequence of a CO<sub>2</sub>-O<sub>2</sub> molar replacement without CH<sub>4</sub> influence, a process different from the mechanism found for *Fontaine Ardente*.

These seeps will be investigated in more details, to better quantify fluxes to the atmosphere and to better describe underlying processes. Another challenge will be to evaluate if other unknown macro- or micro-seeps can be discovered in this region, to evaluate if natural degassing of Callovian-Oxfordian shale is restricted to the known point sources or active at wider scale.

Acknowledgments:

This work has been partly funded by the French and Canadian Research Agencies ANR and NSERC (bilateral G-Baseline project).