

## Groundwater management of large aquifers in southwestern France by regional hydrodynamic models

Olivier Douez, Olivier Cabaret, Marc Saltel, Léna Abasq, Arnaud Wuilleumier, Nicolas Pedron

### ► To cite this version:

Olivier Douez, Olivier Cabaret, Marc Saltel, Léna Abasq, Arnaud Wuilleumier, et al.. Groundwater management of large aquifers in southwestern France by regional hydrodynamic models. 43rd IAH International Congress “Groundwater and society: 60 years of IAH”, Sep 2016, Montpellier, France. 2016. <hal-01329626>

HAL Id: hal-01329626

<https://hal-brgm.archives-ouvertes.fr/hal-01329626>

Submitted on 9 Jun 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Groundwater management of large aquifers in southwestern France by regional hydrodynamic models

ABSTRACT n° 1453

Authors :

Name	Email	Country
Douez Olivier	o.douez@brgm.fr	France
Cabaret Olivier	o.cabaret@brgm.fr	France
Saltel Marc	m.saltel@brgm.fr	France
Abasq Léna	l.abasq@brgm.fr	France
Wuilleumier Arnaud	a.wuilleumier@brgm.fr	France
Pedron Nicolas	n.pedron@brgm.fr	France

Topic : Topic 7 - groundwater uses and managing groundwater use conflicts

Sub Topic :

Keywords : groundwater management, hydrodynamic models, conflicts over water

Comments :

Conflict of Interest : 0

Date of submission : 2016-02-23

ABSTRACT [ 327 Words ]

In southwestern France, groundwater resources are withdrawn for various uses (drinkable water, irrigation, industry, geothermal energy, forestry, conservation of wetlands, shellfish breeding among others) which can sometimes cause conflicts between users. To help the management of these resources, regional hydrodynamic models have been developed. These models have been a support to the public policies for over 20 years.

In the northern region, groundwater is in close relation with rivers, especially during periods of low-water levels. Indeed, groundwater used for irrigation impact not only the stream flows, but also the water supply for the second large wetland of France: the Marais Poitevin. The developed models have permitted to test the impact of several scenarios of water withdrawals. These tests help to determinate the amount of water that can be pumped to respect defined objectives as water levels in piezometer and river flows. These models were also used to test the impact of the eventual implementation of water tanks (400,000 to 800,000 m<sup>3</sup>).

Further south, the major problem is the important reduction of the level of the Eocene aquifer in the department of Gironde. In this area, the regional hydrodynamic model has been developed since 1990. Its development allows the management of deep groundwater resources and contributes to validate strategies of exploitation based on different simulations. This model also allows to answer problems of overexploitation and to analyse areas where water savings could be done to avoid this overexploitation, and to estimate the impact of new resources.

The extreme southern region has a significant particularity: on two different sites, groundwater is used for the storage of gas. The cyclic injection and withdrawal of gas impacts significantly the aquifer level. The proposed model is adapted for the groundwater resources knowledge and management of this case.

Finally, these models were used to evaluate impacts of climate change on groundwater resources in order to allow authorities to evaluate strategies to adapt to this change (Project Explore 2070 for example).