

Compaction-related Polygonal Fracture System and mechanical stratigraphy in the Campanian tight chalk of South Landing (Yorkshire, UK)

Cécile Allanic, Nico Hardebol, Ophélie Faÿ-Gomord, Daniel Ammeraal, Michaël Verbiest, Bertrand Gauthier, Giovanni Bertotti, Rudy Swennen, Eric Lasseur, France Champenois

► **To cite this version:**

Cécile Allanic, Nico Hardebol, Ophélie Faÿ-Gomord, Daniel Ammeraal, Michaël Verbiest, et al.. Compaction-related Polygonal Fracture System and mechanical stratigraphy in the Campanian tight chalk of South Landing (Yorkshire, UK). 25ème Réunion des sciences de la Terre (RST 2016), Oct 2016, Caen, France. 2016. <hal-01355575>

HAL Id: hal-01355575

<https://hal-brgm.archives-ouvertes.fr/hal-01355575>

Submitted on 23 Aug 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Compaction-related Polygonal Fracture System and mechanical stratigraphy in the Campanian tight chalk of South Landing (Yorkshire, UK)

Cécile ALLANIC, Nico HARDEBOL, Ophélie FAÏ-GOMORD, Daniel AMMERAAL, Michaël VERBIEST, Bertrand GAUTHIER, Giovanni BERTOTTI, Rudy SWENNEN, Eric LASSEUR, France CHAMPENOIS.

The Santonian-Campanian Flamborough Head chalk cliffs (Yorkshire, UK) are an outstanding outcrop of once deeply-buried tight-chalk (until 1200 m). Flamborough Chalk showed that porosity ranges from 9 to 20% and permeability from 0.04 to 0.2 mD (e.g. Faÿ-Gomord et al., 2016; Sagi et al., 2013). These outcrops are likely to be representative of sub-surface chalk reservoirs in nearby onshore and offshore regions, e.g. the North Sea. The zone had been strongly affected by transpressive tectonic deformations during Tertiary; which has been widely studied (e.g. Bell et al., 1999; Starmer, 2013; Welch et al., 2014), but the precece fracture network, far from these intense deformation zone, has not been investigated yet.

Evidence of synsedimentary faulting associated with polygonal-shaped fractures network was revealed by the observation of platform images (acquired at low altitude and used in photogrammetry). These feature argue for isotropic horizontal state of stress during compaction. Thus, the measured multiple fracture orientations were active simultaneously and mechanically interacting, instead of the intersection of multiple regional sets of different ages reported in the literature. Compaction-related stylolitization occurred in chalk synchronously or soon after synsedimentary deformation, from 800m to 1200m, the maximum burial depth estimated for these outcrops, under the same stress field (sub-vertical σ_1 , and sub-horizontal σ_2 , σ_3). It results in numerous horizontal stylolithic planes and small cemented synsedimentary faults with small displacements often used as preferential pressure-dissolution planes with vertical stylolithic peak.

This fracture pattern, observed in high resolution aerial images and in the field, is also studied using a mechanical stratigraphy approach (e.g. D Jacquemyn et al., 2012). The mechanical units have been characterized using the DigiFract software on high resolution photogrammetry panoramas (Bisdorn et al., 2014). The mechanical interfaces have been defined and their strength quantified. It appears that sedimentological bedding is non-existent or invisible and that the fracture spacing is strongly controlled by the mechanical unit thickness. The mechanical units are mostly delimited by compaction features, such as stylolites and marl-seams, which accumulate insoluble residue, clay essentially. However not all insoluble residue layers behave as mechanical interfaces and the relationship between the thickness of the insoluble residues and the strength of the mechanical interfaces have also been investigated.

Keywords: chalk, synsedimentary faulting, Polygonal Fracture System, photogrammetry mechanical stratigraphy