



**HAL**  
open science

# THE PERCHED PALEOSURFACES OF THE PYRENEES -CANTABRIAN BELT

Robert Wyns, Jessica Uzel

► **To cite this version:**

Robert Wyns, Jessica Uzel. THE PERCHED PALEOSURFACES OF THE PYRENEES - CANTABRIAN BELT. Source to sink: A long term perspective of sediment budgets and sources characterization, Société Géologique de France, Nov 2016, Rennes, France. hal-01366890

**HAL Id: hal-01366890**

**<https://brgm.hal.science/hal-01366890>**

Submitted on 15 Sep 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# THE PERCHED PALEOSURFACES OF THE PYRENEES -CANTABRIAN BELT

WYNS, Robert<sup>1</sup>, UZEL, Jessica<sup>2</sup>

A Source to Sink Total & BRGM collaboration

1: BRGM & ISTO, UMR 7327, BP 36009, 45060 Orléans cedex 2 ([r.wyns@brgm.fr](mailto:r.wyns@brgm.fr))

2: Géosciences Rennes, UMR 6118 – OSUR, 35042 Rennes cedex France

Perched paleosurfaces of the Central and Eastern Pyrenees have been first described by geomorphologists for a long time (Biro, 1937; Lagasquie, 1982; Calvet, 1996) and more recently by geologists (Bosch et al., 2015; Monod et al., 2015; Bosch, in progress). These studies are of peculiar interest since they provide robust constraints for the timing and mechanisms of relief creation in a collision belt.

Pyrenean paleosurfaces have been recognized over large areas in the Axial Zone between the Aure valley to the West and the Cerdagne and Capcir grabens to the East. The actual elevation of these surfaces varies between 1000 and 2500 m above sea level. In Cerdagne, the graben filling has been dated of Vallesian age (Middle to Late Miocene). Based on this constraint, the entire Pyrenean perched paleosurfaces have been considered of Miocene age. Since the palynofacies of the Vallesian sediments is interpreted as a near-sea level deposit, the present elevation of the paleosurfaces should have been acquired after the Vallesian assuming that all paleosurfaces have nearly a Miocene age. In the vicinity of Cerdagne and Capcir grabens, the central Pyrenees paleosurface appears connected to the uppermost terraced paleosurface, suggesting an age prior to the Vallesian times.

A common feature of the Pyrenean paleosurfaces is the preservation of thick lateritic weathering profiles. In the Eastern Pyrenees, the paleosurfaces display a terraced pattern with three or more levels, the highest being likely the oldest. In the Central Pyrenees (from the Aston Massif to the East to the Neouvielle Massif to the West), the single paleosurface is apparently offset along numerous faults. Field work will be conducted in order to collect structural data allowing a detailed tectonic evolution to be reconstructed.

In the Louron valley (close to the Aure valley), the central Pyrenean paleosurface is apparently downthrown along N160 trending faults facing the valley that define a graben-like feature with an overall NS trend. Field work will be conducted to check the structural relationships of such possible features with the reverse faults and thrusts linked to the Pyrenean compression. Preliminary investigations show that fault planes here bear shallow dipping slickensides indicating a probable transtensional mechanism. A second set of normal faults trending N100 to N130, parallel to the belt, seems to offset the central Pyrenean paleosurface. These faults are observed in the Massif of Aston (SW of Ax les Thermes) where a detailed tectonic analysis will be conducted.

Paleosurfaces will be followed from the Aure valley (central Pyrenees) to the East in order to decipher their origin and evolution: are they terraced surfaces or a single surface offset along recent faults having a general N160 and N100-130 orientation? Special attention will be paid to the occurrence of the preserved lateritic weathering profiles that allows reconstructing the lateral correlation of the paleosurface remnants. Finally, we plan to map these paleosurface remnants from the central Pyrenees to the Cantabrian belt with a special attention to possible faults that might offset some portions. An attempt to date the weathering profiles using syn-weathering minerals (adulares) will be conducted in the meantime. These data will collectively help better understanding the significance of paleosurfaces in collision belts.

*This work is part of the project "Source to Sink", we thank Total and BRGM.*