

Spectral induced polarization of Na-montmorillonite dispersions

Philippe Leroy, Maximilian Weigand

► **To cite this version:**

Philippe Leroy, Maximilian Weigand. Spectral induced polarization of Na-montmorillonite dispersions. Clay Conference 2017, Sep 2017, Davos, Switzerland. hal-01528386

HAL Id: hal-01528386

<https://hal-brgm.archives-ouvertes.fr/hal-01528386>

Submitted on 29 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spectral induced polarization of Na-montmorillonite dispersions

Philippe Leroy^{1,2*} and Maximilian Weigand³

¹ BRGM, French Geological Survey, 45060 Orléans, France.

² Department of Earth Sciences, Uppsala University, 75236 Uppsala, Sweden.

³ Department of Geophysics, Steinmann Institute, Bonn University, 53115 Bonn, Germany.

*Corresponding author and mailing address:

Philippe Leroy

BRGM

3 Avenue Claude Guillemin

45060 Orléans Cedex 2, France

E-mail: p.leroy@brgm.fr

Tel: +33 (0)2 38 64 39 73

Abstract

Montmorillonite (Mt) clays have a high specific surface area and surface charge, which confer them remarkable adsorption properties. Nevertheless, their electrochemical and aggregation behavior are not completely elucidated because of the complexity of their microstructural and interfacial properties. In this work, the conductive and dispersive properties of Na-Mt suspensions of weight fractions 0.5 to 5.2% were investigated for the first time using the spectral induced polarization method. A four-electrode system was used to reduce errors introduced by electrode polarization and contact resistances. Complex conductivity spectra in the low-frequency range of 0.1 Hz to 45 kHz were successfully described using a triple layer model of the basal surface of Mt and a complex conductivity model that considers conduction of the diffuse layer and polarization of the Stern layer. Aggregate size distributions were inferred from inverted relaxation time distributions. We found that the negative and permanent surface charge of the basal plane of Na-Mt controls its quadrature (imaginary) conductivity, which is not very sensitive to pH and salinity (NaCl) in the 100 Hz to 45 kHz frequency range. For lower frequencies, the sudden increase of the quadrature conductivity at the highest salinities was explained by considering coagulation of Na-Mt particles.

Keywords: Na-montmorillonite, dispersions, spectral induced polarization, conductivity, dielectric permittivity, basal plane, surface charge, Stern layer, diffuse layer.