

Using structural analysis derived from LIDAR to generate realistic DEM cuttings for rockfall hazard assessment

Clara Levy, Thomas Dewez, Bastien Colas, Didier Hantz

► **To cite this version:**

Clara Levy, Thomas Dewez, Bastien Colas, Didier Hantz. Using structural analysis derived from LIDAR to generate realistic DEM cuttings for rockfall hazard assessment. Journées Aléas Gravitaires (JAG), Oct 2017, Besançon, France. 2017, <www.journeesaleasgravitaires.fr>. <hal-01591404>

HAL Id: hal-01591404

<https://hal-brgm.archives-ouvertes.fr/hal-01591404>

Submitted on 21 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Using structural analysis derived from LIDAR to generate realistic DEM cuttings for rockfall hazard assessment

Clara Levy¹, Thomas Dewez¹, Bastien Colas², Didier Hantz³

1, BRGM, Orléans, France (c.levy@brgm.fr)

2, BRGM, Montpellier, France

3, ISTERre, France

Advances in remote sensing methods such as lidar and photogrammetry have allowed acquiring Digital Elevation Models (DEM) at centimetric to metric resolution. Such topographic detail reveals rock face geological structures to allow the semi-automatic structural analysis of mountain slope (Jaboyedoff et al., 2007; Dewez et al., 2016). It is an important step for the stability analysis of unstable rock compartments. Current techniques include the automatic detection of planar surfaces corresponding to visible discontinuities (joints, fractures, etc.), and analysis of their orientation frequency and extend.

With this realistic description of how the rock massifs are divided into blocks (with attitude and spacing), rockfall risk assessment still needs to focus on slope stability analysis.

The work presented here describes the results of the custom code TOPOCRACKS which divides a high resolution topography by sets of fractures and compute the rupture mechanisms and stability factors of resulting block. It is applied on a 1-m-resolution airborne lidar DEM of a natural hillslope in Ota, Corsica. Field structural observations on site revealed the presence of three fracture sets. So for this toy example, each of the fracture sets was deemed to be 1-m apart. Computation of blocks delimited by fracture intersections generates about 26 000 potentially unstable rock compartments in this area of ca. 600x400 m². As overhangs are very rare in this area, the method was developed in 2.5 D, and the following mechanisms of failure were considered: planar sliding, wedge sliding. The stability of each rock compartment was assessed by performing static analysis. Static analysis takes into account cohesion of joints, friction angles at discontinuities and percentage of rock bridges. Results are compared with expert methodology of rockfall risk assessment at the same study site, and show good agreement for the localization of source areas, the size of the biggest possible event, and the more probable mechanisms of rupture per source areas.

Localisation of potentially unstable rock compartments for a study site in Corsica with a high resolution orthophoto on the background





Figure 1: Result of block partitioning of a 1-m resolution DEM in Corsica based on three fracture sets recognized by field investigations. Considering that each fracture set has a 1-m-period, this figure illustrates how rock blocks could fail by planar sliding on single family fracture or by wedge rupture rooted on different pairs of fracture planes.

References:

Dewez, T.J.B., Girardeau-Montaut, D., Allanic, C., and Rohmer, J. (2016), *FACETS : a CloudCompare plugin to extract geological planes from unstructured 3D point clouds*, *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XLI-B5, 799-804, doi:10.5194/isprs-archives-XLI-B5-799-2016, [link](#).

Jaboyedoff M., Metzger R., Oppikofer T., Couture R., Derron M.-H., Locat J. and Turmel D. (2007) *New insight techniques to analyze rock-slope relief using DEM and 3D-imaging cloud points: COLTOP-3D software*. In: *Rock mechanics: Meeting Society's Challenges and demands* (Vol. 1). Eberhardt E., Stead D. and Morrison T. (Eds.), Taylor & Francis, 61-68, 2007, [DOI:10.1201/NOE0415444019-c8](https://doi.org/10.1201/NOE0415444019-c8)

Acknowledgements:

The project is partly funded by the French national project C2ROP.