

Toxicity of nZVI used for nanoremediation towards natural bacterial community from groundwater

Marc Crampon, Catherine Joulian, Patrick Ollivier, Louis Jolly, Jennifer Hellal

► **To cite this version:**

Marc Crampon, Catherine Joulian, Patrick Ollivier, Louis Jolly, Jennifer Hellal. Toxicity of nZVI used for nanoremediation towards natural bacterial community from groundwater. International Conference on Microbial Ecotoxicology, Ecotoxicomic 2017, Nov 2017, Lyon, France. hal-01621455

HAL Id: hal-01621455

<https://hal-brgm.archives-ouvertes.fr/hal-01621455>

Submitted on 23 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Toxicity of nZVI used for nanoremediation towards natural bacterial community from groundwater

Crampon Marc¹, Jouliau Catherine¹, Ollivier Patrick¹, Jolly Louis¹ and Hellal Jennifer¹,

¹ BRGM, D3E department, team BGE, 3 avenue Claude Guillemin, 45100 Orleans, France.

Remediation techniques for toxic/persistent contaminants in groundwater are often technologically difficult. Nanoparticles (NP) such as nZVI (Zero-Valent Iron) applicable as *in-situ* reduction or oxidation agents for groundwater treatment give promising results. However, they may also represent an additional contamination. Our study aims to assess the impact of nZVI on the denitrifying activity, the abundance and the structure of a planktonic nitrate-reducing bacterial community.

Pre-culture of the denitrifying bacterial community was carried out from groundwater samples. It was then allocated in reactors containing substrate (sodium acetate), nitrates and a range of nZVI concentrations (from 0 to 300 mg Fe.L⁻¹). Physical (pH, redox potential), chemical (NO₃⁻ concentrations) and biological (DNA, RNA) parameters were monitored during one week. The size distribution of nZVI and the mortality of bacteria (Live&Dead observations) were also evaluated. Then the bacterial cultures were transplanted in fresh substrates without nZVI to assess the resilience of the bacterial community.

Denitrification activity was stopped in the presence of nZVI at concentrations higher than 30 mg.L⁻¹, demonstrating the toxic effect of nZVI. Live&Dead observations revealed that the viability of bacteria was lower for lower nZVI concentrations. The presence of NP aggregates was higher for higher nZVI concentrations. They could then result in a lower toxicity for bacteria, even if nZVI concentration is high. DNA and RNA analyses allowed us to observe the impact of nZVI on the activity (*narG* gene), abundance and structure of bacterial community. Finally, resilience was worst for concentrations ranging from 50 to 200 mg.L⁻¹.