



**HAL**  
open science

## Characterization of an agricultural site historically polluted by the destruction of arsenic-containing chemical weapons

Fabienne Battaglia-Brunet, Julie Lions, Catherine Joulian, Nicolas Devau, Mickael Charron, Daniel Hube, Jennifer Hellal, Marina Le Guedard, Isabel Jordan, Prosun Bhattacharya, et al.

### ► To cite this version:

Fabienne Battaglia-Brunet, Julie Lions, Catherine Joulian, Nicolas Devau, Mickael Charron, et al.. Characterization of an agricultural site historically polluted by the destruction of arsenic-containing chemical weapons. The 7th International Congress & Exhibition On Arsenic In The Environment, environmental arsenic in a changing world - AS2018, Jul 2018, Pekin, China. hal-01790840

**HAL Id: hal-01790840**

**<https://hal-brgm.archives-ouvertes.fr/hal-01790840>**

Submitted on 28 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Characterization of an agricultural site historically polluted by the destruction of arsenic-containing chemical weapons

F. Battaglia-Brunet<sup>1</sup>, J. Lions<sup>1</sup>, C. Joulian<sup>1</sup>, N. Devau<sup>1</sup>, M. Charron<sup>1</sup>, D. Hube<sup>1</sup>, J. Hellal<sup>1</sup>, M. Le Guédard<sup>2</sup>, I. Jordan<sup>3</sup>, P. Bhattacharya<sup>4</sup>, K. Loukola-Ruskeeniemi<sup>5</sup>, T. Tarvainen<sup>5</sup> & J. Kaija<sup>5</sup>

<sup>1</sup>BRGM, Orléans, France

<sup>2</sup>LEB Aquitaine Transfert, Villenave d'Ornon, France

<sup>3</sup>GEOS Ingenieurgesellschaft mbH, Halsbrücke, Germany

<sup>4</sup>KTH-Groundwater Arsenic Research Group, Department of Sustainable Development, Environmental Science and Engineering, KTH Royal Institute of Technology, Stockholm, Sweden

<sup>5</sup>Geological Survey of Finland, Espoo, Finland

**ABSTRACT:** Arsenic in agricultural soils may represent a risk for crop quality and surrounding water resources. In the frame of AgriAs project, “Evaluation and management of Arsenic contamination in agricultural soil and water”, a former chemical weapons destruction site converted into agricultural land was characterized. The objective of this study was to identify possible links between arsenic concentration and speciation and bio-indicators informing about the bioavailability of the toxic element. Plants lipidic bio-indicator Omega-3 Index showed that toxicity of the soil was not directly correlated with arsenic concentration. Conversely, arsenic level in the soil samples seemed to influence the density of microbes transforming As species.

### 1 INTRODUCTION

Arsenic (As) and its compounds are toxic and ubiquitous in the environment. It has been estimated that over 220 million people are exposed to As from drinking water or food (Naujokas *et al.*, 2013). Arsenic in agricultural soils and water, and its subsequent entering into the food chain cause potential risk to human health. Indeed, As can accumulate in crop plants such as corn or barley. High As concentrations may also reduce crop yields. Production, destruction and storage of As-rich chemical weapons have increased As concentrations in the European environment. The present study, performed in the framework of the WaterJPI AgriAs project, is focused on the elucidation of the links between As concentration and speciation in soils and a range of bio-indicators that could be helpful to understand the behavior and bio-availability of arsenic in agricultural soils.

### 2 METHODS/EXPERIMENTAL

#### 2.1 Site description and sampling

The sampling site is a former chemical ammunition breaking-down facility of the interwar period converted into agricultural land near Verdun, France. The field is located 25 km north-east from Verdun. The field was used as a pasture in 2002 and was cultivated from 2002 to 2015. Since 2015, it is a fallow ground because cultivation was forbidden when the

pollution was detected. The sampling was performed in May 2017. Many plant species could be observed. The diversity was lower in the most polluted places but plants were observed even in those places. Soils were sampled in a reference zone, far from the polluted area, and along a three-zones transect representing a gradient of As pollution. Soils were sampled in the 0–20 cm layer. Each sample was taken as a composite of 5 points from 3 m × 3 m squares.

#### 2.2 Chemical analyzes

Arsenic concentration in the soil surfaces was determined on site with X-ray fluorescence (NITON) apparatus. Speciation of arsenic was performed by HPLC-ICP-MS after H<sub>3</sub>PO<sub>4</sub> extraction. Diphenylarsinic acid was determined by HPLC-DAD and apolar organoarsenicals by GC/MS.

#### 2.3 Bio-indicators

As(III)-oxidizing and As(V)-reducing microbial concentrations were evaluated by Most Probable Number methods (5 tubes), using specific media in microplates. As(III) was visualized in the wells by adding pyrrolidine dithio-carbamate (Thouin *et al.*, 2016). Plants were collected in the four zones described previously. On the reference, low and medium zones, three and five species of plants were collected to measure As bioavailability and the lipidic bioindicator Omega-3 Index. Arsenic bioavailability was analyzed from collected plants by ICP-MS after mineralization by acid

Table 1. Concentration of arsenic species in the soil samples. ND: not detected

Zones	As(V) mg kg <sup>-1</sup>	As(III) mg kg <sup>-1</sup>	Diphelylarsinic acid mg kg <sup>-1</sup>	Triphenylarsnine mg kg <sup>-1</sup>
Reference	15	0.3	ND	ND
Transect	20	0.2	ND	ND
Low				
Transect	192	1.4	ND	ND
Medium				
Transect	756	14.4	2.19	0.95
High				

attack. For the measurement of the lipidic bioindicator, the leaf fatty acid composition was analyzed by GC-FID on the plants collected on the site but also on lettuces grown under controlled conditions on the four collected soils.

### 3 RESULTS AND DISCUSSION

#### 3.1 Arsenic in soil

Arsenic concentration varied from 15 mg kg<sup>-1</sup> in the reference not polluted area of the site, to 775 mg kg<sup>-1</sup> in the highly polluted spot (Table 1).

The main arsenic form was arsenate (As(V)), however As(III) proportion was higher in the polluted spot. Two organic molecules derived from the weapons were detected only in the highly polluted zone: diphenylarsinic acid and triphenylarsine.

#### 3.2 Bio-indicators

The collected plants were *Brassica napus*, *Plantago major*, *Carduus* sp. in the reference zone; *Brassica napus*, *Plantago major*, *Carduus* sp., and *Chamaemelum nobilis* in the low-pollution zone; *Brassica napus*, *Plantago major*, *Carduus* sp. and *Chamaemelum nobilis* in the medium-polluted zone; *Carduus* sp., and *Chamaemelum nobilis* in the high-pollution zone. Unidentified species were also collected in all zones except the highly-polluted one. Arsenic was more phytoavailable on the medium polluted soil than on the reference and the low polluted soils for all the three species collected. On the highly polluted soil, only one species could be analyzed. For this species, the As content was higher in the highly polluted zone. However, regardless of the zones, the As content in plants was very low compared to plants found on other As-polluted sites. In both site plants and lettuces, the lipidic bioindicator was lower on the medium polluted soil. It seems that the medium polluted soil was the most toxic for plants. This result might be interpreted in the light of further soil characterizations. As(III)-oxidizing microbes were more numerous in the highly polluted soils than




Figure 1. Evaluation by MPN methods of As(III)-oxidizing (A) and As(V)-reducing (B) microorganisms in the soil samples.

in the other samples. The same tendency was observed for As(V)-reducing microbes, but less significantly (Fig. 1).

### 4 CONCLUSIONS

Whereas the studied site was polluted one century ago, and was used for agriculture for years, impact of the pollution level is still observable on the diversity of wild plants. Arsenic level seems to exert influence on the soil composition in As-transforming microbes. These results suggest that arsenic was sufficiently bio-available to exert a selection pressure on the microbial communities. Next experimental steps will include the study of the influence of agricultural practices on the fate of As, through a biogeochemical approach including modelling, and connected with risk assessment.

### ACKNOWLEDGEMENTS

AgriAs is co-funded by EU and the Academy of Finland, L'Agence Nationale de la Recherche, Bundesministerium für Ernährung und Landwirtschaft and Forskningsrådet FORMAS under the ERA-NET Cofund WaterWorks2015 Call. This ERA-NET is an integral part of the 2016 Joint Activities developed by the Water Challenges for a Changing World Joint Programme Initiative (Water JPI).

### REFERENCES

- Naujokas, M.F., Anderson, B., Ahsan, H., Vasken Aposhian, H., Graziano, J.H., Thompson, C. & Suk, W.A. 2013. The broad scope of health effects from chronic arsenic exposure: update on a worldwide public health problem. *Environ. Health Persp.* 121, 295–302.
- Thouin, H., Le Forestier, L., Gautret, P., Hube, D., Laperche, V., Dupraz, S., & Battaglia-Brunet, F. 2016. Characterization and mobility of arsenic and heavy metals in soils polluted by the destruction of arsenic-containing shells from the Great War. *Sci. Total Environ.* 550: 658–669.