

Weathering processes in the Gault Clay formation, Paris Basin

Catherine Lerouge, Mathieu Debure, Jacques Brulhet, Christine Fléhoc, Sylvain Grangeon, Catherine Guerrot, Benoît Henry, Nicolas Maubec, Romain Millot, Eric Proust, et al.

► **To cite this version:**

Catherine Lerouge, Mathieu Debure, Jacques Brulhet, Christine Fléhoc, Sylvain Grangeon, et al.. Weathering processes in the Gault Clay formation, Paris Basin. Euroclay 2019, Jul 2019, Paris, France. hal-02073960

HAL Id: hal-02073960

<https://hal-brgm.archives-ouvertes.fr/hal-02073960>

Submitted on 20 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Weathering processes in the Gault Clay formation, Paris Basin

Catherine Lerouge^{1,*}, Mathieu Debure¹, Jacques Brulhet³, Christine Fléhoc¹, Sylvain Grangeon¹, Catherine Guerrot¹, Benoit Henry¹, Nicolas Maubec¹, Romain Millot¹, Eric Proust¹, Philippe Landrein², Benoit Madé², Francis Claret¹, Jean-Charles Robinet²

¹BRGM, 45060 Orléans, France

²Andra, 92298 Chatenay-Malabry, France

³B3J, 77920 Samois sur Seine, France

*c.lerouge@brgm.fr

The Lower Cretaceous Gault marine clay formation outcrops in the area of Brienne-le-Chateau (north-eastern France) currently investigated in the context of a sub-surface repository for low-level radioactive waste. This work aimed at investigating weathering processes (depth variations, chemical reactions) and water transfers in the Gault Clay from present-day to millennial timescale. In this framework, we applied a multi-disciplinary approach to rock and water samples collected in 5 m- deep pits and deep boreholes including the critical zone developed in the Gault clay. The pits were dug along transect on the eastern side of the Aube river that crosscut two successive hillslopes probably corresponding to two alluvial layers; the boreholes were drilled on the ridgetops and in the valleys (Lerouge et al., 2018).

Field work provided evidence of Quaternary surficial formations ranging between ~5 m in valleys and on lower parts of the slopes, 5m to 10 m in filled paleo valleys localized under ridgetops, and less than 1 m on the upper part of the slopes. Slides of surficial formations and of Gault clay moved downslopes. Water comings at different depths in the surficial formations were observed during the dug of the pits. Some waters come also above slide plans at the top of Gault Clay, rather suggesting lateral water transfers. Stable isotopes (¹⁸O, D) of the ground waters and pore waters in Gault clay show evaporation processes down to ~10 m depth.

Powder X-ray diffraction performed on bulk rock and clay fractions, and cation exchange capacity measurements of Gault clay and slide plans in contact with surficial formations give evidence of levels rich in swelling clay and of dissolution of pristine marine carbonates. Some secondary carbonate concretions result from precipitation of recent fluids.

Degassing of samples conditioned under He atmosphere provided evidence of a very high CO₂ production in soil (0-30 cm), and a high CO₂ degassing associated to oxygen peak in the first 2-10 m Gault Clay. The CO₂ degassing increase in weathered Gault clay relative to preserved one results from calcite dissolution due to pyrite oxidation and to organic matter degradation. The $\delta^{13}\text{C}$ of CO₂ indicates that the organic matter degradation is a major source of CO₂ down to 10-12 m, maximum depth at which we observed fossil roots. Then the CO₂ degassing decreases down to a constant value in preserved Gault Clay, where the carbonate system and the mineral assemblage control dissolved carbonates in pore waters.

Strontium (⁸⁷Sr/⁸⁶Sr) and lithium ($\delta^7\text{Li}$) isotopic data of the exchangeable fraction allow well distinguishing surficial formations (~2-5 m) from Gault clay, and altered (~2-20 m) from unaltered Gault Clay (>~15 m).

Combined data rather indicate that weathering processes are occurring as deep as 20 m below the surface with maximum intensity of alteration in the ~2-10 m zone.

Preferred format: Oral

Session number and name: B4 – mudrocks: composition, fabric, and provenance – fundamental research and practical application of the data