

# How both reactive bedrock accessory minerals and vegetation affect the behaviour of REE, U and Th during silicate weathering: Mule Hole watershed, South India

Jean-Jacques Braun, Jean Riotte, Shrema Battacharya, S. Subramanian, Aurélie Violette, Priscia Oliva, Jean-Christophe Maréchal

## ► To cite this version:

Jean-Jacques Braun, Jean Riotte, Shrema Battacharya, S. Subramanian, Aurélie Violette, et al.. How both reactive bedrock accessory minerals and vegetation affect the behaviour of REE, U and Th during silicate weathering: Mule Hole watershed, South India. Goldschmidt 2014, Jun 2014, Sacramento, United States. hal-02271945

HAL Id: hal-02271945

<https://hal-brgm.archives-ouvertes.fr/hal-02271945>

Submitted on 27 Aug 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **How both reactive bedrock accessory minerals and vegetation affect the behaviour of REE, U and Th during silicate weathering : Mule Hole watershed, South India**

JEAN JACQUES BRAUN<sup>1\*</sup>, JEAN RIOTTE<sup>1</sup>,  
SHREMA BATTACHARYA<sup>1</sup>, S. SUBRAMANIAN<sup>1,2</sup>,  
AURELIE VIOLETTE<sup>3</sup>, PRISCIA OLIVA<sup>3</sup>  
AND JEAN-CHRISTOPHE MARECHAL<sup>4</sup>

<sup>1</sup>IFCWS IISc-IRD joint laboratory, Bangalore, India  
jjbraun1@gmail.com (\* presenting author)

<sup>2</sup>Dept. of Materials Engg., IISc., Bangalore, India  
ssmani@materials.iisc.ernet.in

<sup>3</sup>GET, Toulouse, France

<sup>4</sup>BRGM, Montpellier, France, jc.marechal@brgm.fr

REE, U and Th are often found as major elements in the same accessory minerals and secondary phases, both at whole rock and regolith scale. The importance of these phases on biogeochemical budgets is often ignored because of their relatively low abundance of  $\leq 2\%$  by volume. The mobilization, redistribution and fractionation of REE, U and Th during weathering processes have been investigated in the heterogeneous parent gneiss and soils of the Mule Hole small experimental watershed, South India.

The REE-U-Th mineralogy has been studied by coupling SEM-EDX, EPMA and LA-ICPMS analyses. The elemental concentrations have been measured by ICPMS in rocks, regolith, plants and water reservoirs (rainfall, throughfall, overland flow water, pore water, groundwater, stream water). Leaching experiments have been carried out on both fresh gneiss and soil samples to ascertain the kinetics of the elemental sources.

For REE-U-Th, monazite, allanite, baesnesite, xenotime are major bearing minerals and apatite and titanite are minor phases. Low temperature hydrothermal processes have a strong influence on the replacement of monazite and allanite in the fresh rock. Monazite and xenotime, identified in the stream sediments, are more resistant to weathering. Allanite, baesnesite, apatite and titanite are completely broken down in the early stages of weathering. Brockite and thorite have been identified as secondary weathering phases of monazite. The initial mineralogical distribution in bed rock has a significant incidence of the REE, Th and U mobility at the watershed scale.

Hydrological model-based approach permits to estimate the elemental fluxes cycled through the vegetation and regolith of REE, U and Th. Vegetation has a significant effect on the transfer of easily available REE in soils.