



HAL
open science

Implications of the 2740 Ma Cote Gold Au(-Cu) deposit for Archean gold metallogeny and porphyry Au deposit formation in the Archean Swayze greenstone belt, northern Ontario

Daniel J Kontak, Laura Katz, Jocelyn Smith, Blandine Gourcerol, Bruno Lafrance, Benoit Dube, Joseph A Petrus, Robert Creaser, Mostafa Fayek

► **To cite this version:**

Daniel J Kontak, Laura Katz, Jocelyn Smith, Blandine Gourcerol, Bruno Lafrance, et al.. Implications of the 2740 Ma Cote Gold Au(-Cu) deposit for Archean gold metallogeny and porphyry Au deposit formation in the Archean Swayze greenstone belt, northern Ontario. RFG 2018 - Resources for Future Generations, Jun 2018, Vancouver, Canada. hal-02284250

HAL Id: hal-02284250

<https://brgm.hal.science/hal-02284250>

Submitted on 11 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



An IUGS Event



RFG 2018
RESOURCES FOR FUTURE GENERATIONS

PREMIER INTERNATIONAL CONFERENCE ON
ENERGY · MINERALS · WATER · THE EARTH

June 16-21, 2018

Vancouver Convention Centre, BC, Canada

1982 - Implications of the 2740 Ma Cote Gold Au(-Cu) deposit for Archean gold metallogeny and porphyry Au deposit formation in the Archean Swayze greenstone belt, northern Ontario

Daniel Kontak - *Laurentian University*

Laura Katz - *Iamgold Corporation, Gogama, ON*

Jocelyn Smith - *Iamgold Corporation, Gogama, ON*

Blandine Gourcerol - *Bureau de Recherches Géologiques et Minières (BRGM), Orléans, France*

Bruno Lafrance - *Harquail School of Earth Sciences, Laurentian University, Sudbury, ON*

Benoit Dube - *Geological Survey of Canada, Quebec, PQ*

Joe Petrus - *Harquail School of Earth Sciences, Laurentian University, Sudbury, ON*

Robert Creaser - *Department of Earth Sciences, University of Alberta, Edmonton, AB*

Mostafa Fayek - *Department of Geological Sciences, University of Manitoba, Winnipeg MB,*

Ore-deposit models, which are fundamental to mineral exploration, result from integrating robust field observations with high-quality mineral-chemical-isotopic data. New discoveries departing from current models demand explanation, and in some cases new deposit models result (e.g., IOCG). The recent (2009/2010) discovery of the Cote Gold Au(-Cu) deposit in the Archean Swayze greenstone belt, northern Ontario, challenges the age-restricted view of porphyry Cu-Au models. The +8 Moz Au deposit is hosted by a subvolcanic tonalite-diorite intrusive complex where magmatic-hydrothermal breccia bodies occur. Gold mineralization, of disseminated-, fracture- and vein-types, is spatially associated with hydrothermal biotite and muscovite alteration that is similar to potassic and phyllic alteration, respectively, in porphyry-type settings. A robust program involving core logging and field mapping with structural analysis integrated with detailed petrographic studies, geochronology, lithogeochemistry, isotopes, and fluid inclusion studies (microthermometry and evaporate mound SEM-EDS analysis) was used to assess the deposit's origin. That the age of host rocks (U-Pb zircon, titanite) and timing of alteration (U-Pb titanite) and mineralization (Re-Os molybdenite) centre on 2740 Ma and pre-dates the age of regional deformation (ca. 2680 Ma), in addition to the co-spatial nature of mineralization and alteration, suggests a magmatic-hydrothermal connection. Structural analysis of auriferous quartz veins also indicates a relationship to the intrusive centre versus regional deformation, which is further supported by a 2740 Ma molybdenite age for one such gold-mineralized vein. Fluid-chemical data indicates fluid mixing with $\delta^{34}\text{S}$ (py, cpy, moly = $0 \pm 1\%$) and fluid inclusions (mixed H₂O-CO₂ fluid (XCO₂=0.10); Na-K-Ca-Fe-Mn-Cl-F-S chemistry) suggesting a magmatic fluid reservoir whereas $\delta^{18}\text{O}$ (qtz = 7-12‰) also indicates a possible seawater contribution. These observations are best reconciled with a deposit model involving Au(-Cu) mineralization originating from exsolution of magmatic fluids from a high-level, hydrous intermediate magma in the same manner as models for younger porphyry analogues.