

## **Tectonometamorphic evolution of the North Ouaddaï Massif (Am Zoer area, E. Chad)**

Alexis Plunder, Jérémie Melleton, Matthieu Chevillard, Guillaume Vic, I Al-Gadam, E Djedouboum, A.-S André, Abdeltif Lahfid, Philippe Lach, Guillaume Wille, et al.

► **To cite this version:**

Alexis Plunder, Jérémie Melleton, Matthieu Chevillard, Guillaume Vic, I Al-Gadam, et al.. Tectonometamorphic evolution of the North Ouaddaï Massif (Am Zoer area, E. Chad). 2019. hal-02329009

**HAL Id: hal-02329009**

**<https://hal-brgm.archives-ouvertes.fr/hal-02329009>**

Preprint submitted on 23 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## ***Tectonometamorphic evolution of the North Ouaddaï Massif (Am Zoer area, E. Chad).***

A. Plunder<sup>1</sup>, J. Melleton<sup>1</sup>, M. Chevillard<sup>1</sup>, G. Vic<sup>1</sup>, I. Ousman Al-Gadam<sup>2,3</sup>, E. Djedouboum<sup>2</sup>, A.-S. André Meyer A. Lahfid<sup>1</sup>, P. Lach<sup>1</sup>, G. Wille<sup>1</sup>, S. Colin<sup>1</sup>, and O. Rouzeau<sup>1</sup>

<sup>1</sup> BRGM, 3. Av. C. Guillemin, 45000 Orléans, France.

<sup>2</sup> Ministère des mines et de l'industrie, République du Tchad

<sup>3</sup> Laboratoire GéoRessources, Université de Lorraine, 54506 Vandoeuvre-lès-Nancy, France.

The Saharan metacraton was assembled during pre-Neoproterozoic to Neoproterozoic times and highly remobilised during the Pan African stages. The Ouaddaï massif in Eastern Chad represents the core of the Saharan metacraton but our knowledge of the geological history is still poorly documented. We here present the combination of field data with a petrological study and geochronological constraints to decipher the different events that builded up the Ouaddaï massif. The study area is located in the surroundings of Am Zoer at the northern tip of the massif. It mainly corresponds to high-grade gneissic metagranite or metasediments, sometimes migmatitized containing rare mafic granulite. All these metamorphic rocks are intruded by an undated granitic series. Mafic granulites interbedded with the migmatite give Pressure-Temperature conditions of formation around 1.2 GPa and 850°C. Sillimanite-garnet bearing migmatites give contrasting lower Pressure-Temperature conditions of formation around 0.8 GPa and 700°C and graphitic schist gives temperature of at least 640°C. We dated samples using the U/Pb method on zircon. Granitoids yield ages clustering during the Panafrican stage (ca. 620-585 Ma). We also document the inheritance of Paleoproterozoic (ca. 1815 Ma) and Archean (ca. 2810 Ma) events in both granitoids and from detrital zircon in sandstones. The lack of Mezoproterozoic age is coherent with a Gondwanian origin for the sediments. Using our data, we discuss the timing of assembly of the Saharan craton and compare with the nearby massif (Darfour, Tibesti, Mayo Kebi). We also discuss the remobilization of the Saharan Craton during the Neoproterozoic and the associated collisional processes.