***Geochemical and geochronological record of the Andaman
Ophiolite, SE Asia: from back-arc to forearc during subduction
polarity reversal?***

Debaditya Bandyopadhyay^{1,2*}, Biswajit Ghosh¹, Carl Guilmette³, Alexis Plunder^{4,5}, Fernando Corfu⁶, Eldert L. Advokaat^{4,7}, Pinaki C. Bandopadhyay¹, Douwe J.J. van Hinsbergen⁴

1. Department of Geology, University of Calcutta, 35 Ballygunge Circular Road, Kolkata-700019, India
2. Department of Geology, University of North Bengal, Darjeeling-734013, India
3. Département de Géologie et de Génie Géologique, Université Laval, Québec, Québec, Canada
4. Department of Earth Sciences, Utrecht University, Princetonlaan 8A, 3584 CB Utrecht, The Netherlands
5. BRGM, 3 avenue Claude Guillemin, BP 36009, 45060 Orléans Cedex 2, France
6. Department of Geosciences and CEED, University of Oslo, Oslo, Norway
7. now at School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom

*Corresponding author: Debaditya Bandyopadhyay (debaditya.b2r@gmail.com)

Abstract

Ophiolites are widely studied to unravel how new subduction zones form. They may contain crustal and mantle rocks that formed during juvenile stages of intra-oceanic subduction, modifying the pre-existing oceanic lithosphere within which subduction started, and in which a magmatic arc formed upon subduction maturation. Previous geochemical work on the Cretaceous Andaman Ophiolite and its metamorphic sole, located in the forearc of the Sunda subduction zone of south-east Asia, has revealed geochemical signatures that are difficult to reconcile in a single tectonic setting but may rather record different stages in a longer evolution. A recent kinematic reconstruction proposed that the Andaman Ophiolite may have formed during subduction initiation in a former back-arc basin, when the former (Woyla) arc collided with the Sundaland continent. Here, we evaluate whether such a scenario may provide a feasible context to explain the enigmatic geochemical signatures preserved in the Andaman Ophiolite. To this end, we provide new, and review existing geochemical as well as geochronological constraints on the formation of its crustal rocks, as well as the evolution of its mantle portion. We identify mafic magmatic rocks and cogenetic plagiogranites that are consistent with formation in a magmatic arc, whereas other magmatic rocks, as well as metamorphic sole protoliths, have characteristics indicative of a back-arc origin. Three new, and two previous zircon U/Pb ages of arc magmatic rocks give a 99-93 Ma age range, but we also identify an inherited ~105 Ma age. This latter age coincides with Ar/Ar cooling ages of the Andaman metamorphic sole, and with a plagioclase xenocryst age from recent Barren Island volcanics east of Andaman. The geochemical and geochronological constraints of the Andaman ophiolites are straightforwardly explained in the context of the regional kinematic history: (1) The original lithosphere formed in the back-arc basin of the Woyla intra-oceanic arc; (2) Subduction initiation and SSZ ophiolite formation within this basin occurred around or slightly before 105 Ma; (3) This was followed by arc magmatism between 99 and 93 Ma upon subduction maturation.

Middle Andaman (Panchwati coast segment)

NOT TO SCALE

Highlights

- Re-evaluated geochemical characters of Andaman Ophiolite including metamorphic sole
- Magmatic rocks date 98-93 Ma; an inherited ~105 Ma age represents the SSZ spreading
- Results consistent with subduction initiation in back-arc during polarity reversal

Geochemical and geochronological record of the Andaman

Ophiolite, SE Asia: from back-arc to forearc during subduction

polarity reversal?

Debaditya Bandyopadhyay^{1,2*}, Biswajit Ghosh¹, Carl Guilmette³, Alexis Plunder^{4,5}, Fernando Corfu⁶, Eldert L. Advokaat^{4,7}, Pinaki C. Bandopadhyay¹, Douwe J.J. van Hinsbergen⁴

1. Department of Geology, University of Calcutta, 35 Ballygunge Circular Road, Kolkata-700019, India

2. Department of Geology, University of North Bengal, Darjeeling-734013, India

3. Département de Géologie et de Génie Géologique, Université Laval, Québec, Québec, Canada

4. Department of Earth Sciences, Utrecht University, Princetonlaan 8A, 3584 CB Utrecht, The Netherlands

5. BRGM, 3 avenue Claude Guillemin, BP 36009, 45060 Orléans Cedex 2, France

6. Department of Geosciences and CEED, University of Oslo, Oslo, Norway

7. now at School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom

*Corresponding author: Debaditya Bandyopadhyay (debaditya.b2r@gmail.com)

Abstract

Ophiolites are widely studied to unravel how new subduction zones form. They may contain crustal and mantle rocks that formed during juvenile stages of intra-oceanic subduction, modifying the pre-existing oceanic lithosphere within which subduction started, and in which a magmatic arc formed upon subduction maturation. Previous geochemical work on the Cretaceous Andaman Ophiolite and its metamorphic sole, located in the forearc of the Sunda subduction zone of south-east Asia, has revealed geochemical signatures that are difficult to reconcile in a single tectonic setting but may rather record different stages in a longer evolution. A recent kinematic reconstruction proposed that the Andaman Ophiolite may have formed during subduction initiation in a former back-arc basin, when the former (Woyla) arc collided with the Sundaland continent. Here, we evaluate whether such a scenario may provide a feasible context to explain the enigmatic geochemical signatures preserved in the Andaman Ophiolite. To this end, we provide new, and review existing geochemical as well as geochronological constraints on the formation of its crustal rocks, as well as the evolution of its mantle portion. We identify mafic magmatic rocks and cogenetic plagiogranites that are consistent with formation in a magmatic arc, whereas other magmatic rocks, as well as metamorphic sole protoliths, have characteristics indicative of a back-arc origin. Three new, and two previous zircon U/Pb ages of arc magmatic rocks give a 99-93 Ma age range, but we also identify an inherited ~105 Ma age. This latter age coincides with Ar/Ar cooling ages of the Andaman metamorphic sole, and with a plagioclase xenocryst age from recent Barren Island volcanics east of Andaman. The geochemical and geochronological constraints of the Andaman ophiolites are straightforwardly explained in the context of the regional kinematic history: (1) The original lithosphere formed in the back-arc basin of the Woyla intra-oceanic arc; (2) Subduction initiation and SSZ ophiolite formation within this basin occurred around

43 or slightly before 105 Ma; (3) This was followed by arc magmatism between 99 and 93 Ma
44 upon subduction maturation.

45

46 **Keywords**

47 Andaman Ophiolite, Agglomerate, Plagiogranite, U-Pb Zircon age, Tectonic discrimination,
48 Subduction initiation

1 Introduction

Ophiolites, oceanic lithosphere fragments that became uplifted and exposed, normally as a result of underthrusting of (subducting) buoyant continental lithosphere, or due to formation of accretionary prisms (e.g. Dewey, 1976; Wakabayashi and Dilek, 2003), are widely recognized as key geological archive to reconstruct the tectonic evolution of lost oceanic plates and plate boundaries (e.g. Pearce et al., 2003; Reagan et al., 2010; Stern et al., 2012; van Hinsbergen et al., 2015). Extensive geochemical work on modern oceanic settings and ophiolite records has revealed that oceanic crust at mid-ocean ridges, back-arc basins, oceanic plateaus, or forearc setting above subduction zones can be geochemically distinguished (e.g. Dilek and Furnes, 2011; 2014; Pearce, 2008; 2014). Particularly the forearc settings, where so-called ‘supra-subduction zone (SSZ) ophiolites’ form, are of major interest to the geodynamic community because these are thought to signal juvenile stages of subduction (Pearce et al., 1984; Stern and Bloomer, 1992; Shervais, 2001). The majority of ophiolites are of this SSZ type and are conventionally explained as the result of upper plate extension at an incipient subduction zone, leading to synchronous decompression melting of upwelling fertile asthenospheric mantle (Stern and Bloomer, 1992; Shervais, 2001). Typical SSZ ophiolites consist of a sequence starting with mid-ocean ridge basalt (MORB)-like forearc basalts (FAB) (Reagan et al., 2010). Subsequently, devolatilization of the gradually subsiding downgoing plate is inferred to lower the solidus of the mantle wedge, causing higher degrees of melting, and producing evolved arc volcanics (particularly at low melting pressures - boninites), followed by mature arc magmatism (Dilek and Furnes, 2011; Whattam and Stern, 2011; Stern et al., 2012). With this general model, ophiolite stratigraphy has been routinely used as a proxy to reconstruct the mechanism and chronology of subduction initiation that took place in earlier geologic history (e.g. Dilek and Furnes, 2011; Parlak 2016). But, for ophiolites exposed on the Andaman Islands in the east Indian Ocean there

appears to be no single tectonic setting that can explain the observed geochemical results (Bandyopadhyay et al., 2020; Plunder et al., 2020).

Located in the forearc east of the Sunda-Burma subduction zone, between Sumatra and Myanmar, the ophiolite suite of the Andaman-Nicobar Accretionary Ridge can potentially provide insight into subduction initiation in the eastern Neotethyan realm that existed between India, Australia and Eurasia (Fig. 1a, 2). The ophiolites were originally regarded as fragments of subducted Indian plate oceanic lithosphere (Curry et al., 1979; Mukhopadhyay and Dasgupta, 1988; Pal et al., 2003; Curry, 2005) or seamounts thereon (Acharyya et al., 1990; Sengupta et al., 1990; Acharyya, 2007) that were off-scraped and accreted to the upper, Eurasian margin. However, Pal (2011) suggested that both accreted Indian plate MORB mantle and original mantle wedge of the SE Asian plate are exposed, overlain by supra-subduction zone (boninitic) lavas and subsequent island arc tholeiitic (IAT) volcanics reflecting arc maturation. The Andaman Ophiolites also have a metamorphic sole (Pal and Bhattacharya, 2010; Plunder et al., 2020), which typically forms during subduction initiation and is commonly found below supra-subduction zone ophiolites (e.g. Hacker and Gnos, 1997; Agard et al., 2016; Soret et al., 2017; Guilmette et al., 2018). U-Pb zircon ages for plagiogranites with island arc geochemical signatures (Jafri et al., 1995) yielded ages of 93.6 ± 1.3 Ma (Sarma et al., 2010) and 95 ± 2 Ma (Pedersen et al., 2010) show that by this time, the Andaman Ophiolites were located above and close to a subduction zone. Thus, explaining the formation of the Andaman Islands as a regular SSZ ophiolite may be tempting (Pedersen et al., 2010).

However, the existing geochemical and geochronological constraints of the Andaman Ophiolite appear to be more diverse than straightforwardly explained in such a setting (Ghosh et al., 2017; Bandyopadhyay et al., 2020, and references therein). Specifically, the presence of less depleted lherzolitic mantle, the geochemical characteristics of which are more

common for a back-arc basin instead of a SSZ ophiolite forearc spreading centre (Ghosh et al., 2018). In addition, the ophiolite is associated with explosive volcanic agglomerates that are indicative of mature arc volcanism, all within the same ophiolite suite, within a few kilometres from each other (Fig. 1b,c). Moreover, $^{40}\text{Ar}/^{39}\text{Ar}$ cooling ages of the Andaman metamorphic sole were recently shown to give ~105 Ma ages (Plunder et al., 2020), i.e. 10-12 Myr older than the subduction-influenced plagiogranites. This is surprising because globally there is a coincidence between sole cooling and supra-subduction zone ophiolite spreading (Dewey and Casey, 2013; van Hinsbergen et al., 2015). So, does the Andaman ophiolite challenge general ophiolite concepts? Or else, does it juxtapose magmatic and metamorphic rocks formed in different settings at different times?

A recent kinematic reconstruction of the eastern Indian Ocean provides predictions on the sequence of tectonic settings that the Andaman Ophiolites may have experienced through time (Fig. 2). In this paper, we aim to evaluate this reconstruction of the geochemical and geochronological record of the Andaman Sea. The reconstruction first restores the N-S opening of the Cenozoic Andaman Sea Basin to the east of the Andaman Islands, which brings the ophiolites to the west of Sumatra (van Hinsbergen et al., 2011). On western Sumatra lies the intra-oceanic Woyla Arc that collided with the Sundaland continent of SE Asia around 100 Myr ago (Barber et al., 2005; Advokaat et al., 2018). Prior to collision, there was westward subduction below the Woyla Arc (Barber et al., 2005), implying that the subduction zone that produced the Andaman metamorphic soles and arc magmatic rocks formed in the former back-arc region behind the Woyla Arc (Advokaat et al., 2018).

In this paper, we critically re-evaluate the field relationships and geochemical and geochronological constraints on the Andaman Ophiolite and its metamorphic sole, and present new geochemical results for the metamorphic sole and igneous rocks. We also present new zircon U-Pb geochronological results from gabbros and plagiogranites from three new

locations of Middle Andaman, South Andaman, and Rutland Island. Finally, we then evaluate whether the geochemical and geochronological characteristics of the Andaman Ophiolites may be readily explained in the context of the regional plate kinematic evolution.

2 Geological framework

The Andaman and Nicobar archipelago, a chain of islands in the NE Indian Ocean, stretches for over 700 km, with a maximum width of c. 60 km, in the forearc of the Sunda subduction zone (Fig. 1a). This archipelago occurs as the exposed part of a long ridge which extends from the Arakan–Yoma ranges of western Myanmar in the north to Sumatra in the south, and is flanked by the Andaman Sea to the east and the Bay of Bengal to the west (Fig. 1a; Bandopadhyay, 2017). Ophiolitic rocks in the Andaman and Nicobar islands were first reported by Karunakaran et al. (1964) and described as dismembered bodies (Hutchison, 1975). Being located between the Himalayan collision zone and the Indonesian arc system, this ophiolite has the potential to offer insights into the Tethyan tectonics and India-Asia collision as a whole, though natural isolation along with geographical inaccessibility left these islands unexplored for long. In the 1980-1990's a few geological investigations were carried out on the Andaman ophiolite (Ray, 1985; Ray et al., 1988; Acharyya et al., 1990; Jafri et al., 1990, 1995). In the last decade, especially after the great 2004 Sumatra-Andaman earthquake, and after recent volcanic eruptions on Barren Island in the active arc east of the Andaman islands, a significant amount of research has appeared in the international literature (Lay et al., 2005; Meltzner et al., 2006; Pal et al., 2007, 2010; Morishita and Ghosh, 2010; Sheth, 2014). Following that, many workers have focused on the ophiolites and other lithomembers of the islands to understand their evolution and bearing on regional plate

tectonics (Pal et al., 2003; Ghosh et al., 2009, 2017; Pedersen et al., 2010; Sarma et al., 2010; Pal, 2011; Bandopadhyay, 2012; Bandyopadhyay et al. 2020).

The Sunda trench in the Andaman region is accommodating highly oblique convergence between the Indo-Australian and Eurasian plates (Curry, 1989). To the north, the Sunda trench connects to the collisional margin between India and the west-Burma block (Myanmar), and the Main Frontal Thrust of the Himalaya (Fig. 1). The oceanic subduction to the south has resulted in an active volcanic arc extending from Sumatra to the central Myanmar Basin in north Myanmar, including two volcanic islands in the Andaman Sea - Barren and Narcondam (Acharyya, 2007).

The Andaman Sea basin is a pull-apart basin that links two main strike-slip corridors: the Sagaing Fault between the West Burma Block of Myanmar to the northeast, and the West Andaman and Old West Andaman Faults and the Sumatran Fault System to the southwest (Fig. 1a). This fault system separates the West Burma-Andaman forearc sliver from Sundaland and transferred it northward. Since the Late Pliocene, extension in the Andaman Sea was accommodated by oceanic spreading (Kamesh Raju et al., 2004; Curry, 2005). Prior to that time extension was accommodated by crustal stretching, with cooling ages of exhumed dredge samples from the Alcock and Sewell Rises adjacent to the Andaman oceanic basin suggesting Early Miocene extension and exhumation (Frerichs, 1967; Rodolfo, 1969a,b; Curry, 2005). Oligocene extension was documented in the Mergui-North Sumatran basin (Kamili et al., 1976; Polachan and Racey, 1994; Andreason et al., 1997), which suggests that the formation of the Burma-Andaman sliver plate started sometime in the Oligocene, consistent with recent findings of Sagaing Fault activity at 27.5 Ma (Morley and Arboit, 2019). Restoring all this extension brings the southern margin of the West Burma Block adjacent to north Sumatra, and the northern margin of the West Burma Block to the

171 south of the major Eo-Oligocene Mae Ping and Three Pagodas faults that cut NW-SE through
172 mainland Indochina, but not through the West Burma Block (van Hinsbergen et al., 2011).

173 This restoration also brings the Andaman ophiolites to the west of North Sumatra (van
174 Hinsbergen et al., 2011; Advokaat et al., 2018), immediate west of the Woyla Arc. This arc
175 complex consists of basaltic to andesitic volcanic rocks that include xenoliths of radiolarian
176 chert, dykes, and volcanoclastic sandstones lacking quartz, and shales, with no indications that
177 the arc was built on a continental basement (Barber, 2000; Wajzer et al., 1991). It is a
178 Cretaceous (~130-105 Ma) intra-oceanic arc, thought to have formed above a W-dipping,
179 intra-oceanic subduction zone that terminated upon collision with continental Sundaland
180 (Barber et al., 2005). Following collision, the subduction polarity switched and became
181 northeastward, like today, with the accreted Woyla Arc in an upper plate position (Wajzer et
182 al., 1991; Barber, 2000; Barber et al., 2005; Advokaat et al., 2018). Advokaat et al. (2018)
183 postulated that the initiation of subduction recorded in the Andaman Ophiolites is causally
184 related to the arrest of subduction upon collision of the adjacent Woyla Arc.

185 Recently, Plunder et al. (2020) estimated 105-106 Ma Ar/Ar cooling age from
186 amphibolites of dismembered metamorphic soles of the Andaman ophiolite, which is at least
187 10 Myr older than the previous magmatic ages from the plagiogranite. Such metamorphic
188 soles are thought to represent the uppermost slices of downgoing oceanic lithosphere that
189 became metamorphosed and accreted beneath the upper plate lithospheric mantle during
190 ophiolite spreading (e.g. Guilmette et al. 2008; 2009; 2012; van Hinsbergen et al., 2015;
191 Agard et al., 2016; Soret et al., 2017). Whilst metamorphic soles may form well before
192 suprasubduction zone spreading (Guilmette et al., 2018; Pourteau et al., 2019), their cooling
193 ages are systematically synchronous with the age of SSZ spreading (van Hinsbergen et al.,
194 2015). Therefore, Plunder et al. (2020) argued that the 93-95 Ma ages of the plagiogranites do
195 not reflect SSZ spreading of the Andaman ophiolite, but rather the development of a mature

arc in and on already existing SSZ crust, which then likely is 105 Ma old. Such an age was reported from a plagioclase xenocryst hosted in a recent Barren Island lava flow (Ray et al., 2015).

Towards the north, in the Indo-Burman Ranges, there are ophiolites with Early Cretaceous (116-133 Ma) crust (Liu et al., 2016a; Singh et al., 2017; Zhang et al., 2018; Aitchison et al., 2019). Because these are presently a part of the same forearc system as the Andaman-Nicobar Accretionary Ridge, they are widely assumed to have formed at the same subduction zone as the Andaman-Nicobar ophiolites (e.g. Liu et al., 2016a). Advokaat et al. (2018) conceptually explained the southward younging trend in the ophiolites as the result of a southward migration of a triple junction during the polarity reversal documented at Sumatra. However, recent paleomagnetic data from Westerweel et al. (2019) suggested that the Indo-Burman Ranges were located at the same latitude as Sumatra in the Cretaceous to Eocene. This implies that these blocks, and the ophiolites in their forearcs, were parts of different plates, and hence at different plate boundaries. In this paper we focus, therefore, on the Andaman Ophiolite and leave the relation with ophiolites in the Indo-Burman Ranges for future study.

3 Ophiolitic lithologies - field relationships and geochemical fingerprinting

Most of the Andaman islands are occupied by dense tropical forest and exposure is mostly restricted to coastal, river, and rare road outcrops. Exposures reveal ophiolitic rocks and underlying metamorphic sole and mélangé, an overlying Paleogene–Neogene turbidite sequence of clastic sedimentary rocks (Fig. 1b; Bandopadhyay and Carter, 2017). Below we will briefly summarize the field relationship of the different ophiolitic litho-units along with

the published geochemical and geochronological results, with special emphasis on the location-specific field disposition, which we believe holds the clue to resolve the existing gap areas regarding the genesis of this ophiolite.

The Andaman ophiolite suite, including the underlying sole and *mélange* rocks, is found in the hanging wall of the prominent, N-S striking Jarawa (or Jarwa) Thrust that runs over the central-western part of the Andaman Islands (Fig 1b). This thrust juxtaposes the ophiolite suite and overlying Paleogene sedimentary sequence westwards upon an accretionary prism of deep-marine, presumably post-Eocene turbidite sequences (Fig. 1b; Bandopadhyay and Carter, 2017). The Andaman ophiolite suite consists of tectonized, restitic mantle sequences (>700 m thick in North Andaman Island) that host chromitite pods, a plutonic crustal section mainly exposed as cumulate rocks (up to 150 m thick), and volcanic rocks (up to 400 m thick in South Andaman); notably with exposures mostly restricted to the east coast of North, Middle, South Andaman and Rutland Islands (Ghosh et al., 2009, 2013, 2017; Pal, 2011). Identifying the components of an idealized ‘Penrose’ (Anonymous, 1972) architecture of the oceanic lithosphere has been attempted for the Andaman ophiolite (Saha et al., 2010; Pal, 2011). However, it is important to note that while some elements of the Penrose sequence are present (peridotites, gabbros, pillow lavas, cherts, but sheeted dykes are absent), there is nowhere a coherent sequence exposed and contacts between ophiolitic rock units are poorly exposed. This may be a result of tectonic dismemberment, because the sequence has always been incomplete, and/or because there is a multiphase magmatic history preserved that is not typical for a Penrose sequence. In any case, the Andaman ophiolite suite contains different ophiolitic lithomembers, cropped out in various islands, that are put together in an ophiolite stratigraphy (Ghosh et al., 2017; Bandyopadhyay et al., 2020).

3.1 Metamorphic sole

Metamorphic rocks, comprising sheared meta-volcanics and meta-sediments, have been found confined to either the base of the ophiolite, immediately below the mantle peridotite (in North Andaman) or as subangular blocks within the mélangé (in Middle Andaman) (Pal et al., 2003). Pal and Bhattacharya (2010) interpreted these metamorphic rocks as the sub-ophiolitic greenschist facies rocks representing the sole of ophiolite slices (e.g. Wakabayashi and Dilek, 2003). Geochemically, protoliths of the metabasites are inferred to be subduction influenced (Pal and Bhattacharya, 2010). Towards south, along the Panchawati coast, Middle Andaman (*Location 1* – $12^{\circ}35'34''N$; $92^{\circ}57'26''E$; Figure 1b, c), we document different lithotypes occurring as blocks of up to 10 meters or more in a serpentinite matrix (see also Plunder et al., 2020). The rock types include metabasite (amphibolite), peridotite, mafic rocks, and metachert. These were emplaced over Paleogene clastic sedimentary rocks along a post-Paleogene thrust. The peridotite boulders along the coast are inferred to be the same as the to lherzolitic mantle peridotite (Ghosh et al., 2018) exposed in a nearby quarry section (Ghosh et al., 2013). Plunder et al. (2020) estimated peak metamorphic conditions for the sole amphibolites of ~0.9 GPa and ~675°C, followed by cooling below closure temperatures for argon in hornblende (~500°C) constrained by $^{40}\text{Ar}/^{39}\text{Ar}$ ages of 106.4 ± 2.1 Ma and 105.3 ± 1.6 Ma.

3.2 Mantle section

Previous work on the mantle section of the Andaman ophiolites has yielded disparate estimates of its chemical history (Ghosh et al., 2009, 2013, 2018; Pal, 2011). Peridotites are variably serpentinitised (40-80 vol.%) and display strong post-serpentinization deformation across the Andaman Islands. Locally, a crude primary foliation is discerned in the least serpentinitized samples. Petrography and mineral chemistry of the mantle sequence of the

Andaman ophiolite portray a pronounced variation between Rutland Island in the south and Middle and North Andaman towards the north (Ghosh et al., 2013). In Rutland Island and South Andaman, mantle rocks are mainly harzburgite-dominated where chrome spinel and clinopyroxene compositions suggest them to be residue of 14-18% melting of a fertile mantle (e.g. DMM; Workmann and Hart, 2005) in a SSZ environment (Ghosh et al., 2009, 2018). On the other hand, the mantle peridotites in Middle and North Andaman Island are fertile lherzolite that occasionally grades to clinopyroxene-bearing harzburgite, corresponding to <10% mantle melting with minor flux infiltration (Morishita et al., 2018a), with a geochemistry suggesting formation in a back-arc setting (Ghosh et al., 2018). Massive chromitite pods of 8-20 cm size (maximum up to 50 cm) within harzburgite have been reported from Rutland Island (Ghosh et al., 2009) whereas the same pods in lherzolitic mantle rocks have been reported from North-Andaman (Ghosh et al., 2013; Ghosh and Bhatta, 2014).

3.3 Transition between mantle and crustal rocks

Ghosh et al. (2014) interpreted a nearly 6 m thick zone along a road-cut section at Kodiaghat, South Andaman, exposing olivine-rich troctolite with evidence of being overlain by clinopyroxenites, wehrlites and gabbroic rocks, as a MOHO transitional zone (*Location 2 - 11°31'21"N; 92°43'08"E*) (Fig. 1b, 3a,b). Ghosh et al. (2014) described field relationships and clinopyroxene compositions that suggested a genetic link between olivine-rich troctolite, the layered gabbros, and the clinopyroxenite-wehrlite association. They also inferred from the chrome spinel composition that the olivine-rich troctolite was earlier a dunite formed by interaction between melt and a mantle similar to the depleted harzburgite of Rutland Island.

3.4 Crustal section

The lithotypes belonging to the crustal section of the Andaman ophiolite are less abundant than mantle rocks and best exposed in South Andaman along a road section on the way to Chidiyatapu (Fig. 1b), as well as in Rutland Island. The crustal section comprises continuous layered gabbroic rocks at its base, followed up-section by various plutonic rocks collectively named as high-level intrusives, and volcanics at the top. Layered gabbroic rocks, immediately above the cumulate sequence of the MOHO transition zone, are best exposed in the Kodiaghat section, South Andaman (*Location 2 - 11°31'21"N; 92°43'08"E*) (Fig. 1b, 3a,b) (Pal 2011, Ghosh et al., 2014). A wide variation in modal mineralogy has been documented within interlayered olivine gabbros (Ghosh et al., 2014, 2017). A coastal exposure in Bedonabad (*Location 3 - 11°34'01"N; 92°44'10"E*) (Fig. 1b) displays a variety of plutonic and volcanic-hypabyssal rocks. At this location we have found homogeneous gabbro (noncumulate gabbro) and a plagiogranite-diorite suite, together termed as 'high-level intrusives' (Pal et al., 2003). The plagiogranite-diorite suite and volcanic-hypabyssal rocks are often intermingled (Fig. 3c). Locally the plagiogranite appears intrusive within gabbro as irregular bodies (Jafri et al., 1995). Zircon from this plagiogranite yielded an age of 93.6 ± 1.3 Ma (Sarma et al., 2010). Plagiogranite and the non-pillowed volcanic rocks, wherever present, appear brecciated and we describe them as magmatic breccia. Near-vertical thin dykes of altered dolerite are common. We also documented basaltic enclaves within the plagiogranite-diorite suite. Plagiogranites of this location display a volcanic arc geochemical signature (Jafri et al., 1995). In Middle Andaman, an in-situ exposure (*Lat: N12°34'45"; Long: E92°57'41"*) (Fig. 1b,c) of leucocratic rocks belonging to the diorite – granodiorite series is in close association with agglomerates (Fig. 3d), suggesting a genetic linkage between the two. On Rutland Island, plagiogranite (*Location 4 - 11°24'10"N; 92°39'34"E*) (Fig. 1b) is locally exposed along the south-eastern coastal tract of the island, as small dyke-

like bodies showing mutually intrusive relationships with gabbro (Fig. 3e). Sheeted dyke complex has not been observed in the Andaman ophiolites (Ghosh et al., 2017).

Amongst the crustal rocks, volcanic units dominate. Two distinct groups of volcanic rocks – dubbed ‘pillow lavas’ and ‘East Coast volcanics’ were reported from the Andaman ophiolite based on their field characteristics (mainly in South Andaman) and geochemical affinity (Ghosh et al., 2017, and references therein). The ‘East Coast volcanics’, with oceanic island arc affinity, occur as separate thrust slices over the ‘Pillow lavas’ (Ray, 1985; Ray et al., 1988). An alternative classification proposed by Vohra et al. (1989), coined ‘upper lava’ for pillow basalts and ‘lower lava’ for the East Coast volcanic rocks, interpreted to tectonically overlie the former (see table 7.2 of Ghosh et al., 2017).

In South Andaman Island, a coastal exposure (*Location 5 – Corbyn’s Cove, South Andaman; 11°38’28’’N; 92°44’50’’E*) preserves tholeiitic pillow basalts (Ray et al., 1988) (Fig. 1b,3f). Hyaloclastites and (brecciated) pillow lavas with MORB geochemical affinity have been reported from this location (Jafri et al., 2010). Non-pillowed East Coast volcanic rocks are found in the coastal exposure in Bedonabad, South Andaman, (*Location 3 - 11°34’01’’N; 92°44’10’’E*). Ray et al. (1988) also concluded that the East Coast volcanics (basaltic andesite, acid volcanics) are genetically connected with the tholeiitic pillow basalts and formed on a marginal basin floor. Alternatively, a plume origin (Jafri et al., 1990) and a mid-ocean ridge setting (Srivastava et al., 2004) have been also proposed for the South Andaman volcanic rocks. With progressive development of the ophiolite concept, the South Andaman volcanic rocks, with a spectrum of geochemical variation - from MORB to boninites to evolved arc volcanics, have been linked with different stages of subduction by Pal (2011). This was later supported by Bhattacharya et al. (2013) based on volcanic rocks from Rutland Island. Close to the MOHO Transition Zone exposure of the Kodiaghat section, South Andaman (*Location 2 - 11°31’21’’N; 92°43’08’’E*, Fig. 1b) Ghosh et al. (2014)

reported East Coast volcanic samples with a similar geochemical signature to that of calculated melt compositions in equilibrium with clinopyroxene from olivine-rich troctolites and gabbroic rocks.

Near Chidiya Tapu (farther south from location 2, near to Mundapahar beach, *Location 6 - 11°29'33"N; 92°42'18"E*, Fig. 1b) we documented a ca. 100 m long coastal exposure that preserves an extensive agglomerate volcanic facies with a bimodal composition of clasts, consisting of mafic-intermediate volcanic rocks (andesite, dacite) and dominantly plagiogranite (Fig. 3g). The size, angularity and composition of the clasts vary from bomb to block tuff. Pedersen et al. (2010) reported a 95 ± 2 Ma age from a plagiogranite in South Andaman. Based on the location marked on their map and description that “the dated trondhjemite (plagiogranite) was sampled at a coastal exposure at Chidiya Tapu at the southern tip of South Andaman Island, where it occurs together with gabbros and cross-cutting dykes”, we have tried to access the location to document the relationship between volcanics and intrusives. Trondhjemite (plagiogranite) dykes mentioned by Pedersen et al. (2010) are not exposed at Chidiya Tapu and we presume that their 95 ± 2 Ma age dates the plagiogranite blocks contained in the agglomerate and formed during an explosive volcanic eruption. Geochemical investigation further inferred that they formed in a supra-subduction zone environment (Bhattacharya et al., 2020).

Volcanic agglomerates are abundant along the Panchawati coast, Middle Andaman (*Location 1 - 12°35'34"N; 92°57'26"E*; Figure 1b,c), where they form coherent blocks of tens of m across. The interlayered pyroclastic rocks are dominantly composed of andesitic (Pal and Bhattacharya, 2016) blocks of different sizes (m to cm scale), sorted in layers set in a volcanic matrix of andesitic composition (Fig. 3h,i). It is not clear whether these are located within the mélangé, or whether they belong on top of the ophiolite and were overthrust by the mélangé along post-Paleogene thrusts that cut the ophiolite and overlying Paleogene clastic

sequence. The latter option appears more likely as serpentinite-matrix mélanges including metamorphic sole blocks are located adjacent to the agglomerates, which do not appear internally deformed, or metamorphosed, and the agglomerates contain plagiogranites (in South Andaman) that are also found intruding the ophiolite.

Various hemipelagic sedimentary rocks are documented along a ~3 km stretch of the Panchawati coast (*Location 1* $-12^{\circ}35'34''N$; $92^{\circ}57'26''E$; Figure 1b). They constitute a volumetrically minor component in the Andaman ophiolite, occur either alternating with basalt or in short successions (~ 2.5 m thick) immediately overlying the extrusives. They are characterized by thin layers of red and light green mudstone, greenish and red chert, and limestone. Jafri et al. (1993) reported radiolarian cherts from South Andaman in association with pillow basalts (near Bedonabad), conglomerate and grit (near Bamboo flat). Several biostratigraphic age constraints have been proposed from hemipelagic sediments - Early Cretaceous based on radiolarian assemblage (Jafri, 1986); Early Eocene to Paleocene from radiolaria in red cherts and planktonic foraminifera in limestones (Roy et al., 1988); Eocene (Middle) and Cretaceous (Campanian) from radiolarian faunas (Ling and Srinivasan, 1993; Ling et al., 1995, 1996); Late Cretaceous to Paleocene (Pal et al., 2003). However, the field disposition of the cherts, from which this large spectrum of ages was determined, is poorly described in the literature, thus their stratigraphic position and formation with respect to the ophiolite is difficult to assign. Jafri et al. (2006) documented the presence of ash layers in radiolarian cherts in South Andaman and correlated them with a possible explosive volcanic event related to the East Coast volcanics during the Late Cretaceous.

4 Methodology

4.1 *Geochemistry – sampling rationale and analytical techniques*

Whilst extensive geochemical analyses have previously been performed on the ultramafic and mafic rocks of the Andaman Ophiolite suite (Ghosh et al., 2017, and references therein) no detailed and comprehensive geochemical analyses were previously reported from the pyroclastic agglomerates of South Andaman and Middle Andaman, nor for the amphibolites of the metamorphic sole. To aid the tectonic interpretation, we collected representative samples of agglomerates from the Chidiya Tapu coastal exposure (location 6) and Panchawati coast (location 1), as well as amphibolites from Middle Andaman. These amphibolites are studied in detail for metamorphic and thermochronological analysis by Plunder et al. (2020). Petrographic data of the selected samples are shown in figure 4 and Table 1. Based on this, fourteen samples were chosen for whole rock geochemistry (major and trace elements) (Table 2; Fig. 5,6). All samples were analysed at Actlabs (Canada) using the Lithoresearch package methodology¹. Analyses for blanks, replicates and standards are provided in supplementary material.

4.2 *U/Pb geochronology*

4.2.1 *Sampling rationale*

Previous geochronological studies on South Andaman plagiogranites gave 93-95 Ma ages for supra-subduction magmatism (Pedersen et al., 2010; Sarma et al., 2010). Because these plagiogranites are found in association basaltic andesitic agglomerates with island arc

¹ available online

<https://actlabs.com/geochemistry/lithogeochemistry-and-whole-rock-analysis/lithogeochemistry/>

geochemical affinity (see above), which is consistent with the geochemistry of the plagiogranites (Jafri et al., 1995; Bedonabad coast, location 3), Bandyopadhyay et al. (2020) suggested that the subduction initiation below the Andaman ophiolites may be older than the 95 Ma plagiogranite ages, as confirmed by the $^{40}\text{Ar}/^{39}\text{Ar}$ plateau ages of ~105-106 Ma of the metamorphic sole (Plunder et al., 2020), as well as the plagioclase xenocrysts hosted in a Barren Island lava flow (Ray et al., 2015). We have collected samples suitable for U-Pb zircon analysis from the previously undated outcrops in Middle Andaman and Rutland Island to test the robustness of the previous ages.

We also selected a layered gabbro from the Kadiaghat section, South Andaman that is geochemically linked to the East Coast volcanics (Ghosh et al., 2014). Dating the crystallization of this gabbro would provide constraints on the timing of the magmatic event from which resulted the East Coast volcanics. From the collected samples, three (marked in Fig. 1b) - **148A** [*Lat: N11°24'10"; Long: E92°39'34"; Rutland Island (location 4)*]; **SA-GAB1** [*Lat: N11°31'57"; Long: E92°43'25"; South Andaman (location 2)*] and **L-18** [*Lat: N12°34'45"; Long: E92°57'41"; Middle Andaman (location 1)*] proved to be most suitable for further geochronologic investigation. Petrographic description of the samples is in Table 1.

4.2.2 Analytical techniques

Zircon grains were extracted by crushing, milling and separation using a Wilfley table, and concentrated using a combination of magnetic separators and heavy liquids (methylene iodide). Individual grains were selected under an optical microscope and subjected to chemical abrasion (Mattinson, 2005). This was followed by spiking with a ^{202}Pb - ^{205}Pb - ^{235}U tracer, dissolution, and mass spectrometry, after the procedure detailed in Krogh (1973) with modifications described in Corfu (2004). For small amount of Pb measurements

were performed with an ion counting secondary electron multiplier. The obtained data were corrected with fractionation factors for Pb determined from the measured $^{202}\text{Pb}/^{205}\text{Pb}$ of the added spike and for U by 0.12%/amu. Blank corrections were 0.1 pg for U and ≤ 2 pg Pb. Additional common Pb in one fraction was corrected using a composition calculated with the model of Stacey and Kramers (1975). The data were also adjusted for a deficit of ^{206}Pb due to initial deficiency of ^{230}Th (Schärer, 1984) and the tracer was calibrated with reference to the ET100. Plotting and regressions were done with the Isoplot software package (Ludwig, 2009). The decay constants are after Jaffey et al. (1971). Detailed geochronological results are given in the section 5.2, Table 3, and Fig. 7.

5 Results

5.1 Geochemistry

We first note that most of the samples contain more than 1 wt% volatile (LOI), with a few agglomerates containing more than 5 to 10 wt% LOI. Such volatile contents are potentially indicative of elemental mobility during hydrothermalism or metamorphism (for example, Ca, Na and K are susceptible to mobility during greenschist and amphibolite facies and Si, Fe and Mg during high fluid-rock ratios in the hydrothermal systems). Accordingly, all ensuing interpretation and discussion will rest on ratios between elements deemed immobile (e.g. Ti, Zr, Y, Nb etc.) during hydrothermal alteration/metamorphism (e.g. Cann, 1970; Pearce and Cann, 1971; Pearce, 1975; Shervais, 1982; Pearce, 2014).

5.1.1 Major and minor elements

We first characterize our geochemical dataset with major element based binary, Harker-type diagrams (Fig. 5). To this end, we recalculated the bulk rock analyses on an

anhydrous basis to avoid the closure effect of high LOIs, and adjusted FeO and Fe₂O₃ as per
 Fe²⁺/Fe_{tot} = 0.85 (e.g. Schilling et al., 1983). Amongst the five analysed metamorphic sole
 samples from Middle Andaman, three amphibolite samples have a basic-intermediate
 transitional composition with SiO₂ = 50-53 wt% (Fig. 5). Two samples have outlying
 compositions; AN1704a with ~58.5 wt% SiO₂ belongs to the intermediate group (Fig. 5),
 whereas AN1703 is acidic with SiO₂ ~ 90 wt%. Its quartz-rich nature (up to 70%) with less
 than 10% amphibole suggests a non-volcanic protolith (probably meta-chert). Its chemistry
 will hence not be discussed further. The three basic-intermediate samples share similar
 compositions consistent with a primitive basalt protolith, with 1.3-1.5 wt% TiO₂ (Fig. 5a),
 ~13-14 wt% Al₂O₃ (Fig. 5b) and 9-10% CaO (Fig. 5c). Their Mg#, Cr and Ni contents are
 also typical of primitive basalts with values in the range of 0.61-0.63 (Fig. 5d), 170-280 ppm
 (Fig. 5e) and 80-170 ppm (Fig. 5f), respectively. By contrast, sample AN1704a has a much
 lower TiO₂ (0.5 wt%) and CaO (7 wt%) contents, similar Al₂O₃ (13.4 wt%) and significantly
 higher Mg# (0.72), Cr (410 ppm) and Ni (280 ppm) contents, hinting to a different protolith.
 Considering that the primary structures and textures of rocks from the metamorphic sole have
 been obliterated during tectonometamorphism, we tested the chemistry of the studied samples
 in a Mg# vs. SiO₂/Al₂O₃ (Kempton and Harmon, 1992) diagram (Fig. 5g) to identify samples
 that could have had cumulates as a protolith (excluding siliceous sample AN1703). The three
 samples with clustered compositions plot within the field of primitive basalts, confirming that
 their protoliths are representative of magmatic liquids and not cumulates. The outlying
 composition of sample AN1704a appears to result from the accumulation of pyroxene,
 consistent with its higher Cr and Ni contents. The sample still plots close to the field of
 primitive basalts, possibly representing pyroxene-phyric lava.

Intermediate agglomerate samples from Middle and South Andaman have relatively
 clustered compositions except for outlier sample AN1721b, petrographically identified as

brecciated rhyodacite (Table 1, Fig. 4j). This classification is consistent with its very high SiO_2 (~79 wt%), lower TiO_2 (0.27 wt%), Al_2O_3 (11 wt%) and $\text{Mg\#}$ (0.27), with undetectable Ni and Cr (Fig. 5a-f). All other agglomerate samples have intermediate compositions with SiO_2 ~53-58 wt% (Fig. 5a). TiO_2 is also fairly consistent across samples with contents ranging between 0.6 - 1.0 wt% (Fig. 5b). The Al_2O_3 and CaO contents are strongly variable, ranging between 15-20 wt% and 0.5-9 wt% respectively (Fig. 5c,d), and inversely correlated. The very low CaO contents of three Al_2O_3 rich samples (AN1705a, AN1705b, AN1710a) might reflect elemental mobility, although clustered SiO_2 and uncorrelated LOI contents do not support the hypothesis. $\text{Mg\#}$ are variable, ranging from 0.52 to 0.71 (Fig. 5d), and do not correlate with equally variable Cr and Ni contents of 20-580 ppm (Fig. 5e) and 30-250 ppm (Fig. 5f) respectively.

5.1.2 Trace elements

In the immobile element-based Total Alkali-Silica (TAS) proxy diagram (from Pearce, 2014, modified after Floyd and Winchester, 1975), which uses Nb/Y as a proxy for total alkalis ($\text{Na}_2\text{O} + \text{K}_2\text{O}$) and Zr/Ti as a proxy for SiO_2 , all the amphibolite samples fall in or very close to the basalt field (Fig. 6a). Agglomerates samples from Middle and South Andaman fall in the basalt to andesite, basaltic andesite field, except for sample AN1721b, which falls in the rhyolite-dacite field, again consistent with petrographic observation.

To characterize our volcanic rocks in a more effective manner, we plot the obtained geochemical dataset in a N-MORB (Sun and McDonough, 1989) normalized spider diagram where the elements are ordered according to their average incompatibility during mantle melting, which efficiently distinguishes subduction influenced volcanic rocks (i.e. arc origin) from uninfluenced (i.e. MORB; Pearce, 2014; Fig. 6b-d).

All metamorphic sole samples except AN1704a have subparallel patterns and are enriched between 0.8 and 4 times relative to MORB. These samples show no HREE fractionation (flat patterns), are slightly enriched in light rare earth elements (LREEs) and show slight negative Nb, Ta, Hf, Zr and Ti anomalies. Sample AN1704a is more depleted in HREEs and more enriched in LREEs and shows more pronounced negative anomalies (Fig. 6b).

Agglomerates in Middle Andaman show both HREE and LREE fractionation, being depleted in HREEs and enriched in LREEs relative to MORBs. They show pronounced negative Nb, Ta, and Ti anomalies and slight Zr-Hf anomalies ranging from positive for the more fractionated samples to negative for the less fractionated ones (Fig 6c). Sample AN1705c shows stronger LREE enrichment than the rest.

Agglomerates from South Andaman show patterns that are similar to those of volcanics of Middle Andaman and AN1704a (metamorphic sole) with fractionated LREE and HREE patterns and strong Nb-Ti negative anomalies. The Ta content was systematically below the 0.01ppm detection limit. Out of three analysed samples only AN1721b has undepleted HREEs that are enriched from 0.8 to 2 times NMORB, slightly fractionated LREE enriched as much as 4 times NMORB, and have very pronounced Nb-Ti negative anomalies, as well as slight negative Zr-Hf anomalies (Fig. 6d). The contrasting composition of the volcanic rocks is also represented in the chondrite normalized (Sun and McDonough, 1989) REE multi-element spidergram (Fig. 6e-g). Middle Andaman sole samples show an almost flat REE pattern (except AN1704a). Agglomerates from middle and South Andaman (and AN1704a) are generally characterized by strong LREE enrichment and HREE depletion (Fig. 6e-g). We conclude from the above that the protolith of the metamorphic sole (except AN1704a) possesses distinct geochemical characteristics (no HREE fractionation, slightly more LREE-enriched than NMORB, slight negative Nb-Ta-Zr-Hf-Ti anomalies) with respect

to the samples from the Middle and South Andaman agglomerates. Samples of Middle and South Andaman agglomerates have a general fractionated HREE pattern and strong negative Nb-Ta-Ti anomalies similar to that of metamorphic sole sample AN1704a.

5.2 *Geochronology*

Only a few zircon grains were recovered from gabbro sample SA-GAB1. Three short prismatic grains and one tip (grains 1-4, Supplementary fig. 1) were analysed yielding variable $^{206}\text{Pb}/^{238}\text{U}$ ages between about 105 and 98 Ma (Table 3, Fig. 7). The pattern can in principle be explained by variable Pb loss of 105 Ma zircon, but this seems unlikely as the grains were of good quality and were chemically abraded, which commonly removes disturbed domains quite efficiently. The alternative is the presence of more than one generation. This interpretation implies that the youngest date of 98.56 ± 0.33 Ma is a maximum, and possibly the real age for emplacement of the gabbro. We consider 105 Ma the age of an older component picked up at the latest stage of magmatism or melting of pre-existing gabbroic crust. The larger error of the 105 Ma date is a function of the limited amount of Pb (and U) present in that grain. This suggests here were two different magmatic events that occurred around 98 and 105 Ma.

Plagiogranite sample 148A contains an abundant population of equant to prismatic zircon (Supplementary fig. 1). Four analyses show a slight scatter yielding an average $^{206}\text{Pb}/^{238}\text{U}$ age of 98.74 ± 0.32 Ma dating emplacement. Zircon in plagiogranite L-18 occurs mostly as irregular, subsequent, anhedral grains and fragments, only partly showing euhedral faces or complete crystals (Supplementary fig. 1). The four analyses overlap giving an average $^{206}\text{Pb}/^{238}\text{U}$ age of 93.48 ± 0.10 Ma dating magmatic crystallization.

6 Geochemical synthesis – methods and limitations

6.1 *Geochemical characterization of volcanic rocks of Andaman ophiolite*

To resolve the debate on the geodynamic origin of the volcanic rocks, an extensive dataset has been compiled from the existing literature on Andaman and Nicobar Islands (Ghosh et al., 2017). Preliminary geochemical analyses of Ghosh et al. (2017) clearly revealed the potential of such a comparative study in explaining the varied geochemical spectrum of volcanics in a supra-subduction zone environment. Comparison of our new results with the previously published data provides comprehensive insights into the geochemical variability of the Andaman Nicobar volcanic rocks and their relation to the mantle rocks. To this end, we included our geochemical results of agglomerates, of sole (from location 1 and 6) and of two previously published dataset (Pedersen et al., 2010; Sachin et al., 2017) into the compilation of Ghosh et al. (2017) (supplementary table 2). Please note the lower lava and upper lava (as well as pillow basalt and east coast volcanics) distinction was followed after Ghosh et al. (2017, for details see table 7.2). The data sets of Sachin et al. (2017) and Jafri and Sheikh (2013) are from Great Nicobar and Bompoka Island, respectively (See Fig. 1a for location). From field, petrographic and geochemical perspective, the agglomerates from South and Middle Andaman are found to possess more or less similar characteristics and thus considered as a single group in our synthesis. The corresponding major element plots of the compiled dataset are given as overlay in figure 5. Variable and sometimes substantial LOI values have been reported for volcanic rocks, highlighting the possibility of element mobility. For this reason, and because Ghosh et al. (2017) have already attempted a geochemical characterization of the Andaman and Nicobar Islands volcanic rocks based on major elements, we herein focus on trace elements that are deemed immobile in

ophiolitic environments and show a better potential to accurately discriminate tectonic affinities (Dilek and Furnes, 2011; Pearce, 2014).

6.2 Geochemical modelling

Volcanic rocks are susceptible to post/syn extrusion alteration and fractionation, and thus inferring the correct parental melt chemistry is crucial to understand the genesis of the basalt in the source mantle. Although tricky, a reasonable estimate of a primary basalt could be achieved using: (1) a petrogenetic grid for mantle melting (e.g. PRIMELT2 model) (Herzberg and Asimow, 2008, and references therein); (2) thermodynamic phase equilibria model calculations (e.g. pMELTS; Ghiorso et al., 2002; Asimow et al., 2004); or (3) an iterative algorithm that allows to estimate primary basalt compositions using back calculations along the fractional crystallization path of a magma (e.g. PRIMACALC2, Excel based WINDOWS application software; Kimura and Ariskin, 2014). Apart from the estimation of major and trace element compositions of primary melts in a variety of tectonic settings, PRIMACALC2 is capable of calculating various important parameters related to source mantle conditions, like degree of melting and temperature and pressure of melting, which we consider an important proxy for correlating source mantle conditions with the observed geochemical diversity of the volcanic rocks in the Andaman ophiolite.

We could not get successful results for the entire dataset due to program specific limitations. Samples with low Ni (indicative of too much fractionation), or high Ni, due to Si-rich pyroxenite sources in arc basalts, resulting in high NiO in olivines, are outside the desired equilibrium field and hence excluded for further consideration. We had to exclude a few samples because they were deemed to be inappropriate for calculations in COMAGMAT3.72 (for details see Ariskin and Barmina, 2004, and references therein), a subsidiary program used in PRIMACALC2 for back-calculation. Another major problem is

the significant amount of loss on ignition (LOI) (e.g. >4 wt%) in a number of samples. Being a compilation, we do not have the scope to judge the individual analytical techniques and we speculate that such variable amounts of LOI are indicative of extensive alteration and hence could be a potential source of error in estimating source conditions. Considering all these limitations, instead of drawing first order conclusions we use the obtained results as one of the complementary models to supplement our geochemical observations (Fig. 8) and as a basis of further discussion. For this study we use mantle melting temperatures and degree of melting obtained for the volcanic rocks to check the consistency of the geochemical observation (reported in Fig. 8 and Fig. 9a) and further correlate it with degree of mantle melting obtained from peridotite geochemistry.

To obtain the degree of mantle melting directly from the peridotite chemical composition we have performed numerical melting experiments via the alphaMELTS program (Smith and Asimow, 2005) with Depleted MORB Mantle (DMM; Workman and Hart, 2005) as starting material. Results (clinopyroxene composition) of near-fractional isentropic melting of DMM (up to ~14%, which is found to be maximum extent of the melting in the numerical experiment performed here) are shown in primitive mantle-normalized (McDonough and Sun, 1995) and chondrite normalized (Sun and McDonough, 1989) multi-element spidergrams (Fig. 9b and c, respectively).

7 Discussion

7.1 *Contrasting geochemical characters of the Andaman ophiolite volcanic rocks – tectonic discrimination*

Owing to the different elemental property of Th (subduction-mobile) and Nb (subduction-immobile) in most arc systems Th/Nb is an ideal proxy for differentiating MORB lava from subduction-influenced ones (Pearce, 2008). Many of the analyses compiled from the literature do not include Th or Nb, therefore we interpret and compare our own samples with a restrained set of data. As suggested in Pearce (2014) we use this diagram in combination with the V/Ti diagram from Shervais (1982), which highlights the contribution of water to mantle melting. In the Th/Yb - Nb/Yb diagram (Pearce, 2008, 2014), pillow lavas from South Andaman (green circle, Jafri et al., 2010) and Bompoka Island (orange circle, Jafri and Sheikh, 2013) plot within the MORB-OIB array, having apparently been sourced from depleted MORB mantle (Fig. 8a). They also fall within the MORB or slab distal field of the V/Ti diagram (Fig. 8b), from which we infer that they represent basalts erupted from relatively shallow and dry melting of a depleted mantle source, corresponding to a mid-ocean ridge environment. Other reported analyses of “pillow basalt”, sometimes referred as “Upper Lavas” units from Rutland, lack Th or Nb analyses, and plot in the MORB field of V/Ti diagram (Fig. 8b). These include: red circle, Bhattacharya et al. (2013), South Andaman; grey circle, Pal 2011; brown circle, Jafri et al. (1990), based on location and authors’ interpretation; pink circle, Srivastava et al. (2004), based on location and authors’ interpretation; pale yellow diamond, Pedersen et al. (2010). The same characteristics also apply to basalts from Great Nicobar Islands (inverted pale blue triangle, Sachin et al., 2017).

Analyses of East Coast volcanics, also referred as “Lower Lavas”, have geochemical characteristics that differ from the Pillow Basalt unit. In the Th/Yb - Nb/Yb diagram, East

Coast volcanic rocks (yellow triangle, Ghosh et al., 2014) follow an intra-oceanic arc trend, parallel to the mantle array but significantly depleted in Nb (Fig. 8a). Other analyses compiled from the literature lack Nb and/or Th contents but were still plotted on the V-Ti diagram. Lower lava samples from Rutland (red triangle, Bhattacharya et al., 2013) and South Andaman (grey triangle, Pal, 2011; pink triangle, Srivastava et al., 2004, based on location and authors' interpretation) all straddle the Island Arc Tholeiite to MORB – slab distal fields. Interpreted together, such rocks likely resulted from melting of a re-fertilized depleted mantle in a water present environment, corresponding to an intra-oceanic arc or back-arc basin (Fig. 8b).

In the Th/Yb - Nb/Yb diagram, metamorphic sole samples (cyan star, except for AN1704a) plot near the mantle array, but with a slight slab component, as suggested by their negative Nb-Ta-Ti anomalies (Fig. 8a). They plot in the MORB-slab distal field of the V-Ti diagram (Fig. 8b). Considering the non-negligible Nb-Ta-Ti negative anomalies, we interpret the protolith of the metamorphic sole as a basaltic rock that resulted from partial melting of a depleted to enriched mantle source in the presence of low amounts of water and with a slight influence from a slab, which we interpret as a back-arc basin spreading centre. In the Th/Yb - Nb/Yb diagram, agglomerates from both South and Middle Andaman (neon red square) along with AN1704a (metamorphic sole) plot away from MORB-OIB array and thus clearly show subduction influence (Fig. 8a). South Andaman (AN1721a, AN1721b, AN1722b) and one Middle Andaman agglomerate (AN1711) samples show strong intra-oceanic arc signatures and appear to be sourced from an ultra-depleted mantle ($Nb/Yb < 1$). Among these samples, AN1721a (South Andaman) and AN1711 (Middle Andaman), plot in the Island Tholeiite field of the V-Ti diagram, consistent with melting in the presence of water (Fig. 8b). The geochemistry of these samples is akin to that of the East-Coast volcanics (reported by Ghosh et al., 2017, yellow triangle).

In the Ce/Nb - Th/Nb diagram (Saunders and Tarney, 1991), back-arc basin basalts tend to fall in the intermediate zone between the low Th/Nb zone (MORB-OIB) and high Th/Nb (Island Arc basalts) zone (Fig. 8c). Pillow lavas from South Andaman (green circle) have relatively high Ce/Nb and low Th/Nb zone and plot near the N-MORB domain. Agglomerates and East Coast volcanics (yellow triangle) are characterized by high Ce/Nb and high Th/Nb typical of arc volcanics. Interestingly, sole samples plot in the intermediate zone and near to the back-arc basin basalts from the Mariana Trough (Fig. 8c). A similar intermediate Ce/Nb and Th/Nb ratio in pillow basalts from Bompoka Island (orange circle) indicates a back-arc origin. In the La/Nb - Y diagram (Floyd et al., 1991) sole samples along with pillow basalts from Bompoka Island fit in the BABB (back-arc basin basalts) field (Fig. 8d). Agglomerates and East Coast volcanics (Ghosh et al., 2014) are characterized by higher La/Nb ratio and mostly plot in fields of subduction influenced basalts – forearc platform basalts (FAPB) and IAT. All previous diagrams suggest a N-MORB affinity for pillow lavas from South Andaman (green circle), however their origin remains inconclusive in the La/Nb – Y diagram. Saccani (2015) proposed a new discrimination diagram using absolute measures of Th and Nb based on >2000 known ophiolitic basalts and ~560 modern rocks from known tectonic settings. In the Th_N - Nb_N diagram (Fig. 8e; Nb and Th normalized with respect to N-MORB) we can see a clear distinction between two groups of volcanic rocks of the Andaman Islands. Pillow lavas from South Andaman (green circle), Bompoka Island (orange circle) and sole samples plot in the domain of no subduction component. Contrastingly, agglomerates, AN1704a (sole) and East Coast volcanics (yellow triangle) are in the subduction-influenced domain. Furthermore, the BABB field overlaps with various volcanic rocks (IAT, CAB, N-MORB, E-MORB) and is further subdivided as back-arc A and back-arc B domains. Back-arc A indicates BABB characterized by input of subduction or crustal components (e.g. immature intra-oceanic or ensialic back-arc), whereas back-arc B indicates

BABB with no input of subduction or crustal components (e.g. mature intra-oceanic back-arc). Hence, from Fig. 8e we can conclude that agglomerates, AN1704a (sole) and East Coast volcanics (Ghosh et al., 2014) probably formed in an arc – arc proximal back-arc environment whereas sole samples, along with pillow basalts from Bompoka Island (Jafri and Sheikh, 2013) with MORB affinity, represent volcanics of a slab distal back-arc setting.

The sharp geochemical contrast of the two groups of volcanic rocks has been clearly distinguished in these diagrams and further substantiated by the multielement N-MORB and chondrite (Sun and McDonough, 1989) normalized spidergram. In most cases the full suite of trace and rare-earth elements is missing, but wherever available the data are plotted (Supplementary fig. 2a-r). We found three datasets of volcanic rocks with the nearly complete spectrum as summarized in Fig 8f-i. In the N-MORB normalized diagram, sole samples (except AN1704a) plot very close to N-MORB (especially in REEs) and display a very similar character to that of pillow lavas from South Andaman and Bompoka Island. However, the slight negative Nb, Ta, Hf, Zr and Ti anomalies of the metamorphic sole are almost absent in South Andaman pillow basalts and very subtle in Bompoka pillow basalts (Fig. 8f). The variable Nb-Ta-Ti anomalies, ranging from obvious (AN1704a) to negligible (AN1709a), point towards a back-arc setting with variable slab influence whereas pillow basalts are expected to form by a high degree of partial melting of a depleted mantle source at shallow depth, typical of mid ocean ridge. Owing to the presence of a slight Ti anomaly in the pillow basalts the possibility of the presence of a slab distal component (BABB) cannot be ruled out. The chondrite normalized diagram (Fig. 8g) for the same group of samples shows very flat REE patterns. Agglomerate samples (and AN1704a) show a contrasting geochemical behaviour to that of the sole samples (discussed in the section 5.1.2) and are near identical with the East Coast volcanics (Ghosh et al., 2014) except that the later show Ta enrichment with respect to Nb (Fig. 8h,i). Sole samples (except AN1704a), pillow lavas from

South Andaman and Bompoka Island have been previously seen to have a geochemical character very close to N-MORB (Fig. 8a,f,g), however, in the large ion lithophile element (LILE) bearing N-MORB normalized spidergram all three datasets - sole samples, pillow lavas from South Andaman and Bompoka Island - display a significant enrichment of Rb, Ba, K, Sr (Fig. 8j). Enrichment of LILE in back-arc basin basalts has been previously documented by Saunders and Tarney (1991). Such LILE enrichment of pillow basalts from South Andaman and Bompoka Island has been proposed to be indicative of back-arc basin basalt affinity (Jafri and Sheikh, 2013; Jafri et al., 2010). A similar inference could be made for the metamorphic sole, but the robustness of such an interpretation could be questioned since trace elements in these rocks, especially the LILE, are prone to have been significantly modified during metamorphism (Ishikawa et al., 2005).

The results obtained by PRIMACALC2 (Fig. 9a) also clearly demonstrate that the volcanics with arc affinity, agglomerates, East Coast volcanics (Ray et al., 1988; Ghosh et al., 2014), and a few samples from Great Nicobar (Sachin et al., 2017), lower lava (Pal, 2011) are product of higher (14-21%) degrees of melting than those that have MORB (or BABB) affinity (3-10%) – sole samples, pillow basalts from South Andaman (Jafri et al., 1990, 2010) and Bompoka Island (Jafri and Sheikh, 2013). Samples from Corbyn's Cove in Pedersen et al. (2010) show a lower degree of melting than those collected in Chidiya Tapu. The one to one correlation of geochemical diversity and degree of mantle melting could be attributed to the two modes of melting - flux induced (higher degree) melting beneath an arc and decompression (lower degree of melting) beneath an extensional setting (e.g. back-arc).

7.2 *Contrasting petrological characters of the Andaman ophiolite mantle rocks*

Along the stretch of the Andaman islands there is a pronounced variation in terms of petrography and mineral chemistry within the ophiolitic mantle sequence. The restitic

peridotite with forearc affinity of Rutland Island is composed of depleted harzburgite to clinopyroxene-bearing harzburgite whereas Middle and North Andaman are dominated by less depleted lherzolite akin to the geochemical signature of abyssal peridotites (Ghosh et al., 2013). The degrees of melting calculated from chrome spinel chemistry indicate low to moderate values (2–10%) for North and Middle Andaman samples, and high values (14–18%) for Rutland peridotites (Ghosh et al., 2018), which corroborate well the degree of melting calculated for the volcanic rocks (pale green and purple boxes in Fig. 9a). Less incompatible trace elements in clinopyroxenes, HREEs in particular, serve as an important proxy for the extent of partial melting since the mineral contains >85% of the trace element budget of the peridotites (Warren et al., 2009). Keeping this in mind, we compare our numerical melting results with the Andaman peridotite clinopyroxene data from Ghosh et al. (2018) (Fig. 9b,c). The chondrite REE spidergram (Fig. 9c) indicates 3-10% melting of DMM for the North and Middle Andaman peridotites and ~14% melting for peridotites of Rutland Island, which is in accord with the conclusions of Ghosh et al. (2018). Similar inferences may be drawn from the primitive mantle normalized spidergram (Fig. 9b), especially comparing the HREE pattern.

We have compiled and plotted clinopyroxene compositions of peridotites of the Southern Mariana forearc (Supplementary fig. 3.a,e; Chen and Zeng, 2007), Izu-Bonin-Mariana forearc (Supplementary fig. 3.b,f; Parkinson and Pearce, 1998), Tonga forearc (Supplementary fig. 3c,g; Birner et al., 2017) and supra-subduction zone (SSZ) peridotites from ophiolite complexes of SW Turkey (Supplementary fig. 3.d,h; Aldanmaz, 2012) to compare them with peridotites from Rutland Island. In case of the Southern Mariana forearc mantle Chen and Zeng (2007) inferred that highly refractory harzburgites are a product of subduction-related mantle metasomatism via hydrous melt. Parkinson and Pearce (1998) interpreted Izu-Bonin forearc peridotites as a result of interaction of a harzburgitic residue

with an oxidized boninitic melt in an SSZ setting. However, reassessment of Birner et al. (2017) suggests that enrichments in LREE in the samples could be better explained by disequilibrium melting than by melt addition (interaction with boninitic melt). For Tonga forearc peridotites, two distinct groups have been identified by Birner et al. (2017) based on spinel Cr#, fO_2 , and trace element concentrations. Group I is interpreted as residues of large degrees of fractional melting, with minor influence from arc-like melts or fluids, formed during the first stages of subduction initiation whereas Group II is inferred to have formed during flux melting of fertile mantle owing to interaction with subduction-related fluids and/or melts. We find a good match with HREEs of all the above mentioned forearc (SSZ influenced) peridotites, except Group II of the Tonga forearc peridotites. We would also like to point out the close resemblance of Rutland samples with the recently recovered peridotite fragments from Daito Ridge, which is interpreted to be the mantle section of a remnant proto-Philippine Sea island arc (Supplementary fig. 3i,j; Morishita et al., 2018b). The clinopyroxene compositions of compiled forearc peridotites are shown for reference in supplementary fig. 3i,j and also in Fig. 9b,c. Owing to the lack of LREE data of the Rutland peridotites we could not satisfactorily compare the behaviour of such elements. We note one major difference in the primitive mantle normalized multielement pattern from the results of melting experiments, i.e. the enrichment of Nb with respect to Ce in the peridotite samples (Fig. 9b). This behaviour is very prominent in Rutland samples; also, in Tonga Group I peridotites (Supplementary fig. 3c,g) and Daito Ridge peridotites (Supplementary fig. 3i,j). Such Nb enrichment could have resulted from the melt mantle interaction in the upper mantle beneath the subduction zone (Kelemen et al., 1993; Stolz et al., 1996). This line of evidence thus confirms the refractory nature of the Rutland peridotite to have formed in a subduction influenced (arc/forearc) environment as suggested by Ghosh et al. (2013, 2018). Further, we can conclude that peridotites like Tonga forearc Group II, which are indicative of a flux

melting dominated mature stages of subduction, were either not formed, or are not exposed in the Andaman ophiolite. Ghosh et al. (2018) also documented a close resemblance of the North and Middle Andaman peridotite data with back-arc basin peridotite data from the Mariana Trough (Ohara et al., 2002) and Parece Vela back-arc basin (Ohara et al., 2003), and assigned a back-arc origin to them (shown in Fig. 9b,c). Morishita et al. (2018a) also interpreted the peridotites of Middle Andaman to be the result of low-degree of partial melting with minor flux infiltration, which is likely to take place in a back-arc region. González-Jiménez et al. (2020) recently estimated the parental melt composition for the chromitite pods hosted within the North Andaman lherzolites, which is found to be similar to high-Mg island arc tholeiite (IAT) and boninites (; and broadly analogous with the parental melt of chromitites from Rutland Island that inferred to be formed at supra-subduction zone setting (Ghosh et al., 2009). The back-arc geochemical signature of the lherzolic peridotites (Ghosh et al., 2018), coupled with the commonly believed melt-mantle interaction origin for podiform chromitites (Arai, 1997; Zhou et al., 1994), thus suggest that the North Andaman lherzolites, after having formed in the back-arc setting, must have been located in the forearc region, where they interacted with arc affinity melt and eventually formed chromitite pods.

7.3 *Cretaceous geodynamic evolution of the Sunda subduction zone*

Reviewing the previous geochemical and geochronological investigation, Ghosh et al. (2017) explained the formation of the Andaman ophiolite entirely within the context of a single progressive subduction system. However, as pointed out by Bandyopadhyay et al. (2020), this does not adequately explain all our field observations and geochronological constraints. For instance, the 93-95 Ma U-Pb ages for the plagiogranites of South Andaman were inferred to date ophiolite spreading and subduction initiation (Pedersen et al., 2010). However, the 93.6 Ma (Sarma et al., 2010) plagiogranite-diorite suite from location 3

(Bedonabad coast, South Andaman) has an arc affinity, implying subduction maturation, and has been proposed to be genetically linked with the East Coast volcanics (Jafri et al., 1995). So, the meaning of this 93- 95 Ma age is intriguing. The problem becomes obvious when compared with the age of ~106 Ma estimated from plagioclase xenocryst within recently erupted Barren Island volcanic rock (Ray et al., 2015) and the 105-106 Ma cooling age from Andaman ophiolite metamorphic sole (Plunder et al., 2020).

Kinematic restoration of the Andaman Sea back to pre-Oligocene time shifts the Andaman ophiolites to the forearc of Sumatra, immediately west of the Cretaceous Woyla Arc (van Hinsbergen et al., 2011; Advokaat et al., 2018), which collided with and accreted to West Sumatra in the mid Cretaceous (Barber et al., 2005). It has been speculated that this collision supposedly flipped subduction polarity (Advokaat et al., 2018; Plunder et al., 2020). We now integrate field observations and geochemical and geochronological data from the Andaman Ophiolites to test whether the previously speculated polarity reversal could be a possible option to initiate the Sunda subduction system in the west of Sumatra during the Cretaceous.

7.3.1 99-93 Ma arc magmatism

Here, we interpret the age of 98.5 Ma of the younger zircon in layered gabbro (SA-GAB1) from the nearby location (location 2; Kodiaghat section, South Andaman) as dating the emplacement (Figs. 7,10). These gabbros are interpreted to be genetically linked with the Moho transition zone rocks (olivine-rich troctolite, and clinopyroxenite-wehrlite) at the same location. Olivine-rich troctolite is interpreted to have an origin by impregnation of hydrous basaltic melt into a dunite (replacive) protolith that formed earlier by melt-mantle interaction (Ghosh et al., 2014). The composition of chrome spinels from this olivine-rich troctolite shows a clear affinity to the harzburgitic peridotites of Rutland Island (Ghosh et al., 2014).

These peridotites possess forearc geochemical affinity (Ghosh et al., 2013) and are found to have suffered relatively higher degree (~14%) of melting (Fig. 9; Ghosh et al., 2018). The extent of melting that the Rutland peridotites suffered is in concurrence with the degree of melting determined for the East Coast volcanics (Fig. 9a). Ghosh et al. (2014) also showed that melt compositions in equilibrium with clinopyroxene from olivine-rich troctolites and gabbroic rocks fit with the East Coast volcanics. Now we have also found the East Coast volcanics data (Ghosh et al., 2014) to have an arc affinity (Fig. 8). In addition, plagiogranite (148A) from Rutland Island (location 4) gives a 98.7 Ma U/Pb age (Fig. 7). So, arc lithosphere with volcanic rocks (East Coast volcanics), plagiogranite, layered gabbro, lower crustal cumulate and harzburgite had been formed at least by ~99 Ma. Specifically, lithotypes exposed in Bedonabad coast, South Andaman (location 3), thus appear to have formed in an arc setting, where pre-existing gabbroic crust was intruded by arc volcanics and plagiogranites around 93 Ma (Jafri et al., 1995; Sarma et al., 2010). Jafri et al. (1995) presumed that the Andaman plagiogranite, may not be a part of the Andaman ophiolite sequence and may post-date emplacement of the Andaman ophiolite gabbro. Our new geochronological and geochemical results are in accord with this inference.

Agglomerates from Panchawati coast, Middle Andaman (Location 1) and Chidiya Tapu coast, South Andaman (Location 6) unequivocally show arc characteristics (Fig. 8). Interestingly, agglomerates from Chidiya Tapu coast, South Andaman (Location 1) are found to host plagiogranite blocks that presumably have the same age (i.e. 95 ± 2 Ma) as the trondhjemite sample dated by Pedersen et al. (2010) from the same location. Thus agglomerates would have formed during an explosive arc volcanism not earlier than ~95 Ma. The likely cogenetic leucocratic rock L-18 (belonging to the diorite – granodiorite series), associated with Middle Andaman agglomerates, yields a ~93.5 Ma age (Fig. 7). Late Cretaceous explosive arc volcanism has been also predicted from the presence of ash layers

within the radiolarian cherts in South Andaman (Jafri et al., 2006). Accordingly, based on our results from Middle Andaman, South Andaman and Rutland Island, we postulate that the bulk of magmatic rocks, including gabbros, plagiogranites, and agglomerates found on the Andaman Islands formed in a volcanic arc that had developed by 98-99 Ma and remained active until at least ~93 Ma (Fig. 10).

For supra subduction zone magmatism, a tendency of increasing alkalinity away from the trench has been documented (Wilson, 1989, and references therein) and is thought to reflect cooling of the mantle wedge and the progressive deepening of the melting upon maturation of subduction (Stern et al., 2012; Whattam et al., 2011). Thus, the process is expected to yield tholeiitic volcanics in the forearc during subduction zone infancy, followed by gradually more calc-alkaline volcanics. We have tested the alkalinity in the volcanic samples of the Andaman ophiolite with major element based AFM (Irvine and Baragar, 1971) and trace element-based (Ross and Bédard, 2009) discrimination diagrams (supplementary fig. 4) and found that the subduction influenced volcanic rocks mostly plot in the tholeiitic to transitional domain with few in calc-alkaline field. This might suggest that agglomerates of South and Middle Andaman were erupted not very far from the trench in the early stages of arc maturation.

7.3.2 Subduction initiation at or before 105 Ma

Though arc characteristics are preserved in most of the crustal rocks (except pillow basalts) and also in Rutland peridotites, the geochemical signatures of Middle Andaman peridotites do not comply with this feature. Ghosh et al. (2013) reported less depleted lherzolitic mantle from Middle Andaman that shows geochemical signatures of abyssal peridotite. These peridotites have been estimated to have suffered relatively low degrees (3-10%) of melting compared to the Rutland peridotites (Ghosh et al., 2018; Fig. 9). The

geochemistry of the Middle Andaman peridotites is better explained by depletion in a back-arc setting (Fig. 9b,c; Ghosh et al., 2018). The magmatic precursor rocks of the sole appear to be the product of ~7% mantle melting, which fits well within the estimated extent of melting for the Middle Andaman peridotites below which the sole rocks accreted (see Plunder et al., 2020). Geochemically, metamorphic sole amphibolites (reported by Plunder et al., 2020) also show slab-distal, back-arc basin signatures, with the exception of sample AN1704a (Fig. 8) that yielded an arc signature. In summary, metamorphic sole rocks (originally part of the nascent downgoing slab) and nearby mantle rocks (of the upper plate) (Fig. 1c) both possess back-arc characteristics (Fig. 8,9). This suggests that subduction may have started in a back-arc basin, but not far from the arc. Here, we would like to focus on the $^{40}\text{Ar}/^{39}\text{Ar}$ ages of 106.4 ± 2.1 Ma and 105.3 ± 1.6 Ma from two amphibolite samples of the metamorphic sole of Middle Andaman (Plunder et al., 2020). Typically, the $^{40}\text{Ar}/^{39}\text{Ar}$ cooling ages of metamorphic soles coincide with the timing of spreading of SSZ ophiolites, which may suggest that this spreading caused the exhumation of the underlying sole (van Hinsbergen et al., 2015, and references therein). The inherited U/Pb age (~105 Ma) found in one of our magmatic samples (SA-GAB1) suggests crystallization of magmatic zircon at this time, as does the ~106 Ma age found in a xenocrystic plagioclase feldspar in a recent Barren Island volcanic rock (Ray et al., 2015). It has long been thought that $^{40}\text{Ar}/^{39}\text{Ar}$ ages reflect metamorphic sole formation, and that this in combination with synchronous SSZ spreading marks the timing of subduction initiation (e.g. Maffione et al., 2015; Whattam et al., 2011; Stern and Bloomer, 1992). However, recent dating of garnet growth within metamorphic soles of Oman and Turkey with Lu/Hf geochronology has shown that sole growth may predate sole cooling and SSZ ophiolite spreading by >10 Myr (Guilmette et al., 2018; Pourteau et al., 2019). From these arguments, in combination with field observation, geochemical evidences and geochronological constraints we conclude that in the Andaman

region subduction initiated at or before 105-106 Ma, in an oceanic back-arc basin environment (Fig. 11). We again emphasize that the cooling age of metamorphic sole, the inherited age of a plagioclase feldspar xenocryst and the inherited U-Pb age of gabbro are closely bracketed within 105-106 Ma, thus reflecting only a minimum age of subduction initiation. The 99-93 Ma arc magmatic rocks thus post-date sole exhumation, and by inference SSZ ophiolite spreading, by 7-13 Myr (Fig. 10), a time delay between SSZ spreading and arc maturation similar to the well-documented case of the Izu-Bonin-Mariana subduction zone (Stern et al., 2012).

7.3.3 *Matching the geochemistry with tectonic evolution*

Following the above discussion, we here try to assess the tectonic reconstruction reversal, which is suggested to have initiated new subduction in the region (Advokaat et al., 2018; Plunder et al., 2020). Further we will also attempt to reconstruct the stage-wise subduction evolution in Andaman in the regional geodynamic context (Fig. 11).

Stage 1 (before 105 Ma, Fig. 11a): From the regional geodynamic perspective, formation of the Woyla Arc has been suggested to be due to westward subduction that occurred simultaneously with (north)eastward subduction below Sundaland (Eurasian plate) from ~130 to ~100-90 Ma (Barber et al., 2005; Advokaat et al., 2018). From plate kinematic and paleomagnetic analyses, Advokaat et al. (2018) postulated that a double verging subduction (of the now-subducted 'Ngalau' oceanic plate, of which remnants of Upper Triassic limestone (Wajzer et al., 1991) and Triassic and Middle Jurassic radiolarian chert (McCarthy et al., 2001; Barber et al., 2005; Munasi and Putra, 2019) are found in the Ngalau Suture on Sumatra), with two trenches bounding the Woyla Arc and Sundaland, was active in this time period. Collision of the Woyla Arc with Sundaland (Eurasian plate) led to southward migration of the trench-trench-trench triple junction between Woyla, Sundaland, and Ngalau,

requiring a subduction polarity reversal to maintain a stable triple junction (Advokaat et al., 2018). This reversal led to subduction initiation in the back-arc behind the Woyla Arc, explaining the geochemical signatures of the metamorphic sole rocks (location 1) and of the Andaman lherzolites.

Stage 2 (~105 Ma, Fig. 11b): By 105 Ma, a metamorphic sole had formed and was exhuming, presumably simultaneously with SSZ spreading (Plunder et al., 2020). We interpret synchronous magmatism in the upper plate, suggested by the Barren Island 106 Ma xenocryst and the inherited ~105 Ma zircon age documented in this paper, as the reflection of SSZ ophiolite spreading. During this stage, Woyla Arc-Sundaland continent collision may not have been finalized yet, as in the modern equivalent of such a setting found in Timor and Wetar. There, GPS data (Nugroho and Harris, 2009) show that approximately 30% of Australia-Banda convergence is still accommodated between Timor and Australia, where the Banda arc has been colliding with Australia since the late Miocene (Audley-Charles et al., 1979; Tate et al., 2014, 2015, 2017), and the remainder of convergence is accommodated by the Wetar back-thrust that has formed within the last few million years at the northern edge of the Banda arc (Breen et al., 1989).

Stage 3 (<99-93 Ma, Fig. 11c): Newly formed, NE verging subduction evolved to a more mature stage at around 99 Ma and was characterized by cooling of the mantle wedge and the associated deepening of the melting zone. Our new magmatic U-Pb age data of ~98-99 Ma show that new arc volcanoes formed within 7-13 Myr of sole exhumation. This newly formed explosive arc volcano was active at least until 93 Ma and produced most of the exposed rock types in the Andaman islands.

8 Conclusions

We have performed a geochemical and geochronological analysis of magmatic rocks of the Andaman Ophiolites to evaluate the feasibility of a previously speculated process of subduction initiation via polarity reversal. We find that there are two distinct groups of ophiolitic rocks in the Andaman Island. Mantle peridotites as well as mafic amphibolites of the metamorphic sole below the ophiolite have chemical signatures suggestive of an origin in a back-arc basin. Volcanic agglomerates, including plagiogranite xenoliths, suggest formation in a volcanic arc. New U/Pb ages show that arc magmatism occurred between ~99 and 93 Ma, at least 7 Myr later than the cooling of the sub-ophiolitic metamorphic sole that gave ~106-105 Ma hornblende $^{40}\text{Ar}/^{39}\text{Ar}$ ages. The origin of pillow lava sequences remains ambiguous due to the lack of age constraints and it is a topic of future research. However, their MORB geochemical affinity, with a speculated back-arc basin (slab distal) origin, may suggest that they could have originated at any stages of subduction, where ocean floor spreading was taking place with no or little influence from slab.

We interpret the geochemistry of the Andaman Islands as follows: Restoring the Andaman ophiolites back to the Early Oligocene onset of opening of the Andaman Sea basin brings the ophiolites to the west of the Woyla Arc. The latter had collided with Sumatra around 105-95 Ma. We find that the Andaman ophiolites resulted from a subduction polarity reversal behind an arc (the Woyla Arc), after which SSZ spreading around 105 Ma led to metamorphic sole exhumation. By 99-93 Ma a volcanic arc had developed on the Andaman Islands. This proposed mechanism satisfactorily explains the field relationships of the geochemically incoherent ophiolitic rock record in the Andaman Islands. Such a petrochemical manifestation of subduction polarity reversal in rock records has not been described in detail before.

996

997 **Acknowledgments**

998 This work is part of the doctoral research (PhD) program of the first author, supported
999 with DST INSPIRE Fellowship (IF 130148). D.B specially acknowledges the help received
1000 from Dr. Jun- Ichi Kimura (JAMSTEC, Japan) regarding PRIMACALC2 calculation. B.G.
1001 acknowledges financial support received from Science and Engineering Research Board,
1002 DST, India (EMR/2017/000929). D.J.J.v.H., A.P., and E.L.A. were funded through ERC
1003 starting grant SINK (306810) to D.J.J.v.H. D.J.J.v.H. acknowledges NWO Vici grant
1004 865.17.001. The authors thank two anonymous reviewers for their insightful comments and
1005 suggestions. We also acknowledge Michael Jentzer for his comments on an earlier version of
1006 the manuscript. Editorial handling by Greg Shellnutt is greatly acknowledged. We thank the
1007 Ramakrishna mission in Port-Blair for accommodation and Yael Engbers for help during the
1008 fieldwork. All data including supporting information for this manuscript are available at
1009 Figshare <https://doi.org/10.6084/m9.figshare.11396469.v7>.

Figure captions

Figure 1 Tectonic and geological framework of the Andaman ophiolite - (a) Colour-shaded relief image showing the eastern Himalayan syntaxis, the western Burma (Myanmar) region and the Andaman–Sumatra–Java (ASZ) trench in the regional geodynamic framework (modified after Bandyopadhyay et al., 2020). Major tectonic features are redrawn from Awasthi (2017), Imsong et al. (2016) and Liu et al. (2016b). Regions demarcating the Woyla Arc, West Sumatra, are from Advokaat et al. (2018) and West Burma block from van Hinsbergen et al. (2011). Red box demarcates the Andaman Islands and Rutland Island (detailed in Fig. 1b); previous geochronological results are also shown (see text for details). Relief data are from ETOPO1 Global Relief Model (<http://www.ngdc.noaa.gov/mgg/global/>); (b) Geological map of Andaman Islands (modified after Ghosh et al., 2017) over processed-hillshade map (generated using SRTM digital elevation data with a resolution of 1 arc-second from USGS Earth Explorer <https://earthexplorer.usgs.gov/>; modified after Bandyopadhyay et al., 2020). North-South extending Jarwa Thrust is also shown. STDS: South Tibetan Detachment System, MCT: Main Central Thrust, MBT: Main Boundary Thrust, MFT: Main Frontal Thrust; LA: Little Andaman; CN: Car Nicobar; Nic. I: Nicobar Islands; GN: Great Nicobar; EMF: Eastern Margin Fault; DF: Diligent Fault; WAF: West Andaman Fault; BI: Barren Island; NI: Narcondam Island; AR: Alcock Rise; SR: Sewell Rise; WOB: Western Ophiolite Belt; MOB: Myitkyina Ophiolite Belt. Star marks indicate important field locations of Middle and South Andaman and Rutland Island that are discussed in this study (sample locations are marked by green stars, see text for details); (c) Google Earth satellite image showing the salient locations along the Panchawati coast (Middle Andaman).

Figure 2 Schematic plate kinematic reconstruction of the east Indian Ocean from the Cretaceous to the present, with the predicted location of Andaman, modified from Advokaat et al. (2018).

Figure 3 Field disposition of different magmatic rocks in the Andaman Islands - (a) Rock types exposed in the MOHO Transition zone (MTZ) in Kadiaghat, South Andaman (Location 2, see Ghosh et al. 2014 for details); (b) In-situ exposure of layered gabbro in South Andaman, in close proximity of the MTZ location (location 2); (c) Plagiogranite exposure on the Bedonabad coast, South Andaman (location 3), Note the intermingling of mafic rocks with the plagiogranites; (d) In-situ exposure of leucocratic rocks (belonging to the diorite – granodiorite series) in close association with agglomerates on the Panchawati coast, Middle Andaman (location 1); (e) Plagiogranite exposure in Rutland Island (location 4); (f) Pillow lava outcrop at Corbyn’s Cove beach, South Andaman (location 5); (g) Coastal exposure (near Mundapahar beach) of agglomerates in Chidiya Tapu, South Andaman (location 6). Note the predominant occurrence of felsic blocks of varied size; (h, i) Coastal exposure of agglomerates on the Panchawati coast, Middle Andaman (location 1).

Figure 4 Optical photomicrographs showing characteristic petrographic features – (a) plagiogranite (148A), note plagioclase grains are often altered; (b) plagiogranite (BD-1) showing interstitial vermicular and micrographic intergrowths of quartz and plagioclase; (c) layered gabbro (SA-GAB1) composed of olivine, plagioclase, clinopyroxene and showing cumulate texture; (d) cumulate clinopyroxenite (SA-GAB2) with interstitial plagioclase; (e) plagiogranite clast from the agglomerates (CH-3) with hypidiomorphic granular texture, note the occurrence of epidote; (f-k) agglomerates from South Andaman that contains varieties of clasts (e.g. plagiogranite, andesite, epidote-rich clasts etc.) in fine grained volcanic matrix; (l) leucocratic rock belonging to the diorite – granodiorite series (L-18) showing hypidiomorphic granular texture, composed of plagioclase and amphibole (locally replaced by chlorite); (m-p) volcanic rocks (agglomerates) from Middle Andaman, dominantly showing porphyritic texture. Besides plagioclase there are phenocrysts of another phase, possibly clinopyroxene, replaced by chlorite and carbonate (See Table 1 for details).

Figure 5 Bivariate plots (major and minor element based) of geochemical data from Middle and South Andaman volcanic rocks and amphibolites (Metamorphic sole) – (a) $\text{SiO}_2 - \text{TiO}_2$; (b) $\text{SiO}_2 - \text{Al}_2\text{O}_3$; (c) $\text{SiO}_2 - \text{CaO}$; (d) $\text{SiO}_2 - \text{Mg\#}$; (e) $\text{SiO}_2 - \text{Cr}$; (f) $\text{SiO}_2 - \text{Ni}$; (g) $\text{SiO}_2/\text{Al}_2\text{O}_3 - \text{Mg\#}$ (after Kempton and Harmon, 1992); All data have been plotted on an anhydrous basis (Table 2). Compiled data of volcanic rocks (see Supplementary table 2) from Andaman and neighbouring islands are shown as transparent overlays in background, to show the entire spectrum (for symbols see Fig. 8). The grey shades indicate the ultrabasic, basic, intermediate and acid boundaries (after Le Bas et al., 1986, Fig. 2)

Figure 6 Trace and REE geochemistry of Middle and South Andaman volcanic rocks and amphibolites (metamorphic sole). (a) Immobility element-based TAS (Total Alkali-Silica) proxy diagram (from Pearce, 2014, after Floyd and Winchester, 1975); (b-f) N-MORB and chondrite (Sun and McDonough, 1989) normalized spidergram, (b,e) amphibolites (Middle Andaman), (c,f) agglomerates (Middle Andaman) and (d,g) agglomerates (South Andaman).

Figure 7 Geochronological results - Concordia diagram for zircons recovered from sample SA-GAB1, 148A and L-18. Data point error ellipses are 2σ . See table 3.

Figure 8 Tectonic discrimination of volcanic rocks of Andaman ophiolite: (a) $\text{Th/Yb} - \text{Nb/Yb}$ diagram (Pearce, 2008, 2014); (b) $\text{V} - \text{Ti}$ diagram (Shervais, 1982); (c) $\text{Ce/Nb} - \text{Th/Nb}$ diagram (Saunders and Tarney, 1991); (d) $\text{La/Nb} - \text{Y}$ diagram (Floyd et al., 1991), (e) $\text{Th}_N - \text{Nb}_N$ diagram (Saccani, 2015). N-MORB (f,h); (j) with emphasis on LILE, (g,i) chondrite normalized spidergram for volcanic rocks of Andaman ophiolite. OIB: ocean-island basalt; MORB: mid-ocean ridge basalt, N-: normal type, E-: enriched type, P-: plume type, D-: depleted type; MTB: medium-Ti basalt; IAT: island arc tholeiites; BABB: back-arc basin basalts; OFB: oceanic flood basalt; FAPB: forearc platform basalt; AB: alkaline basalt; CAB: calc-alkaline basalt. Data source with colour codes: red: Bhattacharya et al. (2013); yellow: Ghosh et al. (2014); orange: Jafri and Sheikh (2013); green: Jafri et al. (2010); brown: Jafri et

al. (1990); blue: Ray et al. (1988); grey: Pal (2011); pink: Srivastava et al. (2004); light purple: Sachin et al. (2017); pale yellow: Pedersen et al. (2010); neon red: agglomerates (South and Middle Andaman, present study); cyan: metamorphic sole (present study). Details of data source and meaning of symbols are also given in the diagram.

Figure 9 (a) Source mantle conditions of volcanic rocks of Andaman ophiolite - Degree of melting vs Temperature diagrams (calculated using PRIMACALC2; Kimura and Ariskin, 2014). Degree of melting calculated from chrome spinel (Ghosh et al., 2018) is also shown. Pale green band: degree of melting from $Cr\#_{\text{spinel}}$ in peridotites of Rutland; purple band: degree of melting from $Cr\#_{\text{spinel}}$ in peridotites of North and Middle Andaman; symbols are the same as in Fig. 8; (b) primitive mantle; (c) chondrite normalized plot REE patterns of clinopyroxene of the DMM (Workman and Hart, 2005) after melting calculation (using alphaMELTS program, Smith and Asimow, 2005, see text for details, degree of melting are shown in %). Clinopyroxene trace element patterns of Rutland, North and Middle Andaman peridotite (Ghosh et al., 2018) along with the domain of back-arc peridotite (Ohara et al., 2002, 2003) and forearc peridotites (compiled in this study, see supplementary figure 3) are also shown. Owing to extremely low values (normalized) for Nb (in Fig. 9b) and La (in Fig. 9c), numeric values are mentioned in the respective diagrams (original in parentheses).

Figure 10 Geochronological summary of U-Pb zircon ages of magmatic rocks along with Ar-Ar ages from metamorphic sole (Plunder et al., 2020) of Andaman ophiolite; Ar-Ar age from plagioclase xenocryst hosted in Barren lava (Ray et al., 2015) is also shown for reference.

Figure 11 Schematic reconstruction of Cretaceous subduction initiation and evolution of Andaman ophiolite (see text for details).

1107 **Table captions**

1108 **Table 1.** Geographic location and petrographic summary of the studied samples of Andaman
1109 ophiolites

1110 **Table 2.** Bulk rock geochemistry of selected samples of volcanic rocks from Andaman
1111 ophiolites

1112 **Table 3.** U-Pb data for zircon from Andaman samples

References

- Acharyya, S.K., 2007. Collisional emplacement history of the Naga-Andaman ophiolites and the position of the eastern Indian suture. *Journal of Asian Earth Sciences* 29, 229-242.
- Acharyya, S.K., Ray, K.K., Sengupta, S., 1990. Tectonics of the ophiolite belt from Naga Hills and Andaman Islands, India. *Proceedings of the Indian Academy of Sciences- Earth Planetary Sciences* 99, 99-187.
- Advokaat, E.L., Bongers, M.L.M., Rudyawan, A., BouDagher-Fadel, M.K., Langereis, C.G., van Hinsbergen, D.J.J., 2018. Early Cretaceous origin of the Woyla Arc (Sumatra, Indonesia) on the Australian plate. *Earth and Planetary Science Letters* 498, 348-361.
- Agard, P., Yamato, P., Soret, M., Prigent, C., Guillot, S., Plunder, A., Dubacq, B., Chauvet, A., Monié, P., 2016. Plate interface rheological switches during subduction infancy: Control on slab penetration and metamorphic sole formation. *Earth and Planetary Science Letters* 451, 208-220.
- Aitchison, J.C., Ao, A., Bhowmik, S., Clarke, G.L., Ireland, T.R., Kachovich, S., Lokho, K., Stojanovic, D., Roeder, T., Truscott, N., Zhen, Y., Zhou, R., 2019. Tectonic Evolution of the Western Margin of the Burma Microplate Based on New Fossil and Radiometric Age Constraints. *Tectonics* 38, 1718-1741.
- Aldanmaz, E., 2012. Trace element geochemistry of primary mantle minerals in spinel-peridotites from polygenetic MOR–SSZ suites of SW Turkey: constraints from an LA-ICP-MS study and implications for mantle metasomatism. *Geological Journal* 47, 59-76.
- Allen, R., Carter, A., Najman, Y., Bandopadhyay, P.C., Chapman, H.J., Bickle, M.J., Garzanti, E., Vezzoli, G., Andò, S., Foster, G.L., Gerring, C., 2008. New constraints on the sedimentation and uplift history of the Andaman-Nicobar accretionary prism, South

1137 Andaman Island, in: Draut, A.E., Clift, P.D., Scholl, D.W. (Eds.), Formation and
 1138 Applications of the Sedimentary Record in Arc Collision Zones, Geological Society of
 1139 America Special Papers 436. Geological Society of America, p. 0.

1140 Andreason, M.W., Mudford, B., Onge, J.E.S., 1997. Geologic Evolution and Petroleum
 1141 System of the Thailand Andaman Sea Basins. Proceedings of an International Conference on
 1142 Petroleum Systems of SE Asia and Australasia, Indonesian Petroleum Association, 337-350.

1143 Arai, S., 1997. Origin of podiform chromitites. *Journal of Asian Earth Sciences* 15,
 1144 303-310.

1145 Ariskin, A., Barmina, G., 2004. COMAGMAT: Development of a Magma
 1146 Crystallization Model and Its Petrological Applications. *Geochem. Int.* 42, S1–S157.

1147 Asimow, P.D., Dixon, J.E., Langmuir, C.H., 2004. A hydrous melting and fractionation
 1148 model for mid-ocean ridge basalts: Application to the Mid-Atlantic Ridge near the Azores.
 1149 *Geochemistry, Geophysics, Geosystems* 5.

1150 Audley-Charles, M.G., Carter, D.J., Barber, A.J., Norvick, M.S., Tjokrosapoetro, S.,
 1151 1979. Reinterpretation of the geology of Seram: implications for the Banda Arcs and northern
 1152 Australia. *Journal of the Geological Society* 136, 547-566.

1153 Awasthi, N., 2017. Provenance and paleo-weathering of Tertiary accretionary prism-
 1154 forearc sedimentary deposits of the Andaman Archipelago, India. *Journal of Asian Earth*
 1155 *Sciences* 150, 45-62.

1156 Bandopadhyay, P.C., 2012. Re-interpretation of the age and environment of deposition
 1157 of Paleogene turbidites in the Andaman and Nicobar Islands, Western Sunda Arc. *Journal of*
 1158 *Asian Earth Sciences* 45, 126-137.

1159 Bandopadhyay, P.C., 2017. Introduction and history of mapping and research, in:
 1160 Bandopadhyay, P.C., Carter, A. (Eds.), The Andaman–Nicobar Accretionary Ridge: Geology,
 1161 Tectonics and Hazards, Geological Society, London, Memoirs 47, pp. 1-7.

1162 Bandopadhyay, P.C., Carter, A., 2017. Mithakhari deposits, in: Bandopadhyay, P.C.,
 1163 Carter, A. (Eds.), The Andaman–Nicobar Accretionary Ridge: Geology, Tectonics and
 1164 Hazards, Geological Society, London, Memoirs 47, pp. 111-132.

1165 Bandyopadhyay, D., van Hinsbergen, D.J.J., Plunder, A., Bandopadhyay, P.C.,
 1166 Advokaat, E., Chattopadhyaya, S., Morishita, T., Ghosh, B., 2020. Andaman Ophiolite: An
 1167 Overview, in: Ray, J.S., Radhakrishna, M. (Eds.), The Andaman Islands and Adjoining
 1168 Offshore: Geology, Tectonics and Palaeoclimate. Springer International Publishing, Cham,
 1169 pp. 1-17.

1170 Barber, A.J., 2000. The origin of the Woyla Terranes in Sumatra and the Late Mesozoic
 1171 evolution of the Sundaland margin. *Journal of Asian Earth Sciences* 18, 713-738.

1172 Barber, A.J., Crow, M.J., Milsom, J.S., 2005. Sumatra: Geology, Resources and
 1173 Tectonic Evolution. Geological Society of London.

1174 Bhattacharya, A., Pal, T., Ghosh, B., 2013. Characterization of the Accreted Ophiolite
 1175 Slices of Rutland Island, Andaman Sea: Evolution in a Suprasubduction Zone Setting.
 1176 *Ophioliti* 38, 121-142.

1177 Birner, S.K., Warren, J.M., Cottrell, E., Davis, F.A., Kelley, K.A., Falloon, T.J., 2017.
 1178 Forearc Peridotites from Tonga Record Heterogeneous Oxidation of the Mantle following
 1179 Subduction Initiation. *Journal of Petrology* 58, 1755-1780.

1180 Breen, N.A., Silver, E.A., Roof, S., 1989. The Wetar Back Arc Thrust Belt, eastern
 1181 Indonesia: The effect of accretion against an irregularly shaped arc. *Tectonics* 8, 85-98.

1182 Cann, J.R., 1970. Rb, Sr, Y, Zr and Nb in some ocean floor basaltic rocks. Earth and
1183 Planetary Science Letters 10, 7-11.

1184 Chen, J., Zeng, Z., 2007. Metasomatism of the peridotites from southern Mariana fore-
1185 arc: Trace element characteristics of clinopyroxene and amphibole. Science in China Series
1186 D: Earth Sciences 50, 1005-1012.

1187 Corfu, F., 2004. U–Pb Age, Setting and Tectonic
1188 Significance of the Anorthosite–Mangerite–Charnockite–Granite Suite, Lofoten–Vesterålen,
Norway. Journal of Petrology 45, 1799-1819.

1189 Curray, J.R., 1989. The Sunda Arc: A model for oblique plate convergence.
1190 Netherlands Journal of Sea Research 24, 131-140.

1191 Curray, J.R., 2005. Tectonics and history of the Andaman Sea region. Journal of Asian
1192 Earth Sciences 25, 187-232.

1193 Curray, J.R., Moore, D.G., Lawver, L.A., Emmel, F.J., Raitt, R.W., Henry, M.,
1194 Kieckhefer, R., 1979. Tectonics of the Andaman Sea and Burma, in: Watkins, J.S.,
1195 Montadert, L., Dickerson, P.W. (Eds.), Geological and Geophysical Investigations of
1196 Continental Margins, AAPG Memoir 29. American Association of Petroleum Geologists, pp.
1197 189-198.

1198 Dewey, J.F., 1976. Ophiolite obduction. Tectonophysics 31, 93-120.

1199 Dewey, J.F., Casey, J.F., 2013. The sole of an ophiolite: the Ordovician Bay of Islands
1200 Complex, Newfoundland. Journal of the Geological Society 170, 715-722.

1201 Dilek, Y., Furnes, H., 2011. Ophiolite genesis and global tectonics: Geochemical and
1202 tectonic fingerprinting of ancient oceanic lithosphere. Geological Society of America Bulletin
1203 123, 387-411.

1204 Dilek, Y., Furnes, H., 2014. Ophiolites and Their Origins. Elements 10, 93-100.

1205 Floyd, P.A., Kelling, G., Gökçen, S.L., Gökçen, N., 1991. Geochemistry and tectonic
1206 environment of basaltic rocks from the Misis ophiolitic mélange, south Turkey. *Chemical*
1207 *Geology* 89, 263-280.

1208 Floyd, P.A., Winchester, J.A., 1975. Magma type and tectonic setting discrimination
1209 using immobile elements. *Earth and Planetary Science Letters* 27, 211-218.

1210 Frerichs, W.E., 1967. Distribution and Ecology of Foraminifera in the Sediments of the
1211 Andaman Sea. University of Southern California Los Angeles, California, p. 282.

1212 Ghiorso, M.S., Hirschmann, M.M., Reiners, P.W., Kress III, V.C., 2002. The pMELTS:
1213 A revision of MELTS for improved calculation of phase relations and major element
1214 partitioning related to partial melting of the mantle to 3 GPa. *Geochemistry Geophysics*
1215 *Geosystems* 3, 1-35.

1216 Ghosh, B., Bandyopadhyay, D., Morishita, T., 2017. Andaman–Nicobar Ophiolites,
1217 India: origin, evolution and emplacement, in: Bandyopadhyay, P.C., Carter, A. (Eds.), *The*
1218 *Andaman–Nicobar Accretionary Ridge: Geology, Tectonics and Hazards* 47. Geological
1219 Society, London, Memoirs, pp. 95-110.

1220 Ghosh, B., Bhatta, K., 2014. Podiform chromitites in Iherzolitic mantle rocks
1221 (Andaman ophiolite, India): the role of magma/rock interaction and parental melt
1222 composition. *Bulletin de la Société Géologique de France* 185, 123-130.

1223 Ghosh, B., Morishita, T., Bhatta, K., 2013. Significance of chromian spinels from the
1224 mantle sequence of the Andaman Ophiolite, India: Paleogeodynamic implications. *Lithos*
1225 164-167, 86-96. Ghosh, B., Morishita, T., Gupta, B.S., Tamura, A., Arai, S., Bandyopadhyay,
1226 D., 2014. Moho transition zone in the Cretaceous Andaman ophiolite, India: A passage from
1227 the mantle to the crust. *Lithos* 198, 117-128.

1228 Ghosh, B., Mukhopadhyay, S., Morishita, T., Tamura, A., Arai, S., Bandyopadhyay,
1229 D., Chattopadhyaya, S., Ovung, T.N., 2018. Diversity and evolution of suboceanic mantle:

1230 Constraints from Neotethyan ophiolites at the eastern margin of the Indian plate. *Journal of*
1231 *Asian Earth Sciences* 160, 67-77.

1232 Ghosh, B., Pal, T., Bhattacharya, A., Das, D., 2009. Petrogenetic implications of
1233 ophiolitic chromite from Rutland Island, Andaman—a boninitic parentage in supra-
1234 subduction setting. *Mineralogy and Petrology* 96, 59.

1235 González-Jiménez, J.M., Mondal, S.K., Ghosh, B., Griffin, W.L., O'Reilly, S.Y., 2020.
1236 Re-Os Isotope Systematics of Sulfides in Chromitites and Host Lherzolites of the Andaman
1237 Ophiolite, India. *Minerals* 10, 686.

1238 Guilmette, C., Hébert, R., Dostal, J., Indares, A., Ullrich, T., Bédard, É., Wang, C.,
1239 2012. Discovery of a dismembered metamorphic sole in the Saga ophiolitic mélange, South
1240 Tibet: Assessing an Early Cretaceous disruption of the Neo-Tethyan supra-subduction zone
1241 and consequences on basin closing. *Gondwana Research* 22, 398-414.

1242 Guilmette, C., Hébert, R., Dupuis, C., Wang, C., Li, Z., 2008. Metamorphic history and
1243 geodynamic significance of high-grade metabasites from the ophiolitic mélange beneath the
1244 Yarlung Zangbo ophiolites, Xigaze area, Tibet. *Journal of Asian Earth Sciences* 32, 423-437.

1245 Guilmette, C., Hébert, R., Wang, C., Villeneuve, M., 2009. Geochemistry and
1246 geochronology of the metamorphic sole underlying the Xigaze Ophiolite, Yarlung Zangbo
1247 Suture Zone, South Tibet. *Lithos* 112, 149-162.

1248 Guilmette, C., Smit, M.A., van Hinsbergen, D.J.J., Gürer, D., Corfu, F., Charette, B.,
1249 Maffione, M., Rabeau, O., Savard, D., 2018. Forced subduction initiation recorded in the sole
1250 and crust of the Semail Ophiolite of Oman. *Nature Geoscience* 11, 688-695.

1251 Hacker, B.R., Gnos, E., 1997. The conundrum of samail: explaining the metamorphic
1252 history. *Tectonophysics* 279, 215-226.

1253 Herzberg, C., Asimow, P.D., 2008. Petrology of some oceanic island basalts:
 1254 PRIMELT2.XLS software for primary magma calculation. *Geochemistry, Geophysics,*
 1255 *Geosystems* 9.

1256 Hutchison, C.S., 1975. Ophiolite in Southeast Asia. *GSA Bulletin* 86, 797-806.

1257 Imsong, W., Choudhury, S., Phukan, S., 2016. Ascertaining neotectonic activities in the
 1258 southern part of Shillong Plateau through geomorphic parameters and remote sensing data.
 1259 *Current Science* 110, 91-98.

1260 Irvine, T.N., Baragar, W.R.A., 1971. A Guide to the Chemical Classification of the
 1261 Common Volcanic Rocks. *Canadian Journal of Earth Sciences* 8, 523-548.

1262 Ishikawa, T., Fujisawa, S., Nagaishi, K., Masuda, T., 2005. Trace element
 1263 characteristics of the fluid liberated from amphibolite-facies slab: Inference from the
 1264 metamorphic sole beneath the Oman ophiolite and implication for boninite genesis. *Earth and*
 1265 *Planetary Science Letters* 240, 355-377.

1266 Jaffey, A.H., Flynn, K.F., Glendenin, L.E., Bentley, W.C., Essling, A.M., 1971.
 1267 Precision Measurement of Half-Lives and Specific Activities of ²³⁵U and ²³⁸U. *Physical*
 1268 *Review C* 4, 1889-1906.

1269 Jafri, S.H., 1986. Occurrence of Hagistrids in chert associated with Port Blair Seies,
 1270 South Andaman, India. *Journal of the Geological Society of India* 28, 41-43.

1271 Jafri, S.H., Balaram, V., Govil, P.K., 1993. Depositional environments of Cretaceous
 1272 radiolarian cherts from Andaman-Nicobar Islands, northeastern Indian Ocean. *Marine*
 1273 *Geology* 112, 291-301.

1274 Jafri, S.H., Balaram, V., Ramesh, S.L., 1990. Geochemistry of Andaman-Nicobar
 1275 island basalts: A case for a possible plume origin. *Journal of Volcanology and Geothermal*
 1276 *Research* 44, 339-347.

1277 Jafri, S.H., Charan, S.N., Govil, P.K., 1995. Plagiogranite from the Andaman ophiolite
1278 belt, Bay of Bengal, India. *Journal of the Geological Society* 152, 681-687.

1279 Jafri, S.H., Rao, M.V.S., Ramesh, S.L., 2006. Occurrence of ash beds in radiolarian
1280 cherts from South Andaman Island, Bay of Bengal, India: Evidence for Late Cretaceous
1281 explosive volcanism. *Current Science* 91, 1614-1615.

1282 Jafri, S.H., Sarma, D.S., Sheikh, J.M., 2010. Hyaloclastites in pillow basalts, South
1283 Andaman Island, Bay of Bengal, India. *Current Science* 99, 1825-1829.

1284 Jafri, S.H., Sheikh, J.M., 2013. Geochemistry of pillow basalts from Bompoka,
1285 Andaman–Nicobar islands, Bay of Bengal, India. *Journal of Asian Earth Sciences* 64, 27-37.

1286 Kamesh Raju, K.A., Ramprasad, T., Rao, P.S., Ramalingeswara Rao, B., Varghese, J.,
1287 2004. New insights into the tectonic evolution of the Andaman basin, northeast Indian Ocean.
1288 *Earth and Planetary Science Letters* 221, 145-162.

1289 Kamili, Z.A., Kingston, J., Achmad, Z., Wahab, A., Sosromihardjd, S., Crausaz, C.U.,
1290 1976. Contribution to the Pre–Baong Stratigraphy of North Sumatra. 5th Annual Convention
1291 Proceedings , Indonesian Petroleum Association 2, 91-108.

1292 Karunakaran, C., Pawde, M.B., Raina, V.K., Ray, K.K., 1964. Geology of the South
1293 Andaman Island, India, Reports 22nd International Geological Congress, New Delhi, India
1294 XI, pp. 79-100.

1295 Kelemen, P.B., Shimizu, N., Dunn, T., 1993. Relative depletion of niobium in some arc
1296 magmas and the continental crust: partitioning of K, Nb, La and Ce during melt/rock reaction
1297 in the upper mantle. *Earth and Planetary Science Letters* 120, 111-134.

1298 Kempton, P.D., Harmon, R.S., 1992. Oxygen isotope evidence for large-scale
1299 hybridization of the lower crust during magmatic underplating. *Geochimica et Cosmochimica*
1300 *Acta* 56, 971-986.

1301 Kimura, J.-I., Ariskin, A.A., 2014. Calculation of water-bearing primary basalt and
1302 estimation of source mantle conditions beneath arcs: PRIMACALC2 model for WINDOWS.
1303 *Geochemistry Geophysics Geosystems* 15, 1494-1514.

1304 Krogh, T.E., 1973. A low-contamination method for hydrothermal decomposition of
1305 zircon and extraction of U and Pb for isotopic age determinations. *Geochimica et*
1306 *Cosmochimica Acta* 37, 485-494.

1307 Lay, T., Kanamori, H., Ammon, C.J., Nettles, M., Ward, S.N., Aster, R.C., Beck, S.L.,
1308 Bilek, S.L., Brudzinski, M.R., Butler, R., DeShon, H.R., Ekström, G., Satake, K., Sipkin, S.,
1309 2005. The Great Sumatra-Andaman Earthquake of 26 December 2004. *Science* 308, 1127-
1310 1133.

1311 Le Bas, M.J., Le Maitre, R.W., Streckeisen, A., Zanettin, B., Rocks, I.S.o.t.S.o.I., 1986.
1312 A Chemical Classification of Volcanic Rocks Based on the Total Alkali-Silica Diagram.
1313 *Journal of Petrology* 27, 745-750. Ling, H.Y., Chandra, R., Karkare, S.G., 1996. Tectonic
1314 significance of Eocene and Cretaceous radiolaria from South Andaman Island, northeast
1315 Indian Ocean. *Island Arc* 5, 166-179.

1316 Ling, H.Y., Sharma, V., Singh, S., Mazumdar, D., Mahapatra, A.K., 1995. Cretaceous
1317 and Middle Eocene Radiolaria from Ejected Sediments of mud Volcanoes of Baratang Island
1318 in Andaman Sea of the Northeastern Indian Ocean. *Journal of Geological Society of India* 45,
1319 463-469.

1320 Ling, H.Y., Srinivasan, M.S., 1993. Significance of Eocene Radiolaria from Port Blair
1321 Group of South Andaman Island, India. *Journal of The Palaeontological Society of India* 38,
1322 1-5.

1323 Liu, C.-Z., Chung, S.-L., Wu, F.-Y., Zhang, C., Xu, Y., Wang, J.-G., Chen, Y., Guo, S.,
1324 2016a. Tethyan suturing in Southeast Asia: Zircon U-Pb and Hf-O isotopic constraints from
1325 Myanmar ophiolites. *Geology* 44, 311-314.

1326 Liu, C.-Z., Zhang, C., Xu, Y., Wang, J.-G., Chen, Y., Guo, S., Wu, F.-Y., Sein, K.,
 1327 2016b. Petrology and geochemistry of mantle peridotites from the Kalaymyo and Myitkyina
 1328 ophiolites (Myanmar): Implications for tectonic settings. *Lithos* 264, 495-508.

1329 Ludwig, K.R., 2009. Isoplot 4.1. A geochronological toolkit for Microsoft Excel.
 1330 Berkeley Geochronology Center Special Publication 4, 76.

1331 McCarthy, A. J., Jasin, B., & Haile, N. S., 2001. Middle Jurassic radiolarian chert,
 1332 Indarung, Padang District, and its implications for the tectonic evolution of western Sumatra,
 1333 Indonesia. *Journal of Asian Earth Sciences*, 19(1), 31–44.

1334 Maffione, M., Thieulot, C., van Hinsbergen, D.J.J., Morris, A., Plümper, O., Spakman,
 1335 W., 2015. Dynamics of intraoceanic subduction initiation: 1. Oceanic detachment fault
 1336 inversion and the formation of supra-subduction zone ophiolites. *Geochemistry Geophysics*
 1337 *Geosystems* 16, 1753-1770.

1338 Mattinson, J.M., 2005. Zircon U–Pb chemical abrasion (“CA-TIMS”) method:
 1339 Combined annealing and multi-step partial dissolution analysis for improved precision and
 1340 accuracy of zircon ages. *Chemical Geology* 220, 47-66.

1341 McDonough, W.F., Sun, S.s., 1995. The composition of the Earth. *Chemical Geology*
 1342 120, 223-253.

1343 Meltzner, A.J., Sieh, K., Abrams, M., Agnew, D.C., Hudnut, K.W., Avouac, J.-P.,
 1344 Natawidjaja, D.H., 2006. Uplift and subsidence associated with the great Aceh-Andaman
 1345 earthquake of 2004. *Journal of Geophysical Research: Solid Earth* 111, B02407.

1346 Morishita, T., Ghosh, B., 2010. The 26 December 2004 Sumatra-Andaman Tsunami at
 1347 the South Andaman Island: 5 years later. *Journal of the Geological Society of Japan* 116, V-
 1348 VI.

1349 Morishita, T., Yoshikawa, M., Tamura, A., Guotana, J.M., Ghosh, B., 2018a. Petrology
1350 of Peridotites and Nd-Sr Isotopic Composition of Their Clinopyroxenes from the Middle
1351 Andaman Ophiolite, India. *Minerals* 8, 410.

1352 Morishita, T., Tani, K.-I., Soda, Y., Tamura, A., Mizukami, T., Ghosh, B., 2018b. The
1353 uppermost mantle section below a remnant proto-Philippine Sea island arc: Insights from the
1354 peridotite fragments from the Daito Ridge. *American Mineralogist* 103, 1151-1160.

1355 Morley, C.K., Arboit, F., 2019. Dating the onset of motion on the Sagaing fault:
1356 Evidence from detrital zircon and titanite U-Pb geochronology from the North Minwun
1357 Basin, Myanmar. *Geology* 47, 581-585.

1358 Mukhopadhyay, M., Dasgupta, S., 1988. Deep structure and tectonics of the burmese
1359 arc: constraints from earthquake and gravity data. *Tectonophysics* 149, 299-322.

1360 Munasri & Putra, A. M., 2019. First evidence of Middle to Late Triassic radiolarians in
1361 the Garba mountains, South Sumatra, Indonesia. *Island Arc*, 28(3), e12298.

1362 Nugroho, H., Harris, R., Lestariya, A.W., Maruf, B., 2009. Plate boundary
1363 reorganization in the active Banda Arc–continent collision: Insights from new GPS
1364 measurements. *Tectonophysics* 479, 52-65.

1365 Ohara, Y., Fujioka, K., Ishii, T., Yurimoto, H., 2003. Peridotites and gabbros from the
1366 Parece Vela backarc basin: Unique tectonic window in an extinct backarc spreading ridge.
1367 *Geochemistry Geophysics Geosystems* 4.

1368 Ohara, Y., Stern, R.J., Ishii, T., Yurimoto, H., Yamazaki, T., Petrology, 2002.
1369 Peridotites from the Mariana Trough: first look at the mantle beneath an active back-arc
1370 basin. *Contributions to Mineralogy and Petrology* 143, 1-18.

1371 Pal, T., 2011. Petrology and geochemistry of the Andaman ophiolite: melt–rock
1372 interaction in a suprasubduction-zone setting. *Journal of the Geological Society* 168, 1031-
1373 1045.

1374 Pal, T., Bhattacharya, A., 2010. Greenschist-facies sub-ophiolitic metamorphic rocks of
1375 Andaman Islands, Burma–Java subduction complex. *Journal of Asian Earth Sciences* 39,
1376 804-814.

1377 Pal, T., Bhattacharya, A., 2016. Evolution of Andaman ophiolite in the backdrop of
1378 Andaman- Java subduction—a comprehensive approach integrating field and Laboratory
1379 data, Geological Survey of India Special Publication.

1380 Pal, T., Chakraborty, P.P., Gupta, T.D., Singh, C.D., 2003. Geodynamic evolution of
1381 the outer-arc–forearc belt in the Andaman Islands, the central part of the Burma–Java
1382 subduction complex. *Geological Magazine* 140, 289-307.

1383 Pal, T., Mitra, S.K., Sengupta, S., Katari, A., Bandopadhyay, P.C., Bhattacharya, A.K.,
1384 2007. Dacite–andesites of Narcondam volcano in the Andaman Sea — An imprint of magma
1385 mixing in the inner arc of the Andaman–Java subduction system. *Journal of Volcanology and*
1386 *Geothermal Research* 168, 93-113.

1387 Pal, T., Raghav, S., Bhattacharya, A., Bandopadhyay, P.C., Mitra, S.K., Renjit, M.L.,
1388 Sankar, M.S., Ghosh, B., 2010. The 2005–2006-eruption of the Barren Volcano, Andaman
1389 Sea: Evolution of basaltic magmatism in island arc setting of Andaman–Java subduction
1390 complex. *Journal of Asian Earth Sciences* 39, 12-23.

1391 Parkinson, I.J., Pearce, J.A., 1998. Peridotites from the Izu–Bonin–Mariana Forearc
1392 (ODP Leg 125): Evidence for Mantle Melting and Melt–Mantle Interaction in a Supra-
1393 Subduction Zone Setting. *Journal of Petrology* 39, 1577-1618.

1394 Parlak, O., 2016. The tauride ophiolites of Anatolia (Turkey): A review. *Journal of*
1395 *Earth Science* 27, 901-934.

1396 Pearce, J.A., 1975. Basalt geochemistry used to investigate past tectonic environments
1397 on Cyprus. *Tectonophysics* 25, 41-67.

1398 Pearce, J.A., 2008. Geochemical fingerprinting of oceanic basalts with applications to
1399 ophiolite classification and the search for Archean oceanic crust. *Lithos* 100, 14-48.

1400 Pearce, J.A., 2014. Immobile Element Fingerprinting of Ophiolites. *Elements* 10, 101-
1401 108.

1402 Pearce, J.A., Cann, J.R., 1971. Ophiolite origin investigated by discriminant analysis
1403 using Ti, Zr and Y. *Earth and Planetary Science Letters* 12, 339-349.

1404 Pearce, J.A., Dilek, Y., Newcomb, S., 2003. Supra-subduction zone ophiolites: The
1405 search for modern analogues, in: Dilek, Y., Newcomb, S. (Eds.), *Ophiolite concept and the*
1406 *evolution of geological thought* 373. Geological Society of America, pp. 269-293.

1407 Pearce, J.A., Lippard, S.J., Roberts, S., 1984. Characteristics and tectonic significance
1408 of supra-subduction zone ophiolites, in: Kokelaar, B.P., F., H.M. (Eds.), *Marginal Basin*
1409 *Geology: Volcanic and Associated Sedimentary and Tectonic Processes in Modern and*
1410 *Ancient Marginal Basins*, Geological Society, London, Special Publications 16, pp. 77-94.

1411 Pedersen, R.B., Searle, M.P., Carter, A., Bandopadhyay, P.C., 2010. U–Pb zircon age
1412 of the Andaman ophiolite: implications for the beginning of subduction beneath the
1413 Andaman–Sumatra arc. *Journal of the Geological Society* 167, 1105-1112.

1414 Plunder, A., Bandyopadhyay, D., Ganerød, M., Advokaat, E.L., Ghosh, B.,
1415 Bandopadhyay, P., van Hinsbergen, D.J.J., 2020. History of subduction polarity reversal
1416 during arc-continent collision: constraints from the Andaman Ophiolite and its metamorphic
1417 sole. *Tectonics* 39, e2019TC005762.

1418 Polachan, S., Racey, A., 1994. Stratigraphy of the Mergui Basin, Andaman Sea:
1419 Implications for Petroleum Exploration. *Journal of Petroleum Geology* 17, 373-406.

1420 Pourteau, A., Scherer, E.E., Schorn, S., Bast, R., Schmidt, A., Ebert, L., 2019. Thermal
 1421 evolution of an ancient subduction interface revealed by Lu–Hf garnet geochronology,
 1422 Halilbaşı Complex (Anatolia). *Geoscience Frontiers* 10, 127-148.

1423 Ray, J.S., Pande, K., Bhutani, R., 2015. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of subaerial lava
 1424 flows of Barren Island volcano and the deep crust beneath the Andaman Island Arc, Burma
 1425 Microplate. *Bulletin of Volcanology* 77, 57.

1426 Ray, K.K., 1985. East Coast Volcanics: a new suite in the ophiolite of Andaman
 1427 Islands. *Records of the Geological Survey of India* 116, 83-87.

1428 Ray, K.K., Sengupta, S., van Den Hul, H.J., 1988. Chemical characters of volcanic
 1429 rocks from Andaman ophiolite, India. *Journal of the Geological Society* 145, 393-400.

1430 Reagan, M.K., Ishizuka, O., Stern, R.J., Kelley, K.A., Ohara, Y., Blichert-Toft, J.,
 1431 Bloomer, S.H., Cash, J., Fryer, P., Hanan, B.B., Hickey-Vargas, R., Ishii, T., Kimura, J.-I.,
 1432 Peate, D.W., Rowe, M.C., Woods, M., 2010. Fore-arc basalts and subduction initiation in the
 1433 Izu-Bonin-Mariana system. *Geochemistry, Geophysics, Geosystems* 11.

1434 Rodolfo, K.S., 1969a. Bathymetry and Marine Geology of the Andaman Basin, and
 1435 Tectonic Implications for Southeast Asia. *GSA Bulletin* 80, 1203-1230.

1436 Rodolfo, K.S., 1969b. Sediments of the Andaman Basin, northeastern Indian Ocean.
 1437 *Marine Geology* 7, 371-402.

1438 Ross, P.-S., Bédard, J.H., 2009. Magmatic affinity of modern and ancient subalkaline
 1439 volcanic rocks determined from trace-element discriminant diagrams. *Canadian Journal of*
 1440 *Earth Sciences* 46, 823-839.

1441 Roy, D.K., Acharyya, S.K., Ray, K.K., Lahiri, T.C., Sen, M.K., 1988. Nature of
 1442 occurrence and depositional environment of the oceanic pelagic sediments associated with
 1443 the ophiolite assemblage, South Andaman Island. *Indian Minerals* 42, 31-56.

1444 Saccani, E., 2015. A new method of discriminating different types of post-Archean
1445 ophiolitic basalts and their tectonic significance using Th-Nb and Ce-Dy-Yb systematics.
1446 *Geoscience Frontiers* 6, 481-501.

1447 Sachin, R., Verma, S., Pal, T., 2017. Petrochemical and petrotectonic characterisation
1448 of ophiolitic volcanics from Great Nicobar island Andaman-Sumatra belt. *Journal of the*
1449 *Geological Society of India* 90, 85-92.

1450 Saha, A., Dhang, A., Ray, J., Chakraborty, S., Moecher, D., 2010. Complete
1451 preservation of ophiolite suite from south Andaman, India: A mineral-chemical perspective.
1452 *Journal of Earth System Science* 119, 365-381.

1453 Sarma, D.S., Jafri, S.H., Fletcher, I.R., McNaughton, N.J., 2010. Constraints on the
1454 Tectonic Setting of the Andaman Ophiolites, Bay of Bengal, India, from SHRIMP U-Pb
1455 Zircon Geochronology of Plagiogranite. *The Journal of Geology* 118, 691-697.

1456 Saunders, A., Tarney, J., 1991. Back-arc basins, in: Floyd, P.A. (Ed.), *Oceanic Basalts*.
1457 Springer US, Boston, MA, pp. 219-263.

1458 Schärer, U., 1984. The effect of initial²³⁰Th disequilibrium on young U-Pb ages: the
1459 Makalu case, Himalaya. *Earth and Planetary Science Letters* 67, 191-204.

1460 Schilling, J.G., Zajac, M., Evans, R., Johnston, T., White, W., Devine, J.D., Kingsley,
1461 R., 1983. Petrologic and geochemical variations along the Mid-Atlantic Ridge from 29
1462 degrees N to 73 degrees N. *American Journal of Science* 283, 510-586.

1463 Sengupta, S., Ray, K.K., Acharyya, S.K., de Smeth, J.B., 1990. Nature of ophiolite
1464 occurrences along the eastern margin of the Indian plate and their tectonic significance.
1465 *Geology* 18, 439-442.

1466 Shervais, J.W., 1982. Ti-V plots and the petrogenesis of modern and ophiolitic lavas.
1467 *Earth and Planetary Science Letters* 59, 101-118.

1468 Shervais, J.W., 2001. Birth, death, and resurrection: The life cycle of suprasubduction
 1469 zone ophiolites. *Geochemistry Geophysics Geosystems* 2, 2000GC000080.

1470 Sheth, H., 2014. What drives centuries-long polygenetic scoria cone activity at Barren
 1471 Island volcano? *Journal of Volcanology and Geothermal Research* 289, 64-80.

1472 Singh, A.K., Chung, S.-L., Bikramaditya, R.K., Lee, H.Y., 2017. New U–Pb zircon
 1473 ages of plagiogranites from the Nagaland–Manipur Ophiolites, Indo-Myanmar Orogenic Belt,
 1474 NE India. *Journal of the Geological Society* 174, 170-179.

1475 Smith, P.M., Asimow, P.D., 2005. AdiaBAT_1ph: A new public front-end to the
 1476 MELTS, pMELTS, and pHMELTS models. *Geochemistry, Geophysics, Geosystems* 6,
 1477 0.1029/2004GC000816.

1478 Soret, M., Agard, P., Dubacq, B., Plunder, A., Yamato, P., 2017. Petrological evidence
 1479 for stepwise accretion of metamorphic soles during subduction infancy (Semail ophiolite,
 1480 Oman and UAE). *Journal of Metamorphic Geology* 35, 1051-1080.

1481 Srivastava, R.K., Chandra, R., Shastry, A., 2004. High-Ti type N-MORB parentage of
 1482 basalts from the south Andaman ophiolite suite, India. *Journal of Earth System Science* 113,
 1483 605-618.

1484 Stacey, J.S., Kramers, J.D., 1975. Approximation of terrestrial lead isotope evolution
 1485 by a two-stage model. *Earth and Planetary Science Letters* 26, 207-221.

1486 Stern, R.J., Bloomer, S.H., 1992. Subduction zone infancy: Examples from the Eocene
 1487 Izu-Bonin-Mariana and Jurassic California arcs. *Geological Society of America Bulletin* 104,
 1488 1621-1636.

1489 Stern, R.J., Reagan, M., Ishizuka, O., Ohara, Y., Whattam, S., 2012. To understand
 1490 subduction initiation, study forearc crust: To understand forearc crust, study ophiolites.
 1491 *Lithosphere* 4, 469-483.

1492 Stolz, A.J., Jochum, K.P., Spettel, B., Hofmann, A.W., 1996. Fluid- and melt-related
1493 enrichment in the subarc mantle: Evidence from Nb/Ta variations in island-arc basalts.
1494 *Geology* 24, 587-590.

1495 Sun, S.-s., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic
1496 basalts: implications for mantle composition and processes, Geological Society, London,
1497 Special Publications, in: Saunders, A.D., Norry, M.J. (Eds.), *Magmatism in the Ocean Basins*
1498 42, pp. 313-345.

1499 Tate, G.W., McQuarrie, N., Tiranda, H., van Hinsbergen, D.J.J., Harris, R.,
1500 Zachariasse, W.J., Fellin, M.G., Reiners, P.W., Willett, S.D., 2017. Reconciling regional
1501 continuity with local variability in structure, uplift and exhumation of the Timor orogen.
1502 *Gondwana Research* 49, 364-386.

1503 Tate, G.W., McQuarrie, N., van Hinsbergen, D.J.J., Bakker, R.R., Harris, R., Jiang, H.,
1504 2015. Australia going down under: Quantifying continental subduction during arc-continent
1505 accretion in Timor-Leste. *Geosphere* 11, 1860-1883.

1506 Tate, G.W., McQuarrie, N., van Hinsbergen, D.J.J., Bakker, R.R., Harris, R., Willett,
1507 S., Reiners, P.W., Fellin, M.G., Ganerød, M., Zachariasse, W.J., 2014. Resolving spatial
1508 heterogeneities in exhumation and surface uplift in Timor-Leste: Constraints on deformation
1509 processes in young orogens. *Tectonics* 33, 1089-1112.

1510 van Hinsbergen, D.J.J., Kapp, P., Dupont-Nivet, G., Lippert, P.C., DeCelles, P.G.,
1511 Torsvik, T.H., 2011. Restoration of Cenozoic deformation in Asia and the size of Greater
1512 India. *Tectonics* 30.

1513 van Hinsbergen, D.J.J., Peters, K., Maffione, M., Spakman, W., Guilmette, C.,
1514 Thieulot, C., Plümper, O., Gürer, D., Brouwer, F.M., Aldanmaz, E., Kaymakcı, N., 2015.
1515 Dynamics of intraoceanic subduction initiation: 2. Suprasubduction zone ophiolite formation

1516 and metamorphic sole exhumation in context of absolute plate motions. *Geochemistry*
1517 *Geophysics Geosystems* 16, 1771-1785.

1518 Vohra, C.P., Haldar, D., Roy, A.K.G., 1989. The Andaman–Nicobar ophiolite complex
1519 and associated mineral resources – current appraisal, in: Ghose, N.C. (Ed.), *Phanerozoic*
1520 *Ophiolites of India*, pp. 281-315.

1521 Wajzer, M.R., Barber, A.J., Hidayat, S., Suharsono, 1991. Accretion, collision and
1522 strike-slip faulting: the Woyla group as a key to the tectonic evolution of North Sumatra.
1523 *Journal of Southeast Asian Earth Sciences* 6, 447-461.

1524 Wakabayashi, J., Dilek, Y., 2003. What constitutes ‘emplacement’ of an ophiolite?:
1525 Mechanisms and relationship to subduction initiation and formation of metamorphic soles, in:
1526 Dilek, Y., Robinson, P.T. (Eds.), *Ophiolites in Earth History*, Geological Society, London,
1527 *Special Publications* 218, pp. 427-447.

1528 Warren, J.M., Shimizu, N., Sakaguchi, C., Dick, H.J.B., Nakamura, E., 2009. An
1529 assessment of upper mantle heterogeneity based on abyssal peridotite isotopic compositions.
1530 114.

1531 Westerweel, J., Roperch, P., Licht, A., Dupont-Nivet, G., Win, Z., Poblete, F., Ruffet,
1532 G., Swe, H.H., Thi, M.K., Aung, D.W., 2019. Burma Terrane part of the Trans-Tethyan arc
1533 during collision with India according to palaeomagnetic data. *Nature Geoscience* 12, 863-
1534 868.

1535 Whattam, S.A., Stern, R.J., *Petrology*, 2011. The ‘subduction initiation rule’: a key for
1536 linking ophiolites, intra-oceanic forearcs, and subduction initiation. *Contributions to*
1537 *Mineralogy and Petrology* 162, 1031-1045.

1538 Wilson, M., 1989. Chapter 6 - Island arcs in *Igneous Petrogenesis*, in *Igneous*
1539 *Petrogenesis* Springer Netherlands, 153-190.

1540 Workman, R.K., Hart, S.R., 2005. Major and trace element composition of the depleted
 1541 MORB mantle (DMM). *Earth and Planetary Science Letters* 231, 53-72.

1542 Zhang, J., Xiao, W., Windley, B. F., Wakabayashi, J., Cai, F., Sein, K., Wu, H., &
 1543 Naing, S. 2018. Multiple alternating forearc-and backarc-ward migration of magmatism in
 1544 the Indo-Myanmar Orogenic Belt since the Jurassic: Documentation of the orogenic
 1545 architecture of eastern Neotethys in SE Asia. *Earth-Science Reviews*, 185, 704–731.

1546 Zhou, M.F., Robinson, P.T., Bai, W.J., 1994. Formation of podiform chromitites by
 1547 melt/rock interaction in the upper mantle. *Mineralium Deposita* 29, 98-101.

Figure 1

Figure 1

Figure 2

Figure 2

Figure 3

Figure 3

Figure 4

Figure 4

Figure 5

Figure 5

Figure 6

Figure 6

Figure 7

Figure 7

Figure 8

Figure 8 (cont)

Figure 8

Figure 9

Figure 9

Figure 10

Figure 10

Figure 11

Table 1
Click here to download Table: Table 1.xlsx

Table 1. Geographic location and petrographic summary of the studied samples of Andaman ophiolites							
Sample	Coordinate		Island	Rock type	Petrographic feature	Geochemistry	Geochronology
148A	11°24'10"N	92°39'34"E	Rutland Island	Plagiogranite	Dominantly composed of plagioclase (often saussuritized), amphibole and quartz. Amphibole is replaced by chlorite and quartz is often recrystallized. Intergranular spaces between larger plagioclases are composed of granular aggregate of smaller plagioclase and quartz. (Fig. 4a)		U-Pb Zircon, this study
BD-1	11°34'01"N	92°44'10"E	South Andaman	Plagiogranite	Dominantly composed of plagioclase (commonly saussuritized) and quartz, with hypidiomorphic granular texture as described by Jafri et al. (1995). Minor amounts of amphibole, epidote and chlorite are also present (Fig. 4b). Smaller plagioclase grains appear to be floating between relatively larger quartz grains. Interstitial vermicular and micrographic intergrowths of quartz and plagioclase feldspar are also noted.	Presumably equivalent with plagiogranites as reported by Jafri et al. (1995)	Presumably equivalent with U-Pb Zircon, Sarma et al. (2010)
SA-GAB1	11°31'57"N	92°43'25"E	South Andaman	Layered gabbro	Interlayered gabbroic rock composed of olivine, plagioclase, clinopyroxene and minor amount of opaque minerals with a cumulate texture. Modal variation in plagioclase and olivine demarcate the layering (Fig. 4c). Two types of plagioclase grains - fresh and relatively altered (saussuritized) might be indicative of melt impregnation, as found in olivine rich troctolite in a nearby location (Ghosh et al., 2014). Amphibole is present along the olivine and plagioclase interface.		U-Pb Zircon, this study [Samples SA-GAB1 and SA-GAB2, occur as boulders on the coast near location 2 (Kodiaghat section, south Andaman), where exactly similar lithology is reported in situ (Ghosh et al., 2014). This confirms that boulder samples like SA-GAB1 are not out of context hence we consider them to be representative of in-situ samples.]
SA-GAB2	11°31'57"N	92°43'25"E	South Andaman	Cumulate interlayered rock (clinopyroxenite layers are surrounded by wehrlite on either side)	Clinopyroxene is almost exclusively present in the clinopyroxenite layer (Fig. 4d). Olivine grains are mostly altered (chlorite?) and almost absent in the clinopyroxenite layer and dominated in wehrlite layers. Minor proportion of plagioclase occur interstitially.		
CH-3	11°29'33"N	92°42'18"E	South Andaman	Plagiogranite clast	Plagiogranite clast, collected from the agglomerate, with hypidiomorphic granular texture. Composed of subequal proportions of quartz and plagioclase with minor proportions of epidote (Fig. 4e).		Presumably equivalent with U-Pb Zircon, Pedersen et al. (2010)
CH-4	11°29'33"N	92°42'18"E	South Andaman	Pyroclastic (agglomerate)	Pyroclastic rock made of multiple rock fragments within an andesitic matrix composed of fine-grained plagioclase laths, epidote and a minor amount of pyroxene. The clasts include epidote rich clasts (epidosites?), andesite-dacite clasts (finer grained than the matrix) and leucocratic tonalite-plagiogranite clasts (Fig. 4f,g).		
AN1701	11°29'35"N	92°42'21"E	South Andaman	Pyroclastic (agglomerate)	Pyroclastic rock made of multiple litho fragments in a volcanic matrix. Amongst the fragments we have documented (i) a coarse grained plutonic quartzo-feldspathic rocks (plagiogranite) composed of saussuritized plagioclase, rare intergranular pyroxenes, and late stage epidote with a few opaque minerals, zircon and titanite. Some plagiogranite clasts show graphic intergrowth of quartz and feldspar; (ii) a coarse-grained plagioclase amphibole (along with minor opaque) bearing plutonic rock that could be a former gabbro/dolerite? (iii) a fine grained plagioclase and pyroxene bearing rock which dominates the clast proportion. We found chlorite to replace the former volcanic matrix suggesting some clasts could be andesite; (iv) large clinopyroxene fragment among other ophiolitic lithotypes (Fig. 4h,i).		
AN1719a	11°29'35"N	92°42'21"E	South Andaman	Tuff	Dacitic tuff (?) containing various litho-fragments including coexisting quartz and plagioclase. The plagioclase needles are aligned in places. We also note the presence of minor epidote, carbonate and opaque minerals.		
AN1720	11°29'43"N	92°42'18"E	South Andaman	Rhyolitic tuff	Volcanic rock with a fine-grained matrix containing rounded quartz, plagioclase and aggregates of epidote.		
AN1721a	11°29'59"N	92°42'18"E	South Andaman	Andesite	andesitic flow (?). This sample is made of fragments of variable size and a sub-volcanic andesitic nature floating in an isotropic brownish matrix.	this study	
AN1721b	11°29'59"N	92°42'18"E	South Andaman	Brecciated rhyodacite	The matrix is very fine grained, mainly made of small needles of plagioclase. Phenocrysts of quartz and plagioclase along with fragment of plagiogranite, rare epidote and chlorite to growth late (Fig. 4j).	this study	
AN1722b	11°29'59"N	92°42'18"E	South Andaman	Andesite	Volcanic rock made of plagioclase needles intergrown with amphibole almost without matrix. The texture suggests initial brecciation, possibly due to late stage hydrothermalism (Fig. 4k).	this study	
L-18	12°34'45"N	92°57'41"E	Middle Andaman	Leucocratic rock belonging to the diorite – granodiorite series	Leucocratic rock shows a hypidiomorphic granular texture and is dominantly composed of plagioclase and amphibole. From the textural perspective amphibole appears to be primary and sometimes replaced by chlorite (Fig. 4l).		U-Pb Zircon, this study
AN1703	12°35'44"N	92°57'27"E	Middle Andaman	Meta-chert (?)	Identified as representative of the metamorphic sole (sample AN1709a,b from in-situ exposure of this metamorphic sole rock). Detailed petrographic descriptions of these samples are given in Plunder et al. (2020). Sample AN1703 is a quartz rich rock showing a mild foliation marked by green amphibole. Sample AN1713a is a mafic rock with a nematoblastic texture. It is mainly made of amphibole and plagioclase.	this study	
AN1704a	12°35'27"N	92°57'28"E		Amphibolite (meta-volcanics)		this study	Ar-Ar Amphibole, Plunder et al. (2020)
AN1709a	12°33'44"N	92°58'07"E				this study	Ar-Ar Amphibole, Plunder et al. (2020)
AN1709b	12°33'44"N	92°58'07"E				this study	
AN1713a	12°35'16"N	92°57'34"E				this study	
AN1705a	12°32'26"N	92°58'31"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock, with altered plagioclase (sometimes with zoning pattern still visible) and chlorite rich pseudomorphs after clinopyroxene. The matrix is made of very fine-grained minerals that we identified as chlorite under the microscope and that replace a former glassy matrix. (Fig. 4m).	this study	
AN1705b	12°32'26"N	92°58'31"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock mainly made of plagioclase and pyroxene (now altered). In the former pyroxene we observed fine grained phyllosilicate that could be either chlorite or serpentine. Almost no matrix is present in between the grains. We also observed a few fragments of exotic rocks containing carbonate, quartz and serpentinite, possibly opicalcite.	this study	
AN1705c	12°32'26"N	92°58'31"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock with plagioclase phenocrysts, and some grains totally replaced by a mixture of quartz and carbonate. We interpret them to be former pyroxene due to the presence of a former longitudinal cleavages and the shape. The matrix is very fine grained, made of small plagioclase crystals and a former glassy matrix. Some opaque minerals complete the assemblage.	this study	
AN1710a	12°34'21"N	92°57'50"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock consists of plagioclase phenocrysts, and some pseudomorphs in a fine-grained matrix. As in sample AN1705c we interpret these minerals as pseudomorphs after pyroxene (Fig. 4n).	this study	
AN1710c	12°34'21"N	92°57'50"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock with a fine-grained matrix that is now mainly made of chlorite and opaque mineral. The rock is highly oxidized and contains phenocryst of plagioclase and of former pyroxene that are replaced by a mixture of carbonate and chlorite. There are also late veins of carbonates (Fig. 4o).	this study	
AN1711	12°35'03"N	92°57'42"E	Middle Andaman	Andesite-basaltic andesite	Volcanic rock made of a fine-grained ground mass. The matrix is mainly made of plagioclase and pyroxene now mostly replaced by chlorite. Porphyroblasts are plagioclase (up to 5 mm) and clinopyroxene. There are many vacuoles filled with nice rosette-shaped chlorite and opaque minerals. Late veins of carbonate crosscut the rock (Fig. 4p).	this study	

Table 2

[Click here to download Table 2.xlsx](#)

Table 2. Bulk rock geochemistry of selected samples of volcanic rocks and amphibolites from Andaman ophiolites														
Andaman (meta-volcanic)		Amphibolite (meta-volcanics)					Volcanic rock							
Sample No	AN1703	AN1704a	AN1709a	AN1709b	AN1713a	AN1705a	AN1705b	AN1705c	AN1710a	AN1710c	AN1711	AN1721a	AN1721b	AN1722b
Latitude	12°35'44"N	12°35'27"N	12°33'44"N	12°33'44"N	12°35'16"N	12°32'25"N	12°32'25"N	12°32'25"N	12°34'21"N	12°34'21"N	12°35'03"N	11°29'59"N	11°29'59"N	11°25'29"N
Longitude	92°57'27"E	92°57'28"E	92°58'07"E	92°58'07"E	92°57'34"E	92°58'37"E	92°58'37"E	92°58'37"E	92°57'50"E	92°57'50"E	92°57'42"E	92°42'18"E	92°42'18"E	92°36'28"E
Island	Middle Andaman										South Andaman			
SiO ₂	90.46	58.84	52.83	52.03	50.65	57.44	55.51	57.69	56.08	53.42	52.78	54.70	78.89	58.09
TiO ₂	0.11	0.48	1.31	1.39	1.52	0.72	0.94	0.77	1.03	0.63	0.76	0.73	0.27	1.01
Al ₂ O ₃	3.17	13.35	13.67	14.29	13.59	18.84	18.74	16.31	19.88	16.17	19.05	15.14	11.35	14.82
Fe ₂ O ₃ [†]	0.37	0.99	1.66	1.68	1.82	1.21	1.34	1.09	1.15	1.36	1.54	1.48	0.36	1.39
FeO [†]	1.90	5.06	8.45	8.55	9.29	6.16	6.86	5.58	5.84	6.91	7.86	7.56	1.85	7.09
MnO	0.06	0.16	0.20	0.19	0.26	0.13	0.12	0.20	0.04	0.36	0.18	0.13	0.02	0.16
MgO	1.12	7.18	7.83	8.08	8.26	6.91	8.81	4.20	8.07	5.37	4.79	6.93	0.39	6.16
CaO	1.34	7.04	9.09	9.05	10.03	0.52	0.54	6.10	0.58	9.13	6.39	8.51	1.18	4.88
Na ₂ O	1.21	5.07	4.05	3.69	3.18	7.33	6.25	7.46	6.98	6.25	6.29	4.58	5.58	6.08
K ₂ O	0.18	1.72	0.80	0.90	1.22	0.56	0.69	0.31	0.18	0.12	0.24	0.09	0.09	0.20
P ₂ O ₅	0.06	0.12	0.12	0.15	0.19	0.19	0.20	0.26	0.18	0.28	0.13	0.14	0.03	0.13
Total**	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Mg#	0.51	0.72	0.62	0.63	0.61	0.67	0.70	0.57	0.71	0.58	0.52	0.62	0.27	0.61
LOI	0.27	0.9	1.31	2.16	1.25	4.56	6.24	6.82	6	10.35	5.43	3.47	1.04	2.95
Total (original)	99.28	99.94	100.2	100.8	100.8	100.2	100.1	100.2	99.92	100.4	100.3	99.38	99.87	100.4
Sc	5	17	42	40	38	16	37	25	16	22	35	35	10	32
V	36	101	302	294	287	162	227	205	168	142	267	266	17	295
Cr	< 20	410	170	280	280	40	240	110	580	130	60	40	< 20	20
Co	11	31	40	41	47	23	31	16	42	22	26	26	3	27
Ni	20	280	80	100	170	50	80	50	250	80	30	30	< 20	30
Cu	10	40	50	40	90	80	100	30	90	40	90	40	10	30
Zn	< 30	80	80	90	90	80	70	60	160	40	90	< 30	< 30	< 30
Ga	3	11	15	16	13	13	15	12	17	10	12	15	10	13
Rb	3	24	9	14	15	9	10	4	4	3	3	< 1	< 1	2
Sr	45	55	164	110	223	363	289	422	162	177	436	140	101	172
Y	5.9	19.5	29.2	27.9	34.3	19.4	15.8	26	14.1	14.4	18.2	17.3	41.1	19.3
Zr	21	55	75	83	93	115	68	181	71	63	46	55	134	67
Nb	< 0.2	3.5	3.4	4.7	2.7	3.5	2	3.1	2.3	1.6	0.4	0.4	1.2	0.6
Cs	< 0.1	< 0.1	0.1	0.1	< 0.1	0.3	0.3	< 0.1	0.3	0.2	< 0.1	< 0.1	< 0.1	< 0.1
Ba	60	124	217	188	582	403	398	297	186	127	174	14	21	64
La	6.67	17.5	4.51	5.76	5.99	5.35	4.4	17.2	4.02	5.3	4.32	5.47	9.57	3.8
Ce	15.1	29.8	11.8	14.3	14.7	13.3	10.9	39.2	9.71	11.9	10.4	15	24.5	9.94
Pr	1.58	4.28	1.82	2.09	2.29	1.92	1.62	5.05	1.52	1.58	1.69	2.46	3.55	1.62
Nd	6.65	18.2	9.33	10.8	11.6	8.65	7.61	20.9	7.44	7.55	8.15	11.7	16.8	8.07
Sm	1.46	4.02	3.09	3.52	3.82	2.49	2.07	4.67	2.24	2.2	2.35	3.11	4.72	2.5
Eu	0.336	0.895	1.15	1.3	1.32	0.714	0.644	1.33	0.636	0.82	0.803	0.993	1.19	0.86
Gd	1.36	3.94	4.31	4.72	5.12	2.9	2.48	4.58	2.72	2.77	2.87	3.34	5.71	3.12
Tb	0.22	0.62	0.77	0.79	0.92	0.52	0.45	0.75	0.43	0.44	0.51	0.52	1.02	0.54
Dy	1.25	3.54	5.3	5	5.99	3.35	2.77	4.65	2.63	2.59	3.34	3.17	6.62	3.62
Ho	0.22	0.66	1.05	1.05	1.26	0.7	0.56	0.92	0.54	0.51	0.64	0.62	1.35	0.68
Er	0.6	1.92	3.12	2.94	3.64	2	1.65	2.59	1.63	1.45	1.93	1.75	4.4	1.97
Tm	0.08	0.288	0.448	0.422	0.565	0.286	0.23	0.367	0.238	0.196	0.292	0.275	0.68	0.293
Yb	0.51	1.77	2.96	2.74	3.93	1.92	1.52	2.42	1.59	1.21	2	1.76	4.74	2.02
Lu	0.08	0.264	0.445	0.429	0.618	0.304	0.244	0.374	0.255	0.185	0.305	0.259	0.749	0.31
Hf	0.5	1.6	2.1	2.1	2.2	2.5	1.8	4.1	1.8	1.6	1.3	1.6	3.6	1.8
Ta	< 0.01	0.25	0.2	0.3	0.11	0.19	0.27	0.16	0.04	0.02	< 0.01	< 0.01	< 0.01	< 0.01
Pb	< 5	6	< 5	< 5	5	< 5	6	7	< 5	< 5	< 5	< 5	< 5	6
Th	1.18	3.04	0.38	0.51	0.39	2.23	0.65	6.3	0.78	0.84	0.53	0.59	1.17	0.33
U	0.1	0.74	0.12	0.14	0.22	0.9	0.38	1.29	0.33	0.32	0.2	0.18	0.33	0.12

[†] FeO and Fe₂O₃ are adjusted as per Fe²⁺/Fe_{tot} = 0.85 (eg. Schilling et al. 1983)

** Total indicates oxides recalculated on an anhydrous basis,original total is also given.

[Click here to download Table: Table 3.xlsx](#)

Table 3. U-Pb data for zircon from Andaman samples																
Properties	Weight	U	Th/U	Pbc	206/204	207/235	2 sigma	206/238	2 sigma	rho	207/206	2 sigma	206/238	2 sigma	207/235	2 sigma
	[ug]	[ppm]		[pg]			[abs]		[abs]			[abs]	[Ma]	[abs]	[Ma]	[abs]
(a)	(b)	(b)	(c)	(d)	(e)	(f)	(f)	(f, g)	(f)	(f)	(f,g)	(f)	(f,g)	(f)	(f)	(f)
SA-GAB1: Layered gabbro, South Andaman (11°31'57"N / 92°43'25"E)																
1 - Z eu sp [1]	1	44	1.06	1.4	51	0.10259	0.01344	0.016473	0.000213	0.21	0.04517	0.00583	105.32	1.35	99.16	12.30
2 - Z tip [1]	1	137	0.60	1.0	157	0.10303	0.00314	0.015689	0.000064	0.28	0.04763	0.00141	100.35	0.41	99.57	2.89
3 - Z eu sp [1]	1	124	0.85	0.9	156	0.10475	0.00344	0.015589	0.000063	0.33	0.04873	0.00155	99.72	0.40	101.15	3.16
4 - Z eu sp [1]	2	68	0.68	0.7	210	0.10019	0.00266	0.015407	0.000051	0.37	0.04717	0.00120	98.56	0.33	96.96	2.45
148A: Plagiogranite, Rutland Island (11°24'10"N / 92°39'34"E)																
5 - Z eu sp [3]	20	68	0.72	1.4	955	0.10246	0.00060	0.015473	0.000035	0.48	0.04803	0.00025	98.98	0.22	99.05	0.56
6 - Z eu lp [6]	22	91	0.65	0.7	2739	0.10195	0.00036	0.015436	0.000034	0.71	0.04790	0.00012	98.74	0.22	98.58	0.33
7 - Z eu sp [1]	18	87	0.36	1.1	1360	0.10149	0.00044	0.015435	0.000034	0.60	0.04769	0.00017	98.74	0.22	98.15	0.41
8 - Z eu eq-sp [12]	26	91	0.64	1.5	1499	0.10221	0.00042	0.015395	0.000034	0.64	0.04815	0.00015	98.49	0.22	98.81	0.38
L18: Leucocratic rock belonging to the diorite – granodiorite series, Middle Andaman (12°34'45"N / 92°57'41"E)																
9 - Z eu+fr [9]	33	364	1.03	1.5	7145	0.09700	0.00029	0.014616	0.000032	0.84	0.04813	0.00008	93.54	0.21	94.00	0.26
10 - Z fr [11]	27	308	1.09	0.8	9831	0.09662	0.00027	0.014615	0.000031	0.83	0.04795	0.00008	93.53	0.20	93.65	0.25
11 - Z eu [8]	22	467	0.97	3.1	2994	0.09665	0.00028	0.014608	0.000031	0.81	0.04799	0.00008	93.49	0.19	93.68	0.26
12 - Z fr [1]	38	395	0.99	1.2	11202	0.09635	0.00026	0.014585	0.000031	0.89	0.04791	0.00006	93.34	0.20	93.40	0.24
a) Z = zircon; eu = euhedral; lp = long prismatic; sp = short prismatic; eq = equant; fr = fragment; [n] = number of grains in fraction; all zircon treated with chemical abrasion (Mattinson, 2005)																
b) weight and concentrations are known to better than 10%.																
c) Th/U model ratio inferred from 208/206 ratio and age of sample																
d) Pbc = total common Pb (initial + blank)																
e) raw data, corrected for fractionation and spike																
f) corrected for fractionation, spike, blank (206/204=18.59; 207/204=15.55) and initial common Pb (based on Stacey and Kramers, 1975); error calculated by propagating the main sources of uncertainty; The U-Pb ratio of the spike used for this work is adapted to 206Pb/238U = 0.015660 for the ET100 solution as obtained with the ET2535 spike at NIGL.																
g) corrected for 230Th disequilibrium according to Schärer (1984) and assuming Th/U magma = 4																