



HAL
open science

Saltwater intrusion and flow reversal at the submarine spring of a Mediterranean karst aquifer : explanation and modelling

Jean-Christophe Maréchal, B. Ladouche, Claudine Lamotte, Benoît Dewandel, Vivien Hakoun, Pierre Perrochet

► To cite this version:

Jean-Christophe Maréchal, B. Ladouche, Claudine Lamotte, Benoît Dewandel, Vivien Hakoun, et al.. Saltwater intrusion and flow reversal at the submarine spring of a Mediterranean karst aquifer : explanation and modelling. IAH Brussels 2021 Congress, Sep 2021, Bruxelles, Belgium. hal-03291111

HAL Id: hal-03291111

<https://hal-brgm.archives-ouvertes.fr/hal-03291111>

Submitted on 19 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Saltwater intrusion and flow reversal at the submarine spring of a Mediterranean karst aquifer : explanation and modelling

Maréchal JC, Ladouche B, Lamotte C, Dewandel, B, Hakoun V, Perrochet P

The Vise spring is the main outlet of a Jurassic karst aquifer located close to Montpellier city, South of France. The Vise spring is submarine, occurring at the bottom of the Thau lagoon at a depth of 30 m. The lagoon, made up of brackish water is connected to the Mediterranean sea. The fresh water from the karst aquifer as a whole, and especially from the spring, contributes to the qualitative state of the Thau lagoon which is well known for shellfish farming activities.


During the last fifty years (from 1967 to 2014), six occasional saltwater intrusions (called “inversac” in French) occurred, inverting the water flow at the submarine spring during a period varying from a few weeks to a few months. This backflooding process at the spring induces a very large saltwater intrusion into the karst aquifer. Given that this aquifer provides several highly important ecosystem services (drinking water supply for the coastal villages, fresh water to the Lagoon, thermal water to Balaruc spa and health resort), a large program of groundwater monitoring has been recently launched particularly at the submarine spring (discharge, temperature, quality) and several neighbouring borewells.

In November 2021, a seventh backflooding event started and is observed with the new monitoring system. From an initial flow rate of about 60 l/s from the aquifer to the lagoon through the spring, the flow inverted to about 350 l/s from the lagoon to the aquifer in a few minutes on the 28th November 2020 at 9:40 AM. This sudden backflooding created a sudden water level rise of about 2.5 meters into the karst confined aquifer. A few months later, the saltwater intrusion into the aquifer is still high, equal to 150 l/s.

A physical mechanism is proposed to explain the sudden inversion of flow and its long duration after that the event has started. The propagation of a piezometric wave through the aquifer is simulated using simple analytical solutions. A preliminary meshed model of the aquifer and its interactions with the lagoon has been developed for identifying the main processes. First attempts for proposing alert indicators are also discussed.

Normal conditions: upward flow

$$H_p \rho_o > H_e \rho_e$$



Inversac conditions: downward flow

$$H_p \rho_o < H_e \rho_e$$

