

Litter decomposition in peatlands is promoted by mixed plants

Fabien Leroy, Sébastien Gogo, Alexandre Buttler, Luca Bragazza, Fatima Laggoun-Défarge

▶ **To cite this version:**

Fabien Leroy, Sébastien Gogo, Alexandre Buttler, Luca Bragazza, Fatima Laggoun-Défarge. Litter decomposition in peatlands is promoted by mixed plants. Journal of Soils and Sediments, 2018, 18, pp.739-749. 10.1007/s11368-017-1820-3 . insu-01598365

HAL Id: insu-01598365
https://insu.hal.science/insu-01598365

Submitted on 2 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

SOILS, SEC # • RESEARCH ARTICLE

Litter decomposition in peatlands is promoted by mixed plants

Fabien Leroy[1,2,3] • Sébastien Gogo[1,2,3] • Alexandre Buttler[4,5,6] • Luca Bragazza[4,5,7] • Fatima Laggoun-Défarge[1,2,3]

[1]Université d'Orléans, ISTO, UMR 7327, 45071, Orléans, France

[2]CNRS/INSU, ISTO, UMR 7327, 45071 Orléans, France

[3]BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France

[4]WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Site Lausanne, Station 2, CH-1015 Lausanne, Switzerland

[5]Laboratory of Ecological Systems (ECOS), Ecole Polytechnique Fédérale de Lausanne (EPFL), School of Architecture, Civil and Environmental Engineering (ENAC), Station 2, CH-1015 Lausanne, Switzerland

[6]Laboratoire de Chrono-Environnement, UMR CNRS 6249, UFR des Sciences et Techniques, 16 route de Gray, Université de Franche-Comté, F-25030 Besançon, France

[7]Department of Life Science and Biotechnologies, University of Ferrara, Corso Ercole l d' Este 32, l-44121 Ferrara, Italy

✉ Fabien Leroy

tel: + 33 2 38 49 45 02

fabien.leroy@univ-orleans.fr

Abstract

Purpose. The carbon sink function of peatlands is primarily driven by a higher production than decomposition of the litter *Sphagnum* mosses. The observed increase of vascular plants in peatlands could alter the decomposition rate and the carbon (C) cycle through a litter mixing effect, which is still poorly studied. Here, we examine the litter mixing effect of a peat moss (*Sphagnum fallax*) and two vascular plants (*Pinus uncinata* and *Eriophorum vaginatum*) in the field and laboratory-based experiment.

Materials and methods. During the laboratory incubation, mass loss, CO_2 production and dissolved organic carbon concentration were periodically monitored during 51 days. The collected data were then processed in a C dynamics model. The calculated enzymatic activity was correlated to the measured β-glucosidase activity in the litter. In the field experiment, mass loss and CO_2 production from litter bags were annually measured for three years.

Results and discussion. Both laboratory and field experiments clearly show that the litter mixture, i.e. *Sphagnum-Pinus-Eriophorum*, had a synergistic effect on decomposition by enhancing the mass loss. Such enhanced mass loss increased the water extractable C and CO_2 production in the litter mixture during the laboratory experiment. The synergistic effect was mainly controlled by the *Sphagnum-Eriophorum* mixture that significantly enhanced both mass loss and CO_2 production. Although the β-glucosidase activity is often considered as a major driver of decomposition, mixing the litters did not cause any increase of the activity of this exo-enzyme in the laboratory experiment suggesting that other enzymes can play an important role in the observed effect.

Conclusions. Mixing litters of graminoid and *Sphagnum* species led to a synergistic effect on litter decomposition. In a context of vegetation dynamics in response to environmental change, such a mixing effect could alter the C dynamics at a larger scale. Identifying the key mechanisms responsible for the synergistic effect on litter decomposition, with a specific focus on the enzymatic activities, is crucial to better predict the capacity of peatlands to act as C sinks.

Keywords β-glucosidase • Carbon dynamics models • Catalysis • CO_2 production • Dissolved organic carbon • Litter mixture effect

1. Introduction

Sphagnum-dominated peatlands accumulate organic matter (OM) as peat at a rate of ca. 20 to 30 g C m^{-2} $year^{-1}$ (Francez 2000; Rydin et al. 2013). At global scale, peatlands are estimated to store about 270 to 547 Pg C as peat (1 Pg = 10^{15} g), representing ca. 15-30% of the world's soil carbon (C) stock in an area accounting for only 3-5% of the land surface (Turunen et al. 2002; Yu et al. 2010). This high rate of C accumulation is due to peculiar environmental and soil conditions, i.e. waterlogging, anoxia, acidity, low temperature, and specific plant species composition that ultimately hamper the microbial litter decomposition so resulting in a net accumulation of OM as peat (e.g. Gorham, 1991; Holden, 2005). In particular, *Sphagnum* mosses have a key role for peat accumulation as they are able to create unfavorable conditions for decomposer activities by producing a recalcitrant litter and by promoting waterlogged and acidic conditions (Van Breemen 1995). As a result, *Sphagnum* mosses gain in competitive ability against vascular plants, whose litter is much more easily decomposable (Bragazza et al. 2007, 2009), thereby acting as effective ecosystem engineers (Van Breemen 1995).

Human-induced environmental changes are expected to modify plant species abundance in peatlands, in particular by favoring vascular plants at the expense of *Sphagnum* mosses under a warmer climate (Buttler et al., 2015; Dieleman et al., 2015). Although previous studies have shown that warming can modify the rate of litter decomposition in peatlands (Thormann et al., 2004; Bragazza et al. 2016) and that decomposition rates of *Sphagnum* are lower than those of vascular plants (Hoorens et al., 2010), few studies have addressed the issue of a litter mixture effect in peatlands (Hoorens et al. 2010; Krab et al. 2013; Gogo et al. 2016). Most of the studies on litter decomposition in peatlands have focused on single plant species, although in natural conditions litters mainly consist of a mixture of multiple plant species (Salamanca et al. 1998). Litters in mixture can decompose faster (synergistic effect) or slower (antagonistic effect) than the same litter type alone (non-additive effect (Gartner and Cardon, 2004). Almost 70% of mixed-species litter exhibited non-additive mass loss with a prevalence of synergistic effects (Gartner and Cardon, 2004). Such modifications of the decomposition rate due to litter mixture could affect the imbalance between primary production and decomposition in peatlands in response to a vegetation dynamics and, ultimately, the capacity of peatlands to act as C sinks. Despite the crucial role of litter decomposition in controlling C dynamics and peat accumulation in peatlands, we have still little understanding of how the mixture litter affects decomposition.

As change in leaf litter quality can affect enzyme production by soil microbes (Hu et al. 2006), litter mixture could also change enzyme activities, which may result in a non-additive effect. For example, β-glucosidase is commonly described as an important exo-enzyme involved in C-cycling, and therefore primarily in the

decomposition of cellulose (Kourtev et al. 2002; Sinsabaugh et al. 2002), so that it could be a key-player in the non-additive effect of litter mixture. Nevertheless, the role of enzymes in relation to litter decomposition still needs to be elucidated.

In order to understand the effect of a litter mixture of *Sphagnum* mosses and vascular plants on OM decomposition and C dynamics in peatlands, we performed a decomposition experiment under laboratory and field conditions. Experimental laboratory data were then used to calibrate a C dynamics model proposed by Gogo et al. (2014), using three C compartments: the solid (mass loss), the dissolved (Water Extractable Organic Carbon, WEOC) and the gaseous components (CO_2-C). In this model, the litter is catalyzed by exo-enzymes at a rate "c" to produce soluble organic C, i.e. WEOC, which is used for respiration at a rate "r" by microorganisms. The model gives a catalysis rate for each litter that contributes to overall catalytic activity. The field experiment was performed to assess whether there was agreement between laboratory and field decomposition tests by focusing on mass loss and CO_2 production. Overall, the aims of this work were: (i) to determine the occurrence of a litter mixture effect for three different plant species during decomposition; (ii) to elucidate the role of β-glucosidase activity during litter decomposition; (iii) to relate the catalysis rate from the C dynamics modeling to the activity of hydrolytic enzymes.

2. Materials and methods

2.1 Study site and litter sampling

Plant litter material for laboratory and field experiments was collected at the Forbonnet peatland in Frasne (France, N46°49'35" E 6°10'20", 840m). The site is a *Sphagnum*-dominated peatland with a mean annual temperature of 7.5°C and annual rainfall amount of 1400 mm (Laggoun-Défarge et al. 2008; Delarue et al. 2011). The vascular plant cover mainly consists of *Eriophorum vaginatum*, *Scheuchzeria palustris*, *Andromeda polifolia*, *Vaccinium oxycoccos* and *Carex limosa* (Buttler et al. 2015) with a recent increased abundance of trees of the genus *Pinus* in response to a decline in peat water content, as observed in many bogs in the study region (Frelechoux et al. 2000). Overall, the moss layer is primarily dominated by *Sphagnum fallax* and *S. magellanicum*. Litter samples of *Eriophorum vaginatum* (E) and *Pinus uncinata* (P) were composed, respectively, of senescent leaves and needles. Litter samples of *Sphagnum fallax* (S) corresponded to the decaying part just below the green photosynthesizing apical part (Bragazza et al. 2007). After having been air-dried, sub-samples of each litter type were oven dried at 50°C for 48 hours in order to calculate the dry weight.

2.2 Laboratory experiment

Sample preparation and incubation - For the laboratory experiment, samples of one gram for seven litter types were prepared. Litter types were formed by single litter of *Sphagnum fallax* (S), *Eriophorum vaginatum* (E) or *Pinus uncinata* (P), and by the following mixtures: *Sphagnum fallax* + *Eriophorum vaginatum* (SE), *Sphagnum fallax* + *Pinus uncinata* (SP), *Pinus uncinata* + *Eriophorum vaginatum* (PE) and *Sphagnum fallax* + *Pinus uncinata* + *Eriophorum vaginatum* (SPE). Litter samples received 20 mL of peatland water overnight in order to inoculate microorganisms (Hoorens et al. 2002). The excess water was removed with a tissue and the litter samples were placed in aluminum cups with small holes at the bottom to allow air circulation. The cups were placed on a 0.4 L pot containing 20 mL of saturated solution of K_2SO_4 to maintain moist conditions. Each pot with its cup was covered with perforated aluminum foil to allow air circulation. All pots were placed in an incubator (Aralab 1200) at 20°C and 95% of relative humidity. A total of 105 litter samples (pots) were prepared in order to have 3 replicates for each of the 7 litter types and 5 periodical harvests. At intervals of 0, 1, 14, 28 and 51 days after incubation, we measured CO_2 production, water extractable organic carbon (WEOC) release, litter mass loss, β-glucosidase activities and C content in 3 replicats of each litter type.

Laboratory measurements - The CO_2 production was measured with a GMP343 Vaisala probe after placing the litter sample (pots) in a 2.34 L chamber. The measurement took 15 minutes. The slope of CO_2 increased over time within the chamber (in μmol CO_2. mol $air^{-1}.sec^{-1}$) and was used to calculate a cumulative C release (g C g^{-1} initial C litter). The WEOC from each sample was obtained after rinsing the incubated litter three times with 50 mL of distilled water and followed the method described in Delarue et al. (2011). The extract was filtered through a 0.45 μm membrane filter. Dissolved organic carbon (DOC) in water extract was determined with a Shimadzu TOC 5000A (Total Organic Carbon Analyzer). The particles on the 0.45 μm filter and the remaining litter were dried at 50°C during two days and then weighed to obtain the remaining mass (mass loss = initial mass – remaining mass) in g OM g^{-1} initial litter mass. The C in each litter sample was then measured with an elementary analyzer (Thermo-FLASH 2000 CHNS/O Analyzer). This normalized the three C pools in g C g^{-1} initial C litter, thus making possible to input data in the C dynamics models. The activity of β-glucosidase was measured by adding to 0.5 mL of water extracted litter, 3.5 mL of 4-Methylumbelliferyl β-D-glucopyranoside, a substrate for β-glucosidase activities which is transformed into 4-Methylumbelliferone (MUF). After one hour of incubation, the concentration of MUF was determined by fluorescence using an excitation wavelength of 330 nm and an emission wavelength of 450 nm and was compared to standard solution. Measured values obtained in the litter mixtures

were compared to calculate additive values (expected) after Hoorens et al. (2010), which are means of the values as measured in the decomposition of the single species litter type.

C dynamic modeling - The experimental design was similar to that published by Gogo et al. (2014, 2016). These authors conceptualized the model assuming that solid OM is catalyzed by exo-enzymes, leading to soluble OM, which is then absorbed by microorganisms and used as an energy source for different microbial functions (enzyme production, maintenance, growth). Then, the soluble organic C is respired and released into the environment in the form of CO_2. The applied model followed Schimel and Weintraub (2003) but was simplified to make it experimentally testable (Supplementary Fig. S6). It is composed of three compartments: (i) the "L" compartment corresponding to the C fraction contained in the litter (solid fraction), (ii) the "W" compartment corresponding to the C fraction in the WEOC (dissolved fraction) and (iii) the "G" compartment corresponding to the C fraction in the cumulative CO_2 released by microbial respiration (gaseous fraction). C flow from the L to the W compartment at a rate corresponding to the exo-enzyme catalysis rate. C flows from the W to the G compartment at a rate corresponding to the respiration rate. Equations describe the simultaneous change in time of the state variable (L, W, and G) and the reaction rates (Gogo et al. 2014; Supplementary Fig. S6). At any time, the sum of all these three fractions is equal to 1. The three fractions corresponding to C stocks (solid, soluble, gaseous) were experimentally measured. The catalysis rate "c" and the respiration rate "r" were tuned at the same time to fit the stock values of the model to those experimentally assessed. When the reaction rates are allowed to change in the course of the experiment, the goodness of fit is improved. The reaction rates were allowed to follow a negative exponential decrease with time. The following parameters describe the shape of the curve: "a+b" = the initial reaction rate, "a" = the final rate, "m" = decay rate with time of the reaction rate (Rovira and Rovira, 2010). Observed rates obtained in litter mixture were compared to calculate additive rates, which are means of the catalysis and respiration rates measured in decomposition of the single species litter type. The root mean square error (RMSE) was calculated for each litter type in order to represent model performance. The RMSE is calculated by squaring individual errors, summing them, dividing the sum by their total number (N), and then taking the square root of this quantity:

$$RMSE \sqrt{\sum \frac{(y_{pred} - y_{meas})^2}{N}}$$

The RMSE was normalized to the mean of observed data and multiplied by 100 to obtain the Normalized Root Mean Square Error (NRMSE) in percentage (%):

$$NRMSE = 100 \times \frac{RMSE}{\bar{y}}$$

2.3 Field experiment

The litterbag experiment was performed at the Forbonnet peatland in order to annually measure mass loss and CO_2 production during three years of field incubation. Litterbags (0.5 mm mesh) were prepared with one gram of litter of S, P, E and in a mixture of the three species (SPE). For the SPE litter mix, each species contributed equally to the final weight of the litter bag. Litter bags were placed vertically in the moss carpet and buried at 5 cm depth on November 2009 in lawns, which represent the major feature in this bog, avoiding hummocks where there was tree encroachment. Twelve samples were prepared for each of the four litter treatments that were periodically collected after one, two and three years of incubation. Litter mass loss was calculated as percentage of remaining mass of initial litter mass. The CO_2 flux was measured immediately after collection in a 835 cm^3 chamber with a Li-Cor LI-8100 analyser and expressed as fluxes (in $\mu g\ CO_2\ h^{-1}\ g^{-1}$ dry weight).

2.4 Statistics

The role of litter mixture in affecting the measured variables was compared to the corresponding additive calculated values (expected) from the litter decomposition of each single species. Two-way ANOVA was applied to investigate differences in the measured variables for both the field and laboratory experiments where the selected factors were "litter types" (observed vs additive) and "time (years or days) of measurements. Post-hoc Tukey's Honestly Significant Differences (HSD) tests were performed to determine significant differences within groups. Differences were considered as a trend for $p < 0.1$ (noted with: -) and significant at $p < 0.05$ and referred to by * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$. Modeled curves for temporal changes of variables in the laboratory incubation experiment were calculated from the C dynamics model. All statistical analyses were done using R 3.1.1 software.

3. Results

3.1 Laboratory experiment

During the laboratory incubation of the single litter type, *Sphagnum fallax* (S) litter was characterized by a mass loss, DOC release and CO_2 production significantly different compared to *Pinus uncinata* (P) and *Eriophorum vaginatum* (E) litter (Fig. 1). The S, E and P litters contained, respectively, 42.4, 50.9 and 47.7 % of C (Table 1). These contents were constant in time and between additive (calculated values) and measured values in mixture litter (Table 1). When compared to the decomposition calculated from the single plant litter (additive mean values), the observed *Sphagnum* + *Pinus* + *Eriophorum* (SPE) mixture showed a significantly higher mass loss (i.e. lower

198 remaining C) ($p<0.001$), a higher WEOC ($p<0.05$) as well as a higher production of CO_2 (p: 0.063) (Fig. 2). The
199 litter mixture of *Sphagnum* + *Eriophorum* (SE) significantly ($p<0.001$) increased mass loss and CO_2 production
200 by 13% after 51 days of incubation, as compared to the additive effect (Fig. 2, Table 2). The C dynamics in the SP
201 and PE mixtures did not significantly differ from the additive effect of the single species (Fig. 2, Table 2) although
202 the litter mixture of *Pinus* and *Eriophorum* (PE) showed a tendency to decrease litter mass loss and CO_2 production
203 (p: 0.069 and 0.097, respectively).
204
205 3.2 Field experiment
206 After three years of field incubation, the decomposition of single litter of *S. fallax* was slower than that of vascular
207 plants ($p<0.001$) (Fig. 3, Table 3). *S. fallax* litter showed significant differences for CO_2 production compared to
208 *Pinus uncinata* ($p<0.001$) and *Eriophorum litter* ($p <0.05$). There was a significant effect of litter mixture during
209 field decomposition, with a mass loss higher (ca. 19%) than that expected from the single species additive effect
210 ($p<0.01$) (Fig. 4), while the CO_2 production in the mixture litter did not differ from the additive effect (Fig. 4,
211 Table 3).
212
213 3.3 Catalysis of C dynamics models and β-glucosidase activities
214 The model fitted well the measured values with a lower NRMSE (sum between 2.82 and 9.41 %; Supplementary
215 Table S5) compared to Gogo et al. (2014). The catalysis rate decreased with incubation time in each litter type
216 (Table 4). In *Sphagnum* litter, this rate was initially very fast and decreased rapidly to become slower than vascular
217 plant litters. The catalysis rates measured in *Sphagnum-Eriophorum* (SE) and *Sphagnum-Pinus-Eriophorum* (SPE)
218 mixtures were always greater than expected from the mean of single litter species at each time of measurement
219 (Table 4). This corresponded to higher C mass loss, WEOC and CO_2 production.
220 At the beginning of incubation, β-glucosidase activity was similar in all litter types but afterward it significantly
221 increased in *Eriophorum* and *Pinus* litters, while remaining constant in *Sphagnum* litter (Supplementary Fig. S7).
222 Mixing litters did not affect β-glucosidase activities so that there were no significant differences between observed
223 and calculated additive values (Fig. 5). Overall, β-glucosidase activities increased with incubation time, whereas
224 the catalysis rate decreased (Fig. 5; Table 4) so that the catalysis rate was inversely correlated to the β-glucosidase
225 activity (r^2: 0.52; $p<0.001$; Supplementary Fig. S8).
226
227 **4. Discussion**

4.1 Occurrence of a synergistic effect

The laboratory incubation experiment clearly showed that the litter mixture of *Sphagnum-Eriophorum* as well as *Sphagnum-Pinus-Eriophorum* had a synergistic effect on mass loss and CO_2 production compared to the corresponding additive effect. Similar results were obtained with a mixture of *Sphagnum* litter and graminoid species (Hoorens et al. 2002; Gogo et al. 2016). Such synergistic interaction on decomposition has been explained by differences in litter chemistry such as N concentration (Hoorens et al. 2002). However, similar N concentration was found in our litter types. Mixing different litters can produce both chemical diversity and microhabitat complexity so supplying an increased diversity of substrates to the decomposers (Gartner and Cardon, 2004). Also, special attention in future studies should be devoted to the improvement of microclimatic condition such as the water content of individual litter in mixture through water flow from the wettest to the driest litter. The hypothesis is that through such water flow the conditions are improved in the driest litter, without decreasing to large extent the conditions of the wettest litter (Gogo et al., 2017).

The field experiment also showed an increase of decomposition by mixing *Sphagnum* + *Eriophorum* + *Pinus* litter. *Sphagnum fallax* decomposed at a slower rate than *Pinus uncinata* and *Eriophorum vaginatum* litter as observed elsewhere by Hobbie (1996). Curiously, we expected to have higher CO_2 production in combination with higher litter mass loss, but, instead, the highest CO_2 production was measured in *Sphagnum* litter with the slowest mass loss. The high capacity of *Sphagnum* litter to maintain capillary water could have contributed to retaining water that had previously percolated through the upper photosynthetically active centimeters of the *Sphagnum* carpet. This percolating water may have been enriched in labile C that could stimulate CO_2 production. Furthermore, the capillary structure of *Sphagnum* is known to host rich microbial communities, including microbial predators, i.e. amoebae (Jassey et al., 2013) in the living apical part and we cannot exclude the possibility that this might have affected the decaying part in the long field run. This could explain the decoupling observed between mass loss and respiration rate.

Laboratory incubation was a short experiment in which environmental factors and C input and output were well controlled. Conversely, field experiments spanned over three years and thus represent litter mass loss as it occurs in natural field conditions. In both laboratory and field experiments, mixing *Sphagnum* with *Eriophorum* and *Pinus* litter had a synergistic effect on litter decomposition. Such non-additive effects have already been reported for other ecosystems (Wu et al. 2013; Zhang et al. 2014), although the reasons are still unclear. Many studies have tried to explain such an effect by nutrient exchanges between litters (Vos et al. 2013), litter chemical quality (Meier and Bowman 2010) or changes in habitat characteristics (Lecerf et al. 2011). Nevertheless, only

very few studies have addressed this effect at microbial scale and established links between microbial communities and litter decomposition (Chapman et al. 2013).

4.2 Role of β-glucosidase in early C dynamics the early decomposition stages???

Contrary to our hypothesis, no significant increases were noticed in β-glucosidase activity between the observed and the additive values during the laboratory experiment, suggesting that, at least in the early stages of decomposition (i.e. the first 28^{h} days), the synergistic mixture effect does not originate from the stimulation of this hydrolytic enzyme.

By comparing the measured enzymatic activities to the modeled catalysis rates, the *Sphagnum* litter showed values similar to those obtained by Gogo et al. (2014) and Gogo et al. (2016), with a fast rate in the early stage of decomposition, and slower but constant values thereafter (< 0.002 $gC.g^{-1}C.d^{-1}$). Contrary to our hypothesis, a negative link was established between modeled catalysis rates and β-glucosidase activities. Such a link between litter enzyme activities and decomposition rates was already assessed with hydrolases and the results showed that this relationship was weak (Allison and Vitousek, 2004). These authors suggested that mass loss occurred independently of enzyme activities with compounds that may leached out of the litter. As this last one, the really different litter types could not allowed to connect the enzymes activities with the decomposition rates. Furthermore, fast catalysis rate in the initial decomposition stage was not related to the β-glucosidase activity but maybe more to the phenol oxidase activity particularly in the case of *Sphagnum* litter (Sinsabaugh et al. 2002). Indeed, it has been shown that phenol oxidase enzymes by degrading inhibitory phenolic compounds, allow other enzymes to act on soil organic C and as a consequence regulate the stability of a vast C store in the soil following the hypothesis of 'enzyme latch' mechanism (Freeman et al., 2001).

Linking microbial extracellular enzyme production and litter decomposition modeled catalysis rates could provide information on the mechanisms of decomposition and non-additive effect (Sinsabaugh et al. 2002). However, using different litter types (which can change litter quality, nutrient availability and pH) such a correlation could not be demonstrated. Enzymatic degradation is still necessary to degrade complex and insoluble litter compound and relating modeled catalysis rates and enzymatic activities are likely to provide powerful models that remains poorly understood.

4.3 Sensitivity of the WEOC compartment

Mixing *Sphagnum* with *Pinus* litters did not increase OM decomposition but affected the WEOC compartment, i.e. the dissolved C pool in the laboratory-based experiment. This dissolved fraction described as the pool that is mostly sensitive to any modification in litter decomposition (e.g. Gogo et al. 2014) is considered as a transitory compartment between solid and gaseous pools, which contains a low C-content compared to the other pools. Thus, an increase in litter decomposition would have a strong effect on this compartment as observed for the *Sphagnum-Eriophorum-Pinus* litter mixture. The *Sphagnum-Eriophorum* mixture did not increase the WEOC content as this pool is a 'dynamic' compartment that is rapidly transformed into a gaseous C pool. The WEOC increase in *Sphagnum-Pinus* litter could indicate a modification in litter decomposition, which was still not perceived in the solid or gaseous compartments however.

4.4 Implication in a scenario of vegetation dynamics

Like Hoorens et al. (2002) and Gogo et al. (2016), our study showed that mixing graminoid and *Sphagnum* litters together induces a synergistic effect on decomposition. This has important implication for estimating litter decomposition rates in peatlands. These species-specific interactions appeared to be modulated by the changing climate through modifications of litter chemistry (Hoorens et al., 2002). In the context of global change and its role in causing a shift in the relative abundance of *Sphagnum* and vascular plants in peatlands (Dieleman et al., 2015), synergistic effects of mixed litters on decomposition is likely to play a critical role for the C dynamics at ecosystem level.

5 Conclusions

On the global scale, the C-sink capacity of *Sphagnum* peatlands could be affected by the litter mixture effect. This calls for further studies on non-additive effects of litter mixture, with a focus on elucidating the specific enzymatic mechanisms behind such interactions, with the ultimate goal of incorporating them in global biogeochemical models.

Acknowledgement This work was funded as part of the PEATWARM initiative through an ANR (French National Agency for Research) grant (ANR-07-VUL-010) and was also supported by the Labex VOLTAIRE (ANR-10-LABX-100-01). The authors gratefully acknowledge the financial support provided to the PIVOTS project by the Région Centre – Val de Loire (ARD 2020 program and CPER 2015 -2020). The authors are also indebted to the Regional Scientific Council of Natural Heritage of the Franche-Comte Region for permitting access to Le Forbonnet site. The authors would like to thank E. Rowley-Jolivet for revision of the English version.

References

Allison S, Vitousek P (2004) Extracellular enzyme activities and carbon chemistry as drivers of tropical plant litter decomposition. Biotropica 36(3): 285-296.

Bragazza L, Siffi C, Iacumin P, Gerdol R (2007) Mass loss and nutrient release during litter decay in peatland: the role of microbial adaptability to litter chemistry. Soil Biol Biochem 39(1):257-267

Bragazza L, Buttler A, Siegenthaler A, Mitchell E (2009) Plant litter decomposition and nutrient release in peatlands. In: "Carbon cycling in Northern peatlands", edited by: Baird A, Belyea L, Comas X, Reeve R, Slater L. Geophysical Monograph. 184:99-110

Bragazza L, Buttler A, Robroek BJM, Albrecht R, Zaccone C, Jassey VEJ, Signarbieux C (2016) Persistent high temperature and low precipitation reduce peat carbon accumulation. Glob Ch Biol 22, 4114–4123

Buttler A, Robroek B, Laggoun-Défarge F, Jassey V, Pochelon C, Bernard G, Delarue F, Gogo S, Mariotte P, Mitchell E, Bragazza L (2015) Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland. J Veg Sci 26(5):964-974

Chapman S, Newman G, Hart S, Schweitzer J, Koch G (2013) Leaf litter mixtures alter microbial community development: mechanisms for non-additive effects in litter decomposition. PLoS ONE 8(4):e62671

Delarue F, Laggoun-Défarge F, Buttler A, Gogo S, Jassey V, Disnar J-R (2011) Effects of short-term ecosystem experimental warming on water-extractable organic matter in an ombrotrophic *Sphagnum*-peatland. Org Geochem 42:1016-1024

Dieleman, C, Branfireun B, McLaughlin J, Lindo Z (2015) Climate change drives a shift in peatland ecosystem plant community: implications for ecosystem function and stability. Glob Change Biol 21(1):388-395

Francez A-J (2000) La dynamique du carbone dans les tourbières à *Sphagnum*, de la sphaigne à l'effet de serre. Année Biologique 39:205-270

342 Frelechoux F, Buttler A, Gillet F (2000) Dynamics of bog pine dominated mired in the Jura mountains,
343 Switzerland: a tentative scheme based on synusial phytosociology. Folia Geobotanica 35(3):273-288
344 Gartner, T. B., & Cardon, Z. G. (2004). Decomposition dynamics in mixed-species leaf litter. Oikos, 104(2), 230-
345 246.
346 Gogo S, Francez A-J, Laggoun-Défarge F, Gouélibo N, Delarue F, Lottier N (2014) Simultaneous estimation of
347 actual litter enzymatic catalysis and respiration rates with a simple model of C dynamics of *Sphagnum*-
348 Dominated peatlands. Ecosystems 17(2):302-316
349 Gogo S, Laggoun-Défarge F, Merzouki F, Mounier S, Guirimand-Dufour A, Jozja N, Hughet A, Delarue F,
350 Défarge C (2016) In situ and laboratory non-additive litter mixture effect on C dynamics of *Sphagnum rubellum*
351 and *Molinia caerulea* litters. J Soils Sediments 16(1):13-27
352 Gogo S, Leroy F, Zocatelli R, Bernard-Jannin L, Laggoun-Défarge F (2017) Role of litter decomposition
353 sensitivity to water content in non-additive litter mixture effect: theoretical demonstration and validation with
354 a peatland litter experiment. Geophysical Research Abstract 19: EGU2017-12297, EGU general assembly 2017
355 Gorham, E. (1991). Northern peatlands: role in the carbon cycle and probable responses to climatic warming.
356 Ecological applications, 1(2), 182-195.
357 Hobbie S (1996) Temperature and plant species control over litter decomposition in Alaska tundra. Ecol Monogr
358 66(4):503-522
359 Hoorens B, Stroetenga M, Aerts R (2010) Litter mixture interactions at the levels of plant functional types are
360 additive. Ecosystems 13:90-98
361 Hoorens B, Aerts R, Stroetenga M (2002) Litter quality and interactive effects in litter mixtures: more negative
362 interactions under elevated CO_2? J Ecol 90:1009-1016
363 Hu Y, Wang S, Zeng D (2006) Effects of single Chinese fir and mixed leaf litters on soil chemical, microbial
364 properties and soil enzyme activities. Plant Soil 282(1-2):379-386
365 Jassey V, Chiapusio G, Binet P, Buttler A, Laggoun-Défarge F, Delarue F, Bernard N, Mitchell E, Toussaint M-
366 L, Francez A-J, Gilbert D (2013) Above- and belowground linkages in *Sphagnum*-peatland: climate warming
367 affects plant-microbial interactions. Glob Change Biol 19:811-823
368 Kourtev P, Ehrenfeld J, Huang W (2002) Enzyme activities during litter decomposition of two exotic and two
369 native plant species in hardwood forests of New Jersey. Soil Biol Biochem 34(9):1207-1218
370 Krab E, Berg M, Aerts R, van Logtestijn R, Cornelissen J (2013) Vascular plant litter input in subarctic peat bogs
371 changes Collembola diets and decomposition patterns. Soil Biol Biochem 63:106-115

Laggoun-Défarge F, Gilbert D, Buttler A, Epron D, Francez, A-J, Grasset L, Guimbaud C, Mitchell E, Roy J-C (2008) Effects of experimental warming on carbon sink function of a temperate pristine mire: the PEATWARM project. Proceedings of 13th International Peat Society Congress, June 2008, Tullamore, Ireland, pp 599-602

Lecerf A, Marie G, Kominoski J, Leroy C, Bernadet C, Swan C (2011) Incubation time, functional litter diversity, and habitat characteristics predict litter-mixing effects on decomposition. Ecology 92:160-169

Limpens J, Berendse F, Blodau C, Canadell J, Freeman C, Holden J, Roulet N, Rydin H, Schaepman-Strub G (2008) Peatlands and the carbon cycle: from local processes to global implications - a synthesis. Biogeosciences 5:1475-1491

Meier C, Bowman W (2010) Chemical composition and diversity influence non-additive effects of litter mixtures on soil carbon and nitrogen cycling: implications for plant species loss. Soil Biol Biochem 42:1447-1454

Rovira P, Rovira R (2010) Fitting litter decomposition datasets to mathematical curves: towards a generalised exponential approach. Geoderma 155:529-543

Rydin H, Jeglum J, Jeglum, J (2013) The biology of peatlands, 2nd ed. Oxford University Press

Salamanca E F, Kaneko N, Katagiri S (1998) Effects of leaf litter mixtures on the decomposition of *Quercus serrata* and *Pinus densiflora* using field and laboratory microcosm methods. Ecol Eng 10(1):53-73

Schimel J, Weintraub, M (2003) The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. Soil Biol Biochem 35(4):549-563

Sinsabaugh R, Carreiro M, Alvarez S (2002) Enzyme and microbial dynamics of litter decomposition. Enzymes in the Environment, Activity, Ecology and Applications. Marcel Dekker, New York, Basel, pp 249-265

Strakova P, Niemi R, Freeman C, Peltoniemi K, Toberman H, Heiskanen I, Fritze H, Laiho R (2011) Litter type affects the activity of aerobic decomposers in a boreal peatland more than site nutrient and water table regimes. Biogeosciences 8:2741-2755

Thormann M, Bayley S, Currah R (2004) Microcosm tests of the effects of temperature and microbial species number on the decomposition of Carex aquatilis and Sphagnum fuscum litter from southern boreal peatlands. Canadian journal of microbiology 50(10): 793-802.

Turunen J, Tomppo E, Toloken K, Reinikainen A (2002) Estimating carbon accumulation rates of undrained mires in Finland-application to boreal and subarctic regions. The Holocene 12:69-80

Van Breemen N (1995) How *Sphagnum* bogs down other plants. Trends Ecol Evol 10:270-275

Vos V, Van Ruijven J, Berg M, Peeters E, Berendse F (2013) Leaf litter quality drives litter mixing effects through complementary resource use among detritivores. Oecologia 173:269-280

Wu D, Li T, Wan S (2013) Time and litter species composition affect litter-mixing effects on decomposition rates. Plant Soil 371(1-2):355-366

Yu Z, Loisel J, Brosseau D, Beilman D, & Hunt S (2010) Global peatland dynamics since the Last Glacial Maximum. Geophys Res Lett 37(13)

Zhang L, Wang H, Zou J, Rogers W, Siemann E (2014) Non-native plant litter enhances soil carbon dioxide emissions in an invaded annual grassland. PLoS one 9(3):e92301

Table captions

TAB. 1: C content (%), N content (%) and C:N ratio for the *Sphagnum* (S), *Pinus* (P), *Eriophorum* (E), *Sphagnum-Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus-Eriophorum* (SPE) litters.

TAB 2. Levels of significance from the post-hoc tests for comparison of *Sphagnum-Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus-Eriophorum* (SPE) litter mixture effect (additive vs observed) by incubation time on the remaining solid C, water extractable organic carbon (WEOC) and cumul of CO_2 - C production during the laboratory experiements. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

TAB 3. Levels of significance from the post-hoc tests for comparison of *Sphagnum-Pinus-Eriophorum* (SPE) litter mixture effect (additive vs observed) by incubation time on the remaining solid C and CO_2 production during the field experiements. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

TAB. 4: Catalysis rate (mg C .g^{-1}C.d^{-1}) obtained from the C-fluxes model over time for *Sphagnum* (S), *Pinus* (P), *Eriophorum* (E), *Sphagnum-Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus -Eriophorum* (SPE). The model was calibrated with data from the laboratory incubation experiment. Calculate additive rates values are means from the catalysis rates measured in the decomposition of the single species litter.

Figure captions

FIG. 1: Mean (±SE, n=3) remaining solid C (a), water extractable organic carbon (WEOC) (b) and CO_2 - C production (c) in single species litter decomposition of *Sphagnum fallax* (○, S), *Pinus uncinata* (△, P) and *Eriophorum vaginatum* (◇, E) during the laboratory incubation. Lines represent the corresponding fited curves from the model for each type of litter. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

FIG. 2: Additive (■) and observed (□) mean (±SE, n=3) of remaining solid C (i), WEOC (ii) and cumulative CO_2 - C (iii) of *Sphagnum+Pinus+Eriophorum* (a), *Sphagnum+Eriophorum* (b), *Pinus+Eriophorum* (c) and *Sphagnum+Pinus* (d) litter mixture decomposition during the laboratory incubation. Additive values are the weighed means from the values measured in the decomposition of the single species litter. Lines represent the corresponding fitted curves from the model. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01,***p < 0.001).

FIG. 3: Mean (±SE, n=12) remaining mass (a) and CO_2 production (b) of *Sphagnum fallax* (○, S), *Pinus uncinata* (△, P) and *Eriophorum vaginatum* (◇, E) litter from field litter bags experiment. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

FIG. 4: Additive (■) and observed (□) mean (±SE, n=12) of remaining litter mass (a) and CO_2 production (b) from litter mixtures of *Sphagnum*, *Pinus* and *Eriophorum* (SPE) during field litter bags experiment. Additive values were calculated as the weighed mean of the values from litter decomposition of single plant species litter. Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

FIG. 5: Additive (■) and observed (□) β-glucosidase activities of *Sphagnum+Pinus+Eriophorum*, *Sphagnum+Eriophorum*, *Pinus+Eriophorum* and *Sphagnum+Pinus* litter mixtures (±SD, n=3). Asterisks represent significant differences (NS = not significant, - p< 0.1, *p < 0.05, **p < 0.01, ***p < 0.001). Only values from 0 to 28 days of incubation are given because a contamination occurred for the β-glucosidase activities samples at 51 days.

454 **Tables**

455 Table 1.

Composition	Litters						
	S	P	E	SP	SE	PE	SPE
C (%)	42.4	50.9	47.7	46.8	45.2	49.7	46.9
N (%)	1.8	1.7	1.9	1.6	1.9	1.9	1.7
C:N	23.7	29.2	25.8	30.1	23.3	26.9	26.8

456

457 Table 2.

		Main effect		Post hoc test				
		Litter type (expected vs observed	Time (days)	0	1	14	28	51
SP	Solid C		***					
	WEOC	-						
	Cumul CO_2-C		***					
SE	Solid C	***	***					***
	WEOC							
	Cumul CO_2-C	***	***					***
PE	Solid C	-	***					
	WEOC		***					
	Cumul CO_2-C	-	***					
SPE	Solid C	*	***					
	WEOC	*	***					
	Cumul CO_2-C	-	***					

458

459 Table 3

		Main effect		Post hoc test		
		Litter type (expected vs observed	Time (years)	1	2	3
SPE	Solid C	**	***			*
	CO_2 production		***			

460

461

462 Table 4

Litters	Time (days)

		0	1	14	28	51
S		7.87	6.61	1.91	1.66	1.65
P		2.57	2.54	2.18	1.84	1.40
E		4.02	3.51	2.22	2.20	2.20
SP	observed	3.44	3.33	2.34	1.84	1.54
	additive	3.05	5.18	1.83	1.62	1.61
SE	observed	3.57	3.51	2.86	2.34	1.77
	additive	3.41	3.34	2.57	2.00	1.44
PE	observed	3.79	3.5	1.87	1.62	1.59
	additive	3.47	3.26	2.08	1.88	1.85
SPE	observed	3.29	3.23	2.63	2.11	1.46
	additive	3.10	3.05	2.45	1.93	1.31

463

Figures




Fig. 1



(i) Solid C (gC g-1 initial C litter)
(ii) WEOC (gC g-1 initial C litter)
(iii) Cumul CO2-C (gC g-1 initial C litter)
(a) Sphagnum-Pinus-Eriophorum
(b)Sphagnum-Eriophorum
(c) Pinus-Eriophorum
(d) Sphagnum-Pinus
additive
observed
***
*
NS
–
Time (days)

Fig. 2



a)
b)
S–P ***
S–E ***
P–E ***
S–P ***
S–E *
P–E NS
initial dry mass)
O2.h-1.g-1 DW)

Fig. 3



Sphagnum−Pinus−Eriophorum
**
Remaining mass (% initial dry mass)
Time(Years)
additive
observed
Sphagnum−Pinus−Eriophorum
NS
CO2 production (µg CO2 .h−1 .g−1 DW)
Time (Years)

Fig. 4



Sphagnum−Pinus−Eriophorum
NS
Sphagnum−Eriophorum
NS

Fig. 5


Supplementary data

Table


Supplementary Table S5: Percentage normalized root mean square error (% NRMSE) for the solid (L), dissolved (W), gaseous (G) pools and the sum of them for the *Sphagnum* (S), *Pinus* (P), *Eriophorum* (E), *Sphagnum-Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus -Eriophorum* (SPE) models of C dynamics.

Pools	Litters						
	S	P	E	SP	SE	PE	SPE
L	0.28	0.19	0.11	0.11	0.13	0.09	0.1
W	0.28	4.08	1.89	0.19	0.48	0.84	5.54
G	5.75	5.14	2.49	2.52	2.52	2.51	1.98
Sum	6.31	9.41	4.49	2.82	3.13	3.45	7.61

Figures




Supplementary FIG. S6: Model of the C flow in the litter decomposition process. Three compartments corresponding to the solid (litter), aqueous (WEOC), and gaseous (cumulative C-CO_2 respired) forms of C are indicated. Solid lines indicate the rates of catalysis and respiration. The L pool flows into the WEOC at the catalysis rate "c" and the WEOC is respired at the rate "r" (adapted from Gogo et al. 2014).



548

549 Supplementary FIG. S7: β-glucosidase activities in *Sphagnum* (∘), *Pinus* (∆) and *Eriophorum* (◊) litter during
550 the laboratory incubation (±SD, n=3). Only values from 0 to 28 days of incubation are given because a
551 contamination occurred for the β-glucosidase activities samples at 51 days.



552

553 Supplementary FIG. S8: Linear regression between catalysis rates and β-glucosidase activities. Each point
554 represents the β-glucosidase activities measured associated to the catalysis rate obtained for each litter type at a
555 time *t*. The r^2 was was calculated with all litters at the exception of *Sphagnum* ones (●).